






UC-NRLF



B 4 314 727



Bioscience & Natural Resources Library





FLORA

OF THE

Vicinity of San Francisco.

By

H. H. BEHR, M. D.



SAN FRANCISCO, CAL.,

1888.


QK149

B4

BIOLOGY
LIBRARY
G



Faint, illegible handwritten text, possibly a name or title.



PREFACE.

The territory to which this book applies extends from Sonoma to Santa Clara, and from Niles to the Pacific Ocean.

Since the year 1884, when the Synopsis of Genera was published, there have been added several genera, which up to that time had escaped observation. Continued investigations will lead perhaps to the discovery of some more; certainly, they will lead to the addition of species hitherto unobserved in the territory. I have followed as much as possible the admirable work of Asa Gray and Sereno Watson "Botany of California," and from their views I have deviated only when opportunities for studying the plants from specimens growing on the spot afforded to me ample grounds for a view differing from that expressed by recognized authorities.

As to many of the species characterized in this book, there exists still a considerable degree of uncertainty. Many of our Californian species split into numerous variations, which mingle frequently with variations of related, equally variable species. Some of these variations owe their existence to hybridization; and this circumstance is probably the reason why several species described and characterized by different authors have not been found

again. In annuals such spurious species will only reappear occasionally.

Questionable types can be investigated only by cultivation. Up to this time California does not possess a botanical garden or experimental grounds where such questions could be definitely settled. Therefore the solution of many problems has been left entirely to enthusiasts who utilize their leisure hours in the cultivation of doubtful species, frequently under great difficulties.

Owing to these circumstances, there is scarcely an author writing on our Flora who has not been compelled to frequently change or correct his previous views, and I hope the present author will be excused if in future times investigations shall lead him to add to, or to retract, some of his statements.

H. H. BEHR, M. D.

LINNÆAN KEY.

MONANDRIA.

I. Monogynia.

1. Acæna.—Calyx lobes valvate. Style terminal. Akene enclosed in the calyx.
2. Hippuris.—Calyx superior. Aquatic.

II. Digynia, Trigynia and Tetragynia.

1. Acæna.—Calyx lobes valvate. Akenes enclosed in the calyx. Styles of akenes terminal.
2. Callitriche.—Fruit a 4-coccous schizocarp. Aquatic.
3. Chenopodium.—Perigonium 3-5-parted. Fruit a utricle enclosed in the persistent perigonium.
4. Festuca.—Glumaceous.
5. Cyperus.—Glumaceous, trigynous.

DIANDRIA.

I. Monogynia.

A. *Floral parts incomplete, inferior.*

1. Salicornia.—Perigonium gamosepalous, opening by a slit and immersed in an excavation of the rachis.

2. Lemna.—Perigonium compressed, growing out of the margin of a thallus. Aquatic.
 3. Scirpus.—Glumaceous. Spikelets imbricate from all sides. Inferior glumæ (1-2) sterile, but not smaller than the rest. Perigonial bristles not exerted. Fruit, a nutlet, apiculate by the persistent base of the style. Aquatic.
 4. Fraxinus. — Ovary 2-celled, laterally compressed. Fruit, a samara, Tree.
 5. Acæna.—Perigonium lobes valvate. Fruit an akene enclosed in the perigonium. Style terminal.
 6. Alchemilla.—Perigonium urceolate, bracteolate, not spinose. Akene enclosed in the perigonium. Style lateral.
- B. *Floral parts complete, inferior, regular.*
1. Fraxinus.—Fruit, a samara. Tree.
 2. Lepidium.—Fruit, a silicula, laterally compressed.
- C. *Floral parts complete, inferior, irregular.*
- a. *Ovary entire.*
 1. Veronica.—Calyx herbaceous 4-parted. Ovary 2-celled. Fruit, an emarginate capsule. Leaves opposite.

2. *Synthyris*.—Calyx herbaceous, 4-parted. Ovary 2-celled.
3. *Cordylanthus*.—Calyx spathaceous, 1-sepalous. Ovary 2-celled.
4. *Utricularia*.—Ovary 1-celled, centrospermous.

b. *Ovary divided into 4 partitions.*

1. *Lycopus*.—Anthers 2-celled. Cells parallel. Aquatic.
2. *Salvia*.—Anthers 1-celled. Filament with a lateral branch, geniculate.
3. *Audibertia*.—Anthers 1-celled. Filament without lateral branch or spur.
4. *Acanthomintha*.—Stamens ascending. Anthers 2-celled. Cells divaricate. A pair of staminodia indicating the place of the upper pair of stamens.

II. *Digynia, Trigynia and Tetragynia.*

1. *Acæna*.—Calyx lobes valvate, tube spinose. Fruit, akenes enclosed in the calyx. Styles terminal.
2. *Alchemilla*.—Calyx urceolate, bracteolate, not spinose. Fruit, akenes enclosed in the calyx.
3. *Chenopodium*.—Perigonium 3-5-parted, not appendiculate. Fruit a compressed utricle, inclosed in the perigonium.
4. *Festuca*.—Glumaceous.

TRIANDRIA.**I. Monogynia.****A. *Floral parts complete, superior.***

1. Plectritis.—Limb of calyx obsolete. Corolla gamopetalous, calcarate.

B. *Floral parts complete, inferior.*

1. Montia.—Calyx 2-sepalous. Corolla cleft on one side, 5-parted.

C. *Floral parts incomplete.***a. *Perigonium petaloid.***

1. Iris.—Perigonium 6-parted, superior, alternate lobes reflexed.
2. Scoliopus.—Perigonium 6-sepalous inferior, alternate lobes reflexed.
3. Brodiaea.—Perigonium 6-parted. inferior. Stamens alternating with staminodia.

b. *Perigonium herbaceous.*

1. Acæna.—Calyx lobes valvate, tube spinose. Fruit an akene enclosed in the calyx. Style terminal.
2. Chenopodium.—Perigonium 3-5-parted, not appendiculate. Fruit a compressed utricle enclosed in the perigonium.

C. *Floral parts glumaceous.*aa. *Spikelets distichous.*

1. *Cyperus*.—Glumæ ∞ , one or the two lowermost smaller and empty.

bb. *Spikelets imbricate on all sides.*

1. *Eleocharis*.—Perigonial bristles not exerted. Caryopsis apiculate by the persistent articulate base of the style.
2. *Fimbristylis*.—Perigonial bristles not exerted (sometimes wanting). Caryopsis apiculate by the persistent continuous base of the style. Aquatic.
3. *Scirpus*.—Perigonial bristles not exerted (sometimes wanting altogether). Caryopsis apiculate by the persistent continuous base of the style. Aquatic.
4. *Eriophorum*.—Perigonial bristles exerted, considerably longer than the glumæ. Aquatic.

Observation.—All grasses which are monogynous are put in Triandria Digynia.

II. *Digynia.*A. *Flowers glumaceous.*

- a. *Spikelets single 1-flowered, imbedded in the notches of the rachis, each covered by its glume.*

I. *Lepturus*.

- b. *Spikelets all sessile on alternating notches of the rachis.*
1. *Lolium*.—Terminal spikelet with 2 glumes, lateral spikelets but one glume.
 2. *Hordeum*.—Spikelets 3 at each notch of the rachis, 1-flowered. Fruit a caryopsis wrapped and grown together with the paleæ.
 3. *Elymus*.—Spikelets 3 at each notch of the rachis, 2— ∞ -flowered.
 4. *Gymnostichum*.—Spikelets 1–3 at each notch of the rachis. Glumæ rudimentary or 0.
 5. *Triticum*.—Spikelet 1 at each notch of the rachis. Glumæ 2, placed right and left (not dorsal and ventral.)
- c. *Spikelets pedicillate, 1–3-flowered, but only 1 flower regularly developed.*
- aa. *Spikelets dorsally compressed.*
1. *Panicum*.—Glumæ 3, the lowermost very small. Involucre 0.
 2. *Setaria*.—Glumæ 3, the lowermost small; bristly involucre below the spikelet.
- bb. *Spikelets laterally compressed. Glumæ 3. The two inferior florets rudimentary, terminal floret only developed.*

1. *Phalaris*.—The 2 inferior florets reduced to 2 awnless paleæ.
 2. *Anthoxanthum*.—The two inferior florets reduced to 2 aristate paleæ larger than the developed terminal floret.
- cc. *Spikelets not dorsally compressed. Glumæ*
2. *When a second floret is present it is rudimentary and terminal.*
- aaa. *Stigma filamentose, exerted on the apex of the spikelet.*
1. *Alopecurus*.—Floret consisting of a single palea.
 2. *Phleum*.—Glumæ carinate, longer than the 2 paleæ of the floret.
 3. *Spartina*.—Glumæ carinate, the inferior shorter than the floret. Paleæ unequal, the superior longer.
- bbb. *Stigma on an elongated style, aspergilli-form and exerted below the apex of the floret.*
1. *Cynodon*.—Glumæ small, spreading. Superior palea linear, concave; inferior palea laterally compressed, oval, chartaceous.
- ccc. *Stigma plumose, exerted from the base of the floret. Style very short.*
1. *Polypogon*.—Glumæ obtuse, but aristate. Paleæ membranaceous.

2. *Agrostis*.—Glumæ acute, the inferior one larger. Paleæ membranaceous.
 3. *Calamagrostis*.—Glumæ acute, the inferior one larger. Paleæ surrounded by silky hair at the base.
 4. *Stipa*.—Glumæ acute, sometimes aristate. Paleæ cartilaginous, the inferior one cylindrically involute. Awn strong, articulate at the base, persistent, developed from the apex of the palea.
 5. *Gastridium*.—Glumæ acute, very much compressed, ventricose at their base. Paleæ membranaceous.
- d. *Spikelets pedicillate, 2-∞-flowered, the uppermost floret frequently rudimentary.*
- aa. *Stigma aspergilliform, exerted below the apex of the floret.*
1. *Phragmites*.—Spikelets ∞-flowered, lowermost floret ♂ or neuter, the other florets ♀ and pilose.
 2. *Hierochloa*.—Spikelets 3-flowered, the two inferior ones ♂ and triandrous, the terminal one ♀ and diandrous.
- bb. *Stigma plumose, exerted near the base of the floret.*
1. *Arrhenatherum*.—Spikelets 2-flowered; inferior floret ♂, aristate; superior floret ♀, awnless.

2. *Holcus*.—Spikelets 2-flowered, inferior floret ♀, awnless; superior floret ♂, aristate.
3. *Aira*.—Spikelets 2-flowered, both ♀; inferior palea 4-dentate at its apex, aristate from its base.
4. *Avena*.—Spikelets 2-∞-flowered. Florets ♀. Inferior palea 2-dentate at its apex, aristate from its median nerve.
5. *Danthonia*.—Spikelets 2-∞-flowered. Florets ♀. Inferior palea 2-cleft, aristate from the fissure.
6. *Melica*.—Spikelets 2-∞-flowered, florets awnless, the inferior or 2 inferior, ♀; the rest rudimentary.
7. *Koeleria*.—Glumæ and paleæ membranaceous. Spikelets 2-∞-flowered. Florets lanceolate, compressed, carinate; inferior palea mucronate-aristate.
8. *Dactylis*.—Glumæ and paleæ herbaceous. Spikelets 3-∞-flowered. Florets ovate, carinate. Inferior palea asymmetric, mucronate-aristate.
9. *Poa*.—Spikelets 2-∞-flowered. Florets ovate, carinate, awnless, deciduous with the joint of the axis.
10. *Eragrostis*.—Spikelets 2-∞-flowered. Florets ovate, carinate, awnless; superior palea and axis persistent.

11. *Glyceria*. Spikelets 2, ∞ -flowered. Florets ovate, convex, awnless. Aquatic grass.
12. *Lophochlæna*. — Spikelets ∞ -flowered, compressed, Glumæ membranaceous. Inferior palea 2-lobed, aristate from the mid-nerve. Superior palea 2-carinate with a pair of dentate appendages.
13. *Briza*.—Spikelets 2- ∞ -flowered, densely distichous. Florets awnless. Inferior palea inflated, with auriculate cordate base.
14. *Festuca*.—Spikelets 2- ∞ -flowered. Florets lanceolate, convex. Superior palea ciliate. Style inserted on the apex of the ovary.
15. *Bromus*.—Spikelets ∞ -flowered. Florets lanceolate. Style inserted below the apex of the ovary on its ventral side.

B. *Flowers not glumaceous.*

1. *Pentacæna*.—Calyx 2-seriate, persistent. External sepals 3, cucullate at the apex, ending in a spine; internal sepals 2, mucronate. Fruit a utricle.

III. *Trigynia.*

1. *Tillæa*.—Calyx 3-sepalous. Corolla 3-petalous. Fruit 3 follicles, each 2-seeded. Aquatic.

2. *Lastarriæa*.—Perigonium tubular, coriaceous, 6-dentate. Ovary 1. Fruit an akene. Involucre 0.
3. *Chorizanthe*.—Involucre 1–3 flowered, coriaceous, 3–6 angular. Ovary 1. Fruit, an akene.
4. *Tricerastes* (*Datisca*).—Calyx superior. Petals 0. Anthers extrorse. Ovary 3-quetrous. Styles 3, each 2-cleft. Ovary 1-celled. Placentæ parietal.
5. *Cyperus*.—Glumaceous. Spiklets distichous.
6. *Eleocharis*.—Glumaceous. Spiklets imbricate. Perigonial bristles not exerted. Caryopsis apiculate by the persistent articulate base of the style.
7. *Fimbristylis*.—Glumaceous. Spikelets imbricate. Perigonial bristles not exerted. Caryopsis apiculate. Style articulate above its base. Aquatic.
8. *Scirpus*.—Glumaceous. Spiklets imbricate. Perigonial bristles not exerted. Caryopsis apiculate by the persistent continuous base of the style. Aquatic.
9. *Eriophorum*.—Glumaceous. Spikelets imbricate. Perigonial bristles exerted, considerably longer than the glumæ. Aquatic.

TETRANDRIA.

I. Monogynia and Digynia.

A. Flowers complete.

a. *Calyx double; the interior calyx adnate to the ovary. Corolla gamopetalous.*

1. *Dipsacus.*

b. *Calyx simple. Corolla gamopetalous, inferior.*

1. *Plantago.*—Corolla-limb regular, 4-parted. Stigma filamentose. Capsule circumscissile.

2. *Gentiana.*—Ovary 1. Placentæ 2. Style short. Stigmas 2, flat and thin. Capsule septicial. Seeds minute.

3. *Microcala (Cicendia).*—Ovary 1. Placentæ 2. Style filiform. Stigma peltate. Seeds minute.

c. *Corolla gamopetalous, superior.*

1. *Galium.*—Corolla rotate. Fruit a 2-coccos schizocarp.

d. *Corolla 4-petalous, inferior.*

1. *Euonymus.*—Petals inserted on a glandular disc. Stamens inserted on the same disc, and alternating with them. Fruit a capsule. Shrub.

2. *Ptelea*.—Stamens alternating with the petals. Ovary on a convex disc, 2-celled. Cells 2-ovulate. Fruit a 2-seeded samara. Shrub.

e. *Corolla 4-petalous, superior.*

1. *Cornus*.—Calyx 4-parted. Fruit a drupe. Tree. Shrub.
2. *Eucharidium*.—Calyx tube extended beyond the ovary, its lobes reflexed. Petals unguiculate, lobed. Fruit a capsule.
3. *Gayophytum*.—Calyx tube not extended beyond the ovary. Anthers versatile. Stamens alternating with staminodia, which are opposite to the petals. Fruit a capsule.

B. *Flowers incomplete.*

a. *Perigonium inferior.*

1. *Majanthemum*.—Perigonium corolline, 4-parted, rotate.
2. *Acæna*.—Calyx lobes valvate, tube spinose. Ovaries 1-2. Style or styles terminal. Fruit 1 or 2 akenes, enclosed in the calyx tube.
3. *Alchemilla*.—Perigonium calycine; limb 8-cleft. Stamens inserted on an annular throat (faux).

4. Sanguisorba.—Perigonium calycine; limb 4-cleft, colored. Stamens inserted on an annular throat (faux).
5. Negundo.—Perigonium very small, 4-dentate. Fruit a 2-seeded samara. Tree. Shrub.

II. Tetragynia.

1. Tillæa.—Calyx 4-sepalous. Corolla 4-petalous. Follicles 4, each 2-seeded.
2. Sagina.—Calyx 4-sepalous. Corolla 4-petalous. Capsule 4-valved. Seeds reniform.
3. Cuscuta.—Corolla gamopetalous, 4-cleft. Ovary 1; ovules few, inserted on the base of the ovary. Capsule circumscissile. Parasitic.
4. Potamogeton. — Perigonium 4-parted. Fruit, 4 sessile drupes. Aquatic.

PENTANDRIA.

I. Monogynia.

A. *Floral parts gamopetalous, inferior.*

a. *Fruit 4 nutlets, each 1-2 seeded.*

aa. *Ovary with 4 sutures finally separating into 4 nutlets.*

1. Heliotropium.—Corolla funnel-shaped, throat naked, limb plicate.

bb.—*Ovary 4-parted, nutlets ventrally adherent to the persistent style.*

1. *Echinospermum*. — Nutlets 3-quetrous, glochidiate on their edges.
2. *Cynoglossum*. — Nutlets compressed, spinose on their entire dorsal surface.
3. *Pectocarya*. — Nutlets in pairs, divergent, carinate, carina dentate.

cc. *Ovary deeply 4-parted, nutlets excavated on their ventral side inserted on a pyramidal disc.*

1. *Amsinckia*. — Nutlets coriaceous, inserted above their base. Cotyledons 2-cleft.

dd. *Ovary deeply 4-parted, nutlets inserted on a flat disc.*

1. *Krynitzkia* (*Eritrichium*). — Nutlets ventrally attached from near their base to a columnar prominence inserted on the otherwise flat disc.
2. *Plagiobothrys* [*Eritrichium*]. — Nutlets attached near the middle of their ventral face to a columnar prominence, inserted on the otherwise flat disc.

b. *Fruit a 1-celled capsule with free central placentation.*

1. *Dodecatheon*. — Corolla partitions reflexed.

2. *Anagallis*.—Corolla rotate, 5-parted, capsule circumscissile.

c. *Ovary 1-celled with 2 parietal placentæ.*

1. *Menyanthes*. — Corolla funnel-shaped. Stigma undivided. Capsule 2-valved, valves bearing the seeds in their middle; placentæ not detached from the exocarp. Aquatic.
2. *Nemophila*.—Calyx with reflexed sinuses. Stamens shorter than the corolla. Stigmas 2. Capsule 2-valved. Placentæ detached from the exocarp.
3. *Ellisia*.—Calyx without reflexed sinuses. Stamens shorter than the corolla. Stigmas 2. Capsule 2-valved. Placentæ detached from the exocarp.
4. *Romanzoffia*. — Calyx without reflexed sinuses. Corolla funnel-shaped. Stigma 1, undivided. Capsule loculicidal. Seeds ∞ .

d. *Capsule 2- ∞ -celled.*

1. *Erythræa*.—Corolla funnel-salver-shaped. Anthers finally spiral. Seeds minute.
2. *Gentiana*.—Ovary 1. Placentæ 2. Style short. Stigmas 2. Capsule septicidal. Seeds minute.

3. *Polemonium*.—Corolla rotate. Stigmas 3. Filaments inserted at equal heights on the tube of the corolla, ascending; anthers incumbent.
4. *Gilia*.—Corolla rotate—funnel-shaped. Stigmas 3. Filaments straight, inserted at equal heights; anthers incumbent.
5. *Collomia*.—Corolla salver-shaped. Stigmas 3. Filaments inserted at unequal heights and exserted.
6. *Convolvulus*.—Corolla funnel-shaped, plicate, 5-angular. Cells of ovary 2-ovulate. Calyx marcescent.
7. *Datura*.—Corolla funnel-shaped, plicate, 5-angular. Calyx transversely deciduous.
8. *Nicotiana*.—Corolla funnel-shaped, plicate, 5-angular. Calyx persistent. Cells of ovary ∞ -ovulate.
9. *Rhododendron*.—Corolla campanulate. Stamens inserted on a disc, not on the corolla. Cells of the anthers opening by pores. Shrubs.
10. *Pentstemon*.—Corolla irregular, funnel-shaped. Stamens of unequal length, one of them sterile. Capsule 2-celled ∞ -ovulate.

e. *Fruit baccate.*

1. Solanum. — Corolla rotate. Anthers opening by apical pores.

B. *Floral parts gamopetalous, superior.*

a. *Fruit capsular.*

1. Samolus. — Staminodia 5, alternating with the stamens. Ovary 1-celled.
2. Campanula. — Corolla 5-lobed. Ovary flat at its apex. Capsule turbinate dehiscent by lateral pores.
3. Heterocodon (Campanula). — Corolla 5-lobed. Ovary flat at its apex. Capsule turbinate, laterally but irregularly dehiscent.
4. Specularia. — Corolla rotate. Capsule elongated, prismatic, dehiscent laterally by valves.
5. Githopsis. — Corolla tubular. Capsule clavate, dehiscent at its apex by the falling off of the base of the style.
6. Downingia. — Corolla irregular. Filaments and anthers united into a tube.

b. *Fruit baccate.*

1. Lonicera. — Corolla irregular. Ovary 2-3-celled, ∞ -seeded.
2. Symphoricarpus. — Ovary 4-celled. Berry 2-seeded. Shrubs.

c. *Floral parts polypetalous, inferior.*aa. *Floral parts irregular.*

1. *Viola*.—Calyx 5-sepalous. Corolla 5-petalous. Capsule 3-valved, placentæ 3; each ∞ -seeded.
2. *Æsculus*.—Calyx 5-dentate. Corolla 4-5-petalous. Capsule 3-valved, 1-seeded. Tree.

bb. *Floral parts regular.*

1. *Acer*.—Fruit a 2-seeded samara. Trees. Shrubs.
2. *Vitis*.—Petals coherent at the apex, and caducous at the base, inserted on a disc. Fruit baccate.
3. *Euonymus*.—Petals 5, inserted on the calyx. Stamens alternating with the petals, inserted on a glandular disc. Fruit capsular. Shrubs.
4. *Rhamnus*.—Petals 4-5 with short claws, inserted on the calyx. Stamens opposite to the petals. Fruit a drupe with several pyrenæ. Trees. Shrubs.
5. *Ceanothus*.—Calyx and disc adnate to the base of the ovary. Stamens opposite to the petals. Petals with long claws cucullate. Fruit a capsule, 3-celled; cells 1-seeded. Capsule septicidal. Trees. Shrubs.

6. Ptelea.—Stamens alternating with the petals. Ovary on a convex disc, 2-celled; cells 2-ovulate. Fruit a 2-seeded samara
Shrubs.
7. Erodium.—Carpidia 5, verticillate round a columnar axis. Styles connate at their apex.
8. Pentacæna.—Sepals 2-seriate, persistent; external sepals 3, with cucullate apex terminating in a spine, internal sepals, mucronate. Ovary 1-ovulate. Style 2-cleft. Fruit a utricle.
9. Frankenia.—Calyx tubular, 4-5 lobed. Petals 4 or 5, unguiculate. Capsule 1-celled, dehiscent by valves. Ovules few, inserted on parietal placentæ.

d. *Floral parts polypetalous, superior.*

1. Ribes.—Petals unguiculate and with the stamens inserted on the calyx limb. Fruit baccate, ∞ -seeded. Shrubs.

e. *Floral parts incomplete.*

1. Glaux.—Perigonium campanulate, 5-lobed, petaloid. Ovary 5-valved.
2. Polygonum.—Perigonium 5-sepalous, persistent. Stigmas 2-3, capitulate. Fruit an akene enclosed in the persistent perigonium.

3. *Acæna*.—Calyx lobes valvate, tube spinose. Style terminal. Akene enclosed in the calyx tube.
4. *Negundo*.—Perigonium small, 5-dentate. Fruit a samara. Tree.
5. *Abronia*.—Flowers capitulate. Involucre ∞ -phyllous. Perigonium gamosepalous, salver-shaped.

II. *Digynia*.

A. *Flowers incomplete*.

1. *Suæda* (*Schoberia*). — Perigonium 5-parted, not appendiculate. Embryo spiral.
2. *Chenopodium*.— Perigonium 5-parted, not appendiculate. Embryo annular.
3. *Acæna*.—Calyx lobes valvate, tube spinose. Fruit, 2 akenes, enclosed in the calyx tube.

B. *Flowers complete, gamopetalous; floral parts inferior*.

a. *Ovaries 2. Seeds ∞* .

1. *Apocynum*.—Ovaries distinct at the base but united by their stigma. Five scales in the tube of the corolla opposite to the corolla lobes.

b. *Ovary 1, with two opposite placentæ.*

1. *Gentiana*.—Lobes of the corolla without nectaria. Capsule ∞ -seeded. Seeds minute.
2. *Nemophila*.—Calyx with reflexed sinuses. Stamens shorter than the corolla. Capsule 2-valved. Placentæ detached from the exocarp.
3. *Ellisia*.—Calyx without reflexed sinuses. Stamens shorter than the corolla. Capsule 2-valved. Placentæ detached from the exocarp.
4. *Phacelia*.—Corolla deciduous. Stamens exserted, equal. Capsule loculicidal.
5. *Emmenanthe*.—Corolla persistent. Stamens exserted, equal. Capsule loculicidal.
6. *Eriodictyon*.—Capsule 4-valved (loculicidal and septicidal). Shrub.

c. *Ovary 1. Seeds few. Placentæ on the base of the ovary.*

1. *Cuscuta*.—Capsule circumscissile. Parasite.

C. *Corolla 5-petaled. Floral parts inferior.*

1. *Heuchera*.—Ovary 1-celled. Capsule ∞ -ovulate, dehiscent by valves. Pla-



(27)

centæ parietal at the margins of the valves.

2. Pentacæna.—Calyx 2-seriate. Petals minute. Ovary 1-ovulate. Fruit a utricule.

D. *Corolla 5-petaled. Floral parts superior.*

a. *Flowers in a simple umbel.*

1. Panax.—Fruit baccate.
2. Hydrocotyle. — Petals entire, acute. Akenes laterally compressed. Costæ filiform.
3. Bowlesia.—Petals obtuse. Akenes ovate, turgid, without costæ, pubescent.
4. Eryngium. — Petals erect, connivent, emarginate at their apex. Akenes terete, obovate, without costæ but squamate and tuberculate.
5. Sanicula. Petals erect, connivent, emarginate at their apex. Akenes spinose.
6. Panax. Fruit baccate.

b. *Flowers in a compound umbel.*

aa. *Endosperm not excavated on its ventral surface (Orthospermæ). Costæ 5, filiform; secondary costæ 0. Akenes laterally compressed.*

aaa. *Petals entire.*

1. Apium. — Petals stellate, rounded, with involute apex.

- bbb. *Petals obcordate with a small inflexed lobe.*
Calyx limb obliterated.
1. Carum.—Petals regular. Styles reclined. Akenes oblong. Costæ filiform. Oil tubes solitary.
 2. Pimpinella.—Petals regular. Styles thin, divergent. Costæ filiform. Oil tubes 3 in each vallecule.
- ccc. *Petals obcordate with a small inflexed lobe.*
Calyx limb 5-dentate.
1. Cicuta.—Endosperm convex. Fruit contracted. Oil tubes solitary.
 2. Sium.—Fruit contracted. Oil tubes superficial, 3 in each vallecule.
 3. Berula.—Fruit contracted. Oil tubes covered by a thickened pericarp, 3 in each vallecule.
- bb. *Endosperm not excavated on its ventral surface (Orthospermæ). Costæ 5, secondary costæ 0. Akenes terete or subterete.*
1. Angelica.—Petals not obcordate, lateral costæ extended into wings.
 2. Selinum.—Calyx limb obliterated. Petals obcordate. Oil tubes solitary. Lateral costæ much broader than the dorsal.
 3. Œnanthe.—Calyx limb 5-dentate. Styles erect. Oil tubes solitary. Carpophore adnate.

- cc. *Endosperm not excavated on its ventral surface (Orthospermæ). Costæ 5, filiform, secondary costæ 0. Akenes dorsally compressed, lenticular.*
1. *Heracleum*.—Petals obcordate. Oil tubes clavate. Lateral costæ of the two akenes touching each other and more distant from the 3 dorsal costæ than these amongst themselves.
 2. *Peucedanum*.—Lateral and dorsal costæ equally distant. Oil tubes superficial.
- dd. *Primary costæ 5, secondary costæ 4.*
1. *Daucus*.—Primary costæ filiform, bristly. Secondary costæ spinose.
- ee. *Endosperm concave on its ventral surface by a longitudinal furrow (Cælospermæ).*
1. *Caucalis*.—Primary costæ spinose; secondary more prominent than the primary.
 2. *Conium*.—Akenes laterally compressed. Costæ undulate, crenulate; secondary costæ 0. Oil tubes 0.
 3. *Deweya*.—Akenes laterally compressed. Costæ prominent. Secondary costæ 0. Oil tubes 2–3 in each vallecule.
 4. *Osmorrhiza*.—Akenes linear, angulate. Carpophore persistent.

C. *Flowers not umbellate.*

1. Boykinia.—Ovary 2-celled. Capsule ∞ -seeded.
2. Heuchera.—Ovary 1-celled. Capsule ∞ -seeded, dehiscent by valves.
3. Ribes.—Petals and stamens inserted on the calyx limb. Petals unguiculate. Fruit baccate, ∞ -seeded. Shrubs.

III. Trigynia.

A. *Floral parts complete, inferior.*

1. Rhus.—Fruit drupaceous, 1-pyrenous. Pyrena 1-seeded. Shrubs.
2. Calandrinia.—Sepals 2, persistent. Petals 5. Capsule 3-valved 2-celled. Placenta central, ∞ -seeded. Succulent.
3. Claytonia.—Sepals 2, persistent. Petals 5. Capsule 3-valved, 1-celled. Placenta central, few-seeded. Succulent.
4. Linum.—Sepals 5. Capsule 3-celled. Cells 2-ovulate.
5. Alsine.—Sepals 5. Petals 5. Capsule 1-celled, 3-valved. Placenta central.

B. *Floral parts complete, superior.*

1. Sambucus. Corolla rotate. Berry 3-seeded (Drupe 3-pyrenous.) Tree. Shrub.
2. Whipplea.—Ovary 3-celled, cells 1-ovulate. Fruit a septicidal capsule.

IV. Pentagynia.

1. Aralia.—Fruit a 5-pyrenous drupe. Pyrenæ 1-seeded.
2. Tillæa. — Calyx 5-parted. Petals 5. Fruit 5-follicles. Aquatic.
3. Linum.—Sepals 5. Capsule 4-5-celled, cells 2-ovulate.
4. Statice.—Calyx limb scarious. Ovary 1-celled, 1-ovulate. Styles not plumose.
5. Armeria.—Calyx limb scarious. Ovary 1-celled, 1-ovulate. Styles plumose.
6. Spergula.—Calyx 5-sepalous. Petals 5, entire. Capsule 1-celled, 5-valved. Seeds winged.
7. Sagina.—Calyx 5-sepalous. Capsule 1-celled, 4-5-valved. Seeds reniform.

V. Polygynia.

1. Myosurus.—Claws of petals longer than the lamina. Carpidia ∞ . Akenes 1-seeded.

HEXANDRIA.**I. Monogynia.****A. *Floral parts complete.***

1. Berberis.—Petals 6, inferior. Fruit a berry. Shrub.
2. Vancouveria.—Petals 6, inferior. Fruit a follicle.

3. *Trillium*.—Sepals 3. Petals 3. Stigmas 3, almost sessile.
 4. *Lythrum*.—Calyx tubular, 8–12-dentate. Petals 4–8.
 5. *Trientalis*.—Calyx 6–7 cleft. Corolla 6–8 parted, spreading. Capsule 6–8-valved, 1-celled. Placenta central.
 6. *Frankenia*.—Calyx tubular, 5-cleft. Petals 5, unguiculate. Capsule 1-celled, dehiscent by valves. Ovules few, inserted on parietal placentæ.
- B. *Floral parts incomplete. Perigonium corolline, gamosepalous.*
1. *Muilla* (*Hesperoscordum*) Perigonium deeply 6-parted, campanulate. Stamens inserted on the base of the perigonium.
 2. *Brodiaea*.—Perigonium 6-lobed, funnel-shaped. Stamens inserted on the throat, alternately sterile.
 3. *Triteleia*.—Perigonium 6-lobed, salver-shaped. Stamens inserted alternately on the throat and on the middle of the tube,
- C. *Floral parts incomplete. Perigonium corolline, 6-sepalous.*
- a. *Style 3-cleft, or 3 sessile stigmas.*
 1. *Fritillaria*.—Sepals of perigonium with a nectarium at the base.

2. Calochortus.—Alternate sepals of different shape; interior sepals larger. Stigmas sessile. Anthers basi-fixed. Capsule 3-celled.

3. Trillium.—External sepals herbaceous, internal corolline. Stigmas almost sessile. Fruit a berry.

b. *Style entire or 0. Stigma entire or 3-lobed.*

aa. *Anthers basi-fixed.*

1. Prosartes.—Fruit baccate; testa of the seeds thin, membranaceous.

2. Clintonia.—Fruit baccate; testa of the seeds crustaceous.

bb. *Anthers versatile.*

1. Lilium.—Sepals with a nectariferous longitudinal groove.

2. Allium.—Nectariferous grooves 0. Flowers umbellate, before spreading enclosed in a spathe.

3. Cyanotris (Camassia).—Sepals 6, 5 ascending, 1 deflexed. Stamens ascending. Style declined, its base persistent. Fruit a loculicidal capsule, several seeded.

4. Chlorogalum. Sepals spreading, ligulate. Stamens spreading. Style deciduous. Fruit a loculicidal capsule. Each cell 2-seeded.
5. Smilacina.—Style short, 3-lobed, persistent. Fruit baccate, few-seeded.

D. *Floral parts incomplete, Perigonium glumaceous.*

1. Juncus.—Style with 3 filiform stigmas. Fruit a capsule, ∞ -seeded.
2. Luzula.—Style with 3 filiform stigmas. Fruit a capsule, 3-seeded.

II. *Trigynia.*

1. Zygadenus.—Perigonium 6-sepalous. Sepals glandular at the base. Anthers extrorse, 1-celled.
2. Xerophyllum.—Perigonium 6-sepalous. Sepals not glandular at the base. Anthers extrorse.
3. Triglochin.—Ovaries 3-6, separated at the base. Stigmas sessile, plumose.
4. Chorizanthe. Involucre 1-3 flowered, tubular. Fruit an akene.
5. Rumex.—Ovary 1, 1-celled, 1-seeded. Stigma plumose.
6. Pterostegia.—Involucre 2-lobed, 1-flowered. Fruit an akene enveloped by the enlarged involucre.

7. *Anemopsis*.—Spadix with a several-leaved involucre. Flowers bracteate. Perigonium 0. Ovaries immersed in the rachis, 1-celled, ∞ -seeded.
8. *Frankenia*.—Calyx tubular, costate, 5-cleft. Petals 5, unguiculate. Capsule 1-celled, dehiscent by valves. Ovules few, inserted on parietal placentæ.

III. Hexagynia—Polygynia.

1. *Triglochin*.—Stigmas sessile, plumose. Aquatic.
2. *Alisma*.—Sepals 3. Petals 3. Aquatic.

HEPTANDRIA.

I. Monogynia. (Number of parts somewhat inconstant.)

1. *Trientalis*.—Calyx 7-cleft. Corolla 7-parted. Stamens inserted on the corolla.
2. *Æsculus*.—Calyx 5-dentate. Petals 4–5, irregular.

II. Trigynia.

1. *Portulaca*.—Corolla superior, Calyx 2-sepalous, deciduous. Petals 4–6. Fruit a circumscissile capsule. Succulent.
2. *Calandrinia*.—Corolla inferior. Calyx 2-sepalous, persistent. Fruit a capsule, 3-valved, ∞ -seeded.

III. Polygynia.

1. *Portulaca*.—Corolla superior. Calyx 2-sepalous, deciduous. Petals 4-6. Fruit a circumscissile capsule. Succulent.

OCTANDRIA.

I. Monogynia.

A. *Floral parts polypetalous, inferior.*

1. *Acer*.—Calyx 4-5-parted. Corolla 4-5 petals. Fruit a samara. Tree.

B. *Floral parts polypetalous, superior.*

1. *Jussiaea*.—Calyx tube not prolonged beyond the ovary, lobes of the calyx limb persistent. Petals inserted on the calyx. Stigma capitate. Fruit a septical capsule. Aquatic.

a. *Seeds with a hairy crown.*

1. *Zauschneria*.—Calyx tube considerably prolonged beyond the ovary; calyx limb petaloid, deciduous. Petals not exceeding the calyx lobes, 2-cleft, erect. Capsule 4-valved, 1-celled, ∞ -seeded.
2. *Epilobium*.—Calyx tube prolonged but little beyond the ovary. Petals inserted on an annular disc on the summit of the calyx tube. Ovary 4-celled. Capsule loculicidal.

b. *Seeds naked.*

1. *Gayophytum*.—Calyx tube not prolonged beyond the ovary. Anthers versatile. Stamens opposite the petals smaller than alternates. Ovary 2-celled. Style short. Capsule 2-celled, 4-valved.
2. *Ænothera*.—Calyx tube prolonged beyond the ovary, lobes of the limb reflexed. Anthers versatile. Ovary 4-celled, ovules ∞ . Style filiform. Capsule 4-celled, loculicidal.
3. *Godetia*.—Calyx tube prolonged beyond the ovary, funnel-shaped, lobes of the limb reflexed. Anthers basi-fixed.
4. *Clarkia*.—Calyx tube prolonged beyond the ovary, lobes of the limb reflexed. Petals unguiculate. Stamens alternately smaller. Anthers basi-fixed.
5. *Boisduvalia*.—Calyx tube prolonged beyond the ovary, lobes of the limb erect. Petals sessile, 2-lobed. Alternate stamens shorter. Anthers basi-fixed.

C. *Floral parts complete, superior. Corolla gamopetalous.*

1. *Vaccinium*.—Stamens inserted on the margin of an epigynous disc. Cells of anthers elongated into a tube at the apex. Fruit a ∞ -seeded berry. Shrub.

D. *Floral parts incomplete, inferior.*

1. *Dirca*.—Perigonium campanulate. Limb obliquely truncate. Style subterminal. Fruit a drupe. Shrub.
2. *Polygonum*.—Perigonium 4-5 sepalous, persistent. Stigmas 2-3. Fruit an akene, enclosed in the persistent perigonium.
3. *Acæna*.—Calyx tube contracted at the throat, angular, the angles armed with glochidate prickles. Stigma penicillate. Akene enclosed in the calyx-tube.

II. *Digynia and Trigynia.*

1. *Polygonum*.—Perigonium 4-5 sepalous, persistent. Fruit an akene enclosed in the perigonium.

III. *Tetragynia.*

1. *Anemopsis*.—Spadix with a several leaved involucre. Perigonium 0. Ovaries immersed in the rachis, 1-celled ∞ -ovuled.

ENNEANDRIA.

1. *Oreodaphne*.—Anthers with valvular dehiscence. Fruit a drupe resting on an enlarged thalamus. Tree.
2. *Eriogonum*.—Involucre ∞ -flowered, campanulate, 6-dentate. Styles 3. Fruit an akene.

3. *Chorizanthe*.—Involucre 1-3-flowered, tubular. Styles 3. Fruit an akene.

DECANDRIA.

I. Monogynia.

- A. *Corolla 4-5 petalous, or petals 0. Never gamopetalous.*
1. *Pyrola*.—Calyx 5-parted, persistent. Hypogynous disc 0. Anthers dehiscent by 2 pores. Seeds minute.
 2. *Cercis*.—Calyx scarcely dentate. Corolla papilionaceous. Anthers versatile. Petals unguiculate, carina 2-petalous, larger than the upper petals. Legume flat ∞ -seeded. Ventral suture winged. Shrub. Tree.
 3. *Thermopsis*.—Calyx cleft to the middle. Corolla papilionaceous. Petals of the carina partly connate, longer than vexillum. Legume linear, compressed.
 4. *Pickeringia*.—Calyx repandly 4-dentate. Corolla papilionaceous. Carina 2-petalous, as long as vexillum. Legume linear, compressed. Shrub.
 5. *Geranium*.—*Carpidia* 5, verticillate around a columnar axis. Styles connate at the apex. Stamens all fertile.
 6. *Erodium*.—*Carpidia* 5, verticillate

- around a columnar axis. Styles connate at the apex. Alternate stamens sterile.
7. *Limnanthes*.—Stamens inserted on a perigynous disc. Carpodia 5, each 1-ovulate. Style entire from its base. Aquatic.
 8. *Jussiaea*.—Petals inserted on the calyx. Stigma capitate. Fruit a septicidal capsule. Aquatic.
 9. *Adenostoma*.—Calyx funnel-shaped, the tube 10-costate. Carpidium 1. Fruit an akene enclosed in the calyx tube. Shrub.
 10. *Acæna*.—Calyx tube contracted at the throat, angular, the angles armed with glochidiate prickles. Petals 0. Stigma penicillate. Fruit an akene enclosed in the calyx tube.

B. *Corolla gamopetalous*.

1. *Rhododendron*.—Calyx 5-parted. Corolla funnel-shaped, rotate. Stamens ascending. Anthers dehiscent by an apical pore. Capsule septicidal. Shrub. Tree.
2. *Gaultheria*.—Calyx 5-cleft. Corolla urceolate. Fruit a spurious berry, that is, a capsule, 5-celled, ∞ -seeded, enclosed in the enlarged and fleshy calyx. Shrub.

3. *Arctostaphylos*.—Corolla urceolate. Ovary 5-celled, cells 1-ovulate. Fruit baccate. Shrub.
4. *Arbutus*.—Corolla urceolate. Ovary 5-celled, cells ∞ -seeded. Fruit a berry. Tree.
5. *Samolus*.—Alternate stamens sterile. Ovary 1-celled, ∞ -seeded with central placentation.

II. Digynia.

1. *Saxifraga*.—Ovary 2-celled, ∞ -seeded. Capsule loculicidal.
2. *Tellima*.—Petals lobed. Ovary 1-celled, ∞ -seeded. Styles short. Stigmas capitate. Capsule dehiscent at the apex.
3. *Tiarella*.—Petals entire. Ovary 1-celled. Styles short. Stigmas simple. Capsule dehiscent to the base; valves unequal.
4. *Acæna*.—Calyx tube contracted at the throat, angular, the angles armed with glochidiate prickles. Petals 0. Stigmas penicillate. Akenes 2, enclosed in the calyx tube.

III. Trigynia.

1. *Silene*.—Calyx gamosepalous, 5-dentate, without bracts at the base. Petals unguiculate. Capsule dehiscent by teeth. Seeds reniform.

2. *Alsine*.—Sepals 5, unchanged in fruit. Petals entire. Capsule dehiscent by 3 valves. Seeds reniform.
3. *Spergularia* (*Lepigonum*). — Sepals 5. Petals entire. Capsule dehiscent by 3 valves. Seeds compressed. Stipules membranaceous.
4. *Arenaria*. — Sepals 5. Petals entire. Capsule dehiscent by 6 valves. Seeds reniform.
5. *Stellaria*. — Sepals 5. Petals 2-cleft. Capsule dehiscent by 6 valves.
6. *Portulaca*.—Corolla superior. Calyx 2-sepalous, deciduous. Petals 4-6. Capsule circumscissile.
7. *Calandrinia*.—Corolla inferior. Calyx 2-sepalous, persistent. Capsule 3-valved, ∞ -seeded.
8. *Whipplea*.—Ovary 3-septate. Cells 1-ovulate. Capsule septicidal.

IV. Tetragynia.

- 1.—*Sagina*.—Sepals 4. Petals entire. Stigmas alternate with the sepals. Capsule dehiscent by 4-5 valves.

V. Pentagynia.

1. *Sagina*.—Sepals 5. Petals entire. Stigmas alternate with the sepals. Capsule dehiscent by 5 valves. Seeds reniform. Stipules 0.

2. *Spergula*.—Sepals 5. Petals entire. Stigmas alternate with the sepals. Capsule dehiscent by 5 valves. Valves opposite to the sepals. Stipules membranaceous.
3. *Cerastium*. — Sepals 5. Petals 2-cleft. Capsule dehiscent by 10 teeth.
4. *Oxalis*.—Stamens monadelphous at their base. Capsule 5-lobed.
5. *Cotyledon* (*Echeveria*).—Calyx 5-parted. Petals coherent by their claws. Ovaries 5. Fruit 5-follicles. Succulent.
6. *Sedum*.—Calyx 5-parted. Petals not coherent by their claws. Ovaries 5. Fruit 5-follicles. Succulent.
7. *Portulaca*.—Corolla superior. Calyx 2-sepalous, deciduous. Petals 4-6. Capsule circumscissile. Succulent.

VI. Polygynia.

1. *Horkelia*.—Calyx companulate, limb 5-parted, augmented by 5 bractlets. ovaries ∞ , inserted on a conical receptacle. Styles subterminal.

DODECANDRIA.

I. Monogynia.

1. *Asarum*.—Perigonium 3-cleft, superior.
2. *Portulaca*.—Calyx 2-sepalous, deciduous. Petals 4-6. Capsule circumscissile. Succulent.

3. Calandrinia.—Corolla inferior. Calyx 2-sepalous, persistent. Capsule 3-valved, ∞ -seeded. Succulent.
4. Lythrum.—Calyx tubular, inferior, 8-12-dentate. Petals 4-6.
5. Adenostoma. — Calyx funnel-shaped, tube 10-costate. Petals 5. Fruit an akene enclosed in the calyx tube. Shrub.
6. Cercocarpus.—Calyx tube cylindrical. Petals 0. Style terminal, long-exserted. Fruit an akene, linear, terete, caudate by the elongated plumose style. Shrub.

II. Pentagynia.

1. Nuttallia.—Ovaries 5, each 2-ovulate. Styles subterminal. Fruit several drupes. Shrub.

ICOSANDRIA.

I. Monogynia.

A. *Floral parts complete, inferior.*

1. Prunus.—Fruit a fleshy drupe. Endocarp not rugose. Shrubs. Tree.
2. Adenostoma.—Calyx funnel-shaped; tube 10-costate. Petals 5. Akene enclosed in the calyx tube. Shrubs.
3. Cercocarpus.—Calyx tube cylindrical. Petals 0. Style terminal, long-exserted. Akene linear, terete, caudate by the elongated exserted plumose style. Shrub.

4. *Portulaca*. — Calyx 2-sepalous, deciduous. Capsule circumscissile.
5. *Calandrinia*. — Calyx 2-sepalous, persistent. Capsule 3-valved, ∞ -seeded.
6. *Eschscholtzia*. — Calyx gamosepalous, circumscissile. Ovary terete. Stigmas 4. Capsule 1-celled, 2-valved, seeds on the margins of the valves.

II. Di-Tri-Tetra-Pentagynia.

1. *Nuttallia*. — Ovaries 5, each 2-ovulate. Style subterminal, subinternal. Fruit several drupes. Shrub.
2. *Photinia* (*Heteromeles*). — Ovaries 2, imperfectly united, half superior. Styles 2. Fruit a pomum, 2-celled. Cells 1-seeded. Tree.
3. *Amelanchier*. — Ovary inferior, 3-5-celled, each cell imperfectly divided. Fruit a pomum, 3-5-seeded. Endocarp bony (drupe). Shrub.
4. *Spiræa*. — Ovaries several, several seeded. Fruit several follicles. Testa of the seeds membranaceous. Shrub.
5. *Neillia*. — Ovaries several, several seeded. Fruit several follicles. Testa of the seed crustaceous. Shrub.
6. *Horkelia*. — Calyx augmented by 5 bractlets. Ovaries on a conical receptacle. Styles subterminal. Fruit akenes.

7. *Mesembrianthemum*.—Calyx partitions unequal. Petals ∞ . Ovary ∞ -celled, ∞ -seeded. Fruit a capsule. Succulent.
8. *Hypericum*.—Calyx 4-5-parted. Petals 4-5. Fruit a capsule.
9. *Mentzelia*.—Ovary inferior. Calyx limb 5-parted, persistent. Capsule 1-celled. Placentæ 3, parietal.

III. *Polygynia*.

1. *Spiræa*.—Ovaries superior, several seeded. Fruit several follicles. Shrub.
2. *Rosa*.—Calyx 5-parted. Petals 5. Fruit akenes enclosed in the calyx tube forming a spurious berry. Shrub.
3. *Calycanthus*.—Calyx cup-shaped, limb ∞ -parted, ∞ -seriate. Petals 0. Fruit ∞ akenes enclosed in the persistent calyx tube. Shrub.
4. *Rubus*.—Calyx 5-parted. Petals 5. Fruit ∞ drupes on a conical receptacle. Shrubs.
5. *Horkelia*.—Calyx campanulate, augmented by 5 bractlets. Styles subterminal. Fruit ∞ akenes.
6. *Potentilla*.—Calyx flattened, augmented by 5 bractlets. Ovaries on a slightly conical receptacle. Styles lateral. Akenes on a herbaceous thalamus.

7. *Fragaria*.—Calyx flattened, augmented by 5 bractlets. Ovaries on a convex receptacle. Styles lateral. Akenes on a fleshy receptacle.
8. *Mesembrianthemum*.—Calyx partitions unequal. Petals ∞ . Ovary ∞ -celled, ∞ -seeded. Fruit a capsule. Succulent.

POLYANDRIA.

I. Monogynia.

A. *Petals 4.*

1. *Dendromecon*.—Calyx 2-sepalous. Ovary linear. Stigma sessile, 2-lobed. Capsule 1-celled, 2-valved, seeds on the margin of the valves. Shrub.
2. *Eschscholtzia*.—Thalamus top-shaped. Calyx gamosepalous, circumscissile. Ovary terete. Stigmas 4. Capsule 1-celled, 2-valved, seeds on the margin of the valves.
3. *Meconopsis*.—Calyx 2-sepalous, caducous. Anthers laterally dehiscent. Ovary 1-celled. Placentæ more than 2. Style short. Stigma radiate.
4. *Argemone*.—Calyx 2-3 sepalous. Petals 4. Anthers extrorse. Ovary 1-celled, placentæ more than 2. Style 0. Stigma radiate.

5. *Actæa*.—Calyx 4-sepalous. Anthers adnate, introrse. Ovary 1. Placenta ventral. Style 0. Berry ∞ -seeded.

B. *Petals 5*.

1. *Helianthemum*.—Calyx double. External sepals 2, sometimes 0; internal 3, contorted in æstivation. Capsule 3-valved.

C. *Petals 6*.

1. *Argemone*.—Calyx 3-sepalous. Filaments filiform. Stigmas radiate.
2. *Platystemon*.—Calyx 3-sepalous. Filaments flattened their entire length, ovary at last separating into its component carpidia.
3. *Platystigma*.—Calyx 3-sepalous. Filaments flattened only at the base, ovary not separating but ripening into a 3-valved capsule.

II. **Di-, Tri-, Tetra-, Penta-, Polygnia.**

A. *Ovaries ventrally dehiscent. Fruit a follicle.*

a. *Flowers irregular.*

1. *Delphinium*. Calyx petaloid. Superior sepal calcarate.

b. *Flowers regular.*

1. *Platystemon*.—Sepals 3, caducous. Petals 6. Filaments flattened. Carpidia at first united into an ∞ -celled ovary.

2. *Pæonia*.—Sepals 5, herbaceous, carpidia distinct from the beginning.
 3. *Aquilegia*.—Sepals petaloid. Petals funnel-shaped, calcarate.
- B. *Ovaries 1-ovulate, indehiscent. Fruit, akenes.*
- a. *Æstivation of sepals imbricate.*
 1. *Ranunculus*.—Petals unguiculate, claws shorter than the lamina, a nectarium near each claw.
 2. *Myosurus*.—Claws of the petals longer than the lamina.
 3. *Anemone*.—Sepals petaloid. Petals 0. Thalamus convex, conical.
 4. *Thalictrum*. Sepals petaloid. Petals 0. Thalamus flat, discoid.
 - b. *Æstivation of sepals valvate.*
 1. *Clematis*.—Sepals 4, petaloid. Petals 0.

DIDYNAMIA.

I. *Gymnospermia.*

- A. *Stamens and style enclosed. An interrupted ring of hairs in the corolla below the insertion of the stamens.*
 1. *Marrubium*.—Akenes flat at their tops.
- B. *Style always exerted. An uninterrupted ring of hairs in the corolla below the insertion of the stamens.*

- a. *Calyx 2-labiate. Stamens parallel.*
1. *Prunella*.—Calyx, closed in fruit.
- b. *Calyx 5-dentate. Stamens parallel.*
1. *Stachys*.—Calyx 5–10-nerved. Lobes of the lower lip of the corolla obtuse. Stamens ascending.
 2. *Sphacele*.—Calyx reticulately veined. Lobes of the corolla 5, all obtuse. Stamens ascending, Shrub.
- c. *No ring of hairs in the corolla.*
1. *Scutellaria*.—Upper lip of the corolla concave. Stamens close together, parallel. Calyx 2-labiate, lips entire.
 2. *Mentha*.—Stamens distant and divergent, but straight. Corolla funnel-shaped, 4-cleft, superior lobe emarginate.
 3. *Micromeria*.—Calyx teeth nearly equal. Stamens ascending.
 4. *Monardella*.—Stamens straight, divergent, exerted. Cells of the anthers at length also divergent. Corolla lobes narrow.
 5. *Pycnanthemum*.—Stamens straight, distant, divergent. Cells of the anthers parallel to the last.
 6. *Pogogyne*.—Stamens ascending, convergent in pairs. Style villous. Calyx deeply 5-cleft.

7. *Acanthomintha*.—Lower pair of stamens ascending. Cells of the anthers divaricate. Calyx 2-labiate, teeth spinose.
8. *Lophanthus*.—Upper pair of stamens longer and declined, lower pair shorter and ascending. Anther cells parallel. Calyx 15-nerved.
9. *Trichostemma*.—Stamens long-exserted, lower pair longer. Tube of corolla slender; limb 5-cleft, lobes oblong, declined.

II. Angiospermia.

A. *Base of the anther cells mucronate.*

a. *Ovary 1-celled ∞-ovulate, placentæ parietal.*

1. *Anoplanthus*.—Flowers without bractlets. Parasitic. Chlorophyll 0.
2. *Aphyllon*.—Flowers with bractlets. Parasitic. Chlorophyll 0.

b. *Ovary 2-celled, ∞-ovulate. Fruit 1-∞-seeded.*

1. *Pedicularis*.—Calyx 5-dentate, superior lip of corolla laterally compressed.

B. *Base of anther cells not mucronate.*

a. *Ovary 1-celled. Placenta ∞-seeded, central.*

1. *Limosella*.—Calyx 5-dentate. Corolla subregular.

b. *Ovary 2-celled.*

1. *Cordylanthus*.—Calyx spathaceous. Cells of anthers distant and of different shape and insertion. Lips of corolla short, upper lip laterally compressed.
2. *Orthocarpus*.—Calyx spathaceous, cleft vertically. Cells of anthers distant and of different shape and insertion. Corolla lips personate, but upper lip the smaller.
3. *Castilleia*.—Calyx spathaceous, cleft vertically. Cells of anthers distant and of different shape and insertion. Corolla ringent, the upper lip the larger.
4. *Mimulus*.—Stigma dilated, petaloid. Calyx angular, angles carinate. Placentæ of the capsule remain united and only separate at last near the apex. Valves of the capsule membranaceous.
5. *Mimetanthe*.—Stigma dilated, petaloid. Calyx campanulate, not angulate. Capsule dehiscent only by the dorsal suture.
6. *Diplacus*. — Stigma dilated, petaloid. Calyx angular; angles carinate. Placentæ of the capsule meeting, but in dehiscence separating their whole length. Valves of the capsule coriaceous. Tube of the corolla funnel-shaped.

7. *Eunanus*. — Stigma dilated, petaloid. Calyx angular, angles carinate. Placentæ of the capsule separate in dehiscence for their whole length. Tube of corolla slender, filiform.
 8. *Collinsia*. — Stigma small. Corolla personate, lower lip 3-lobed, middle lobe laterally compressed, carinate, hiding the stamens.
 9. *Scrophularia*. — Stigma small. Corolla short, globular. Middle lobe of lower lip reflexed.
 10. *Pentstemon*. — Stigma small. Corolla ringent. One sterile stamen besides the didynamous ones.
 11. *Antirrhinum*. — Corolla personate, saccate at the base. Capsule opening by pores.
 12. *Linaria*. — Corolla personate, calcarate. Capsule dehiscent by valves.
 13. *Lippia*. — Calyx 2-cleft. Cells of ovary 1-seeded. Fruit drupaceous, separating into 2 nutlets.
- c. *Ovary 4-celled*.
1. *Verbena*. — Cells of ovary 1-ovulate. Fruit drupaceous, splitting into 4 parts.

TETRADYNAMIA.

I. Siliculosa.

A. Fruit a 2-articulate lomentum.

1. Cakile.—Both articles of the lomentum anceps.

B. Fruit a schizocarp.

1. Senebiera.—Silicula 2-seeded, margin not winged.

C. Fruit an indehiscent silicula.

1. Thysanocarpus.—Silicula 1-seeded, margin winged.

D. Silicula regularly dehiscent, dorsally compressed, septum narrow.

1. Lepidium.—Cells 1-seeded.
2. Capsella.—Cells ∞ -seeded.

E. Silicula regularly dehiscent, laterally compressed, septum broad.

1. Alyssum.—Cells 1-4-ovulate.

II. Siliquosa.

A. Siliqua indehiscent.

1. Raphanus.—Siliqua moniliform.

B. Siliqua regularly dehiscent.

1. Cardamine. — Siliqua laterally compressed. Valves linear, without nerves. Ovules in each cell ∞ , 1-seriate.

2. *Nasturtium*. — Siliqua linear. Valves without nerves. Ovules in each cell ∞ , 2-seriate.
3. *Cheiranthus*. — Siliqua laterally compressed. Septum broad. Valves 1-nerved. Ovules in each cell 1-seriate. Lateral sepals saccate. Petals unguiculate.
4. *Caulanthus*. — Siliqua terete, valves 1-nerved. Ovules in each cell 1-seriate. Lateral sepals saccate. Petals undulate.
5. *Streptanthus*. — Siliqua laterally compressed. Valves 1-nerved. Ovules in each cell 1-seriate. All sepals equal. Petals undulate.
6. *Barbarea*. — Siliqua 4-angular. Valves concave, 1-nerved. Ovules in each cell 1-seriate.
7. *Erysimum*. — Siliqua 4-angular. Valves carinate, 1-nerved. Ovules in each cell 1-seriate.
8. *Tropidocarpum*. — Siliqua dorsally compressed. Valves 1-nerved, carinate. Septum very narrow. Ovules in each cell 2-seriate.
9. *Arabis*. — Siliqua laterally compressed. Valves 1-nerved. Ovules in each cell 1-seriate. Petals not undulate.

10. *Sisymbrium*.—Siliqua linear. Valves 3-nerved, concave. Ovules in each cell 1-seriate.
11. *Brassica*.—Siliqua terete. Valves 3-5-nerved, concave. Ovules in each cell 1-seriate.

MONADELPHIA.

I. Triandria.

1. *Sisyrinchium*.

II. Pentandria.

1. *Erodium*.—Stamens alternating with staminodia.
2. *Downingia*.—Corolla gamopetalous, irregular.

III. Octandria.

1. *Polygala*.

IV. Decandria.

1. *Erodium*.—Alternate stamens sterile. Fruit 5 akenes with persistent styles.
2. *Geranium*.—Stamens all fertile. Fruit 5 akenes with persistent styles.
3. *Oxalis*.—Styles 5. Fruit a 5-lobed capsule.
4. *Lupinus*.—Corolla papilionaceous.
5. *Amorpha*.—Corolla represented by a single petal. Shrub.

V. Polyandria.

1. *Lavatera*. — Calyx double. External calyx 3-6-cleft. Shrub.
2. *Malva*. — Calyx double. External calyx 3-sepalous.
3. *Sidalcea*. — External calyx 0. Stigmas not capitate.
4. *Sphæralcea*. — Stigmas capitate. Cells of capsule 2-ovulate. Shrub.
5. *Sida*. — Stigmas capitate. Cells of capsule 1-ovulate.

DIADELPHIA.

I. Hexandria.

1. *Dicentra*. — Flowers 2-calcarate.

II. Octandria.

1. *Polygala*. — Flowers irregular. Ovary 2-celled. Cells 1-ovulate.

III. Decandria.

A. *Sub-monadelphous*.

1. *Lupinus*. — Anthers alternately oblong and reniform.
2. *Amorpha*. — Corolla represented by a single petal. Shrub.
3. *Psoralea*. — Wings of the corolla united to the carina. Ovary sessile, 1-ovulate. Legume indehiscent, included in the calyx.

B. *Genuinely diadelphous.*

a. *Style not pilose.*

1. *Hosackia*.—Petals unguiculate. Carina rostrate.
2. *Trifolium*.—Petals persistent, adnate to the staminal tube.
3. *Psoralea*.—Wings of the corolla adnate to the carina. Ovary sessile, 1-ovulate. Legume indehiscent, included in the calyx.
4. *Glycyrrhiza*.—Petals not adnate to the staminal tube. Carina 2-petalous.
5. *Astragalus*.—Petals not adnate to the staminal tube. Carina blunt. Legume imperfectly 2-celled.
6. *Melilotus*.—Petals not adnate to the staminal tube. Ovary straight. Legume globose.
7. *Medicago*.—Petals not adnate to the staminal tube. Ovary crescent-shaped. Legume spiral.

b. *Style pilose.*

1. *Lathyrus*.—Style pilose only on the ventral side which is flattened.
2. *Vicia*.—Style pilose on the dorsal side, or all around the apex. Style not spiral.

POLYADELPHIA.

1. *Hypericum*. — Ovary superior. Calyx 4-5-parted. Petals 4-5. Fruit capsular.
2. *Mentzelia*. — Ovary inferior. Calyx limb 5-parted, persistent. Capsule 1-celled. Placentæ 3, parietal.

SYNGENESIA.

I. *Æqualis*.

A. *Flowers ligulate*.

a. *Pappus plumose*.

aa. *Receptacle paleaceous*.

1. *Hypochæris*. — Paleaæ of the receptacle deciduous.

bb. *Receptacle not paleaceous*.

1. *Microseris*. — Pappus paleaceous at base. Akenes more than 5-costate, truncate at apex. Scapigerous.
2. *Stephanomeria*. — Pappus setaceous, somewhat plumose. Akenes 5-costate, truncate at apex and base. Caulescent.
3. *Rafinesquia*. — Bristles of pappus arachnoid, Akenes rostrate,
4. *Malacothrix*. — Pappus 1-seriate. Bristles scabrous, silky, deciduous in a ring. Apex of akenes developed into a crown,

5. Troximon.—Pappus capillary. Bristles deciduous singly, not in a ring. Apex of akenes rostrate; costæ smooth.
6. Taraxacum.—Pappus capillary. Bristles persistent. Apex of akenes elongated, rostrate; costæ roughened.
7. Sonchus.—Akenes flattened, not rostrate. Pappus silky, white.
8. Hieracium.—Akenes terete, not rostrate, Pappus brittle, discolored.

B. *Flowers tubular.*

a. *Receptale naked.*

aa. *Branches of the style club-shaped.*

1. Adenostyles. — Involucre 1-seriate. Branches of the style filiform. Pappus copious, silky, white.
2. Bulbostylis.—Involucre imbricate, 2-3 seriate. Branches of the style filiform. Pappus copious, silky.

bb. *Branches of the style flat and tipped with an appendix.*

aaa. *Parts of pappus ∞.*

1. Bigelovia.—Involucre imbricate. Pappus 1-seriate, capillary, brownish at maturity.

2. Aplopappus.—Involucre imbricate. Pappus 1-seriate. Bristles rigid.
3. Chrysopsis.—Involucre imbricate. Pappus 2-seriate, internal capillary, external scaly.
4. Lessingia.—Ray-flowers enlarged, but not ligulate.
5. Baccharis.—Dioecious. Shrub.

bbb. *Parts of pappus definite in number.*

1. Pentachæta. — Pappus of 5-scabrous bristles, shorter than the corolla, sometimes depauperate.
2. Grindelia.—Pappus aristate.

cc. *Branches of style not flat, but tapering, sometimes with an appendix.*

aaa. *Scales of the involucre herbaceous.*

1. Chænactis.—Pappus paleaceous.
2. Senecio.—Pappus capillary.

bbb. *Scales of the involucre scarious.*

1. Artemisia.—Receptacle slightly convex. Pappus 0. Akenes obovate with small epigynous disc.
2. Tanacetum.—Receptacle slightly convex. Pappus 0. Akenes costate with large epigynous disc.

3. *Matricaria*.—Receptacle cylindrical, hollow.
4. *Baccharis*.—Dioecious.
 - b. *Receptacle paleaceous or setaceous.*
 - aa. *Branches of style connected near the apex,*
1. *Cnicus*.—Receptacle densely bristly. Bristles of the pappus plumose, connected at the base into a deciduous ring. Filaments of stamens distinct.
2. *Silybum*.—Receptacle densely bristly. Bristles of pappus not plumose, 1-seriate.
3. *Carthamus*.—Pappus 0.
4. *Centaurea*.—Receptacle densely bristly. Bristles of pappus not connected at their base into a ring.
 - bb. *Branches of style entirely distinct.*
1. *Bidens*.—Involucre 2-seriate, external spreading, internal erect. Pappus of 2-4 awns.
2. *Madia*.—Pappus 0. Receptacle sometimes only paleaceous near the ray and naked at the disc.

II. Superflua.

A. *Receptacle naked.*

- a. *Anthers obtuse at their base, at least not calcarate.*

aa. *Branches of the style flat and tipped with an appendix.*

aaa. *Parts of pappus ∞ .*

1. *Bigelovia*.—Involucre imbricate; scales dry. Pappus 1-seriate, capillary, at maturity brownish. Ray flowers yellow.
2. *Aplopappus*.—Involucre imbricate, scales foliaceous. Pappus 1-seriate, bristles rigid. Ray flowers yellow.
3. *Chrysopsis*.—Involucre imbricate. Pappus 2-seriate, internal capillary, external scaly. Ray flowers yellow.
4. *Lessingia*.—Ray flowers enlarged but not ligulate.
5. *Solidago*.—Involucre imbricate, scales foliaceous. Pappus silky. Ray flowers yellow.
6. *Aster*.—Involucre imbricate, scales foliaceous. Ray flowers not yellow, 1-seriate.
7. *Erigeron*.—Involucre imbricate, scales foliaceous. Ray flowers not yellow, ∞ -seriate.

bbb. *Parts of pappus definite in number.*

1. *Pentachæta*.—Scales of involucre not herbaceous. Pappus of 5 scabrous bristles shorter than the corolla, sometimes depauperate. Rays yellow.

2. Grindelia.—Scales of the involucre herbaceous. Pappus aristate. Rays yellow.
3. Gutierrezia.—Scales of the involucre herbaceous. Pappus short, paleaceous. Rays yellow.
- bb. *Branches of the style not flat but tapering, sometimes with an appendix.*
 1. Coinogyne.—Involucre regularly imbricate. Scales broad.
 2. Burriellia.—Involucre consisting of 3-5 scales.
 3. Bæria.—Scales of the involucre in a single series. Receptacle conical, minutely muricate. Akenes linear.
 4. Eriophyllum (Bahia).—Scales of the involucre not embracing any akenes. Receptacle convex (not conical). Akene linear, with large terminal areola.
 5. Monolopia.—Scales of involucre united at base. Receptacle conical. Akene oblong, with small terminal areola.
 6. Lasthenia.—Scales of involucre united into a dentate cup. Receptacle conical. Pappus 0.
 7. Rigiopappus.—Scales of involucre linear. Receptacle flat. Pappus 4-5 aristate scales.

8. *Helenium*.—Scales of the involucre reflexed. Pappus of hyaline scales. Akenes turbinate, costate.
9. *Senecio*.—Scales of the involucre 1-seriate, sometimes calyculate. Pappus capillary, copious.
10. *Arnica*.—Scales of involucre lanceolate, linear, all equal. Pappus capillary, 1-seriate.

b. *Anthers calcarate*.

1. *Pluchea*.—Scales of involucre imbricate, herbaceous.
2. *Micropus*.—Scales of involucre scarious, ♀ flowers enclosed in marginal scales, which are laterally compressed and conduplicate. Pappus 0.
3. *Stylocline*.—Scales of involucre scarious. Receptacle cylindrical. ♀ flowers protected by carinate scales.
4. *Filago*.—Scales of involucre scarious. Scales subtending ♀ flowers not carinate.
5. *Anaphalis*.—Scales of involucre scarious. ♀ flowers filiform. Style of ♂ flowers nearly blunt.
6. *Gnaphalium*.—Scales of involucre scarious. ♀ flowers filiform. Style of ♂ flowers 2-cleft.

B. *Receptacle paleaceous.*

a. *Anthers obtuse or sagittate at their base, but never calcarate.*

aa. *Involucre 1-seriate or nearly so.*

1. *Iva*.—♀ flowers reduced to short tubes embracing the base of the style. ♂ flowers: anthers scarcely connate.
2. *Balsamorhiza*.—Akenes of the disc 4-angular. Pappus 0.
3. *Wyethia*.—Akenes 4-angular or laterally compressed. Pappus paleaceous, crown-shaped.
4. *Pugiopappus*.—Akenes villous. Pappus 2-aristate.
5. *Leptosyne*.—Akenes naked, margined. Pappus almost 0.
6. *Madia*.—Involucre in a simple row, each scale wrapped about a laterally compressed akene. Pappus 0.
7. *Holozonia*.—Involucre in a simple row, each scale completely embracing an akene. Pappus of the ray akenes hyaline, of the disc generally 0.
8. *Hemizonia*.—Involucre 1-seriate, each scale embracing with its base half of a ray akene. Pappus of the ray 0.

9. *Lagophylla*.—Ray flowers 5 in number, cuneiform, 3-lobed. Involucre 5 scales, each entirely embracing a ray akene. Receptacle flat.
10. *Layia*.—Capitulum ∞ -flowered. Involucre 1-seriate. Scales acuminate, each entirely enclosing a ray akene. Pappus of the ray 0.
11. *Achyrachæna*.—Ray flowers short and hidden by the scales of the 1-seriate involucre. Receptacle naked in the centre, paleaceous toward the margin. Pappus paleaceous. Paleæ ∞ -seriate, silvery.

bb. *Involucre imbricate.*

1. *Achillea*.—Involucre of few series of scales. Rays short, oval.
2. *Anthemis*.—Involucre ∞ -seriate. Rays long, ligulate.

b. *Anthers calcarate.*

1. *Psilocarphus*.—Scales of the involucre few. ♀ flowers each wrapped in a palea. Pappus 0.
2. *Stylocline*.—Scales of the involucre few. ♀ flowers each bracteate by a carinate palea, but not wrapped in it. Pappus a few caducous bristles.

3. *Evax*.—Scales of the involucre 1-2 seriate. Pappus 0.
4. *Filago*.—Scales of the involucre imbricate. Pappus pilose.

III. Frustranea.

1. *Corethrogyne*.—Receptacle naked.
2. *Centaurea*.—Receptacle bristly.
3. *Bidens*.—Paleæ of the receptacle deciduous.
4. *Helianthus*.—Paleæ of the receptacle persistent. Pappus paleaceous, caducous.
5. *Helianthella*. Paleæ of the receptacle persistent. Pappus 2 persistent awns, besides the caducous paleæ.

IV. Necessaria.

A. *Receptacle naked.*

a. *Anthers obtuse at the base, at least not calcarate.*

aa. *Branches of the style of the ♀ flowers flat and tipped with an appendix.*

1 *Erigeron.*

bb. *Branches of the style not flat, but tapering, sometimes with an appendix.*

1. *Hemizonia*. — Involucre 1-seriate, each scale with its base half embracing a ray akene. Pappus of ray 0.

2. *Holozonia*. — Involucre 1-seriate, each scale completely embracing a ray akene. Pappus of the ray hyaline.
3. *Lagophylla*. — Ray flowers 5, cuneiform, 3-lobed. Involucre of 5 scales, each completely enclosing a ray akene. Receptacle flat. Paleæ 5, between disc and ray. Pappus 0.
4. *Blennosperma*. — Scales of the involucre 1-seriate, united at the base, not embracing the akene. Pappus 0.
5. *Artemisia*. — Involucre imbricate, scarious. Flowers of the ray slender, dentate.
6. *Soliva*. — Flowers of the ray ∞ -seriate, without corolla. Pappus 0.
7. *Nardosmia*. — Pappus pilose.

B. *Receptacle villous.*

1. *Artemisia*.

C. *Receptacle paleaceous.*

a. *Anthers obtuse or sagittate at their base, but not calcarate.*

1. *Lagophylla*. — Ray flowers 5, cuneiform, 3-lobed. Involucre of 5 scales, each completely enclosing a ray akene. Receptacle flat. Paleæ 5, between disc and ray. Pappus 0.
2. *Holozonia*. — Involucre 1-seriate, each scale embracing completely a ray akene. Pappus of the ray hyaline.

3. Hemizonia.—Involucre 1 - seriate, each scale with its base half embracing a ray akene. Pappus of ray akene 0.

4. Madia.—Involucre 1-seriate, each scale completely enwrapping a laterally compressed ray akene. Pappus 0.

b. *Anthers almost distinct.*

1. Iva.

c. *Anthers calcarate.*

1. Evax.—Scales of involucre 1—2-seriate. Pappus 0.

2. Stylocline.—Scales of involucre very few. Flowers of the ray bracteate by a carinate palea, but not enwrapped by it. Pappus a few caducous bristles.

3. Psilocarphus.—Scales of the involucre few. Each ray flower enwrapped in a palea. Pappus 0.

V. Monogamia.

1. Downingia.

GYNANDRIA.

I. Monandria.

a. *Labellum calcarate.*

1. Habenaria.

b. *Labellum not calcarate.*

1. Epipactis.—Labellum geniculate.

2. Corallorhiza. Labellum not geniculate, adnate to the column. Aphyllous and without chlorophyll.
3. Spiranthes.—Labellum not geniculate, embracing the column with its base. Perigonium oblique.

II. Diandria.

1. Cyripedium.

III. Hexandria.

1. Aristolochia.

MONŒCIA.

I. Monandria.

1. Euphorbia. — Involucre campanulate, sometimes 2-phyllous. Flowers umbellate, androgynous. ♂ flowers without perigonium, stipitate, bracteate, ♀ flowers single, central. Ovary 3-celled, styles 3, each 2-cleft.
2. Zostera.—Spathe elongated into a lamina. Spadix androgynous. Ovary 1-celled, 1-ovulate. Fruit a nutlet. Aquatic.
3. Lilæa.—Flowers spicate. ♂, ♀, and androgynous spikes present, besides solitary axillary ♀ flowers. Aquatic.
4. Najas. — Flowers axillary, solitary. ♂ flowers included in a spathe, ♀ flowers naked. Ovary 1, 1-ovulate. Fruit drupaceous. Aquatic.

5. *Zannichellia*.—Flowers axillary. ♂ flower naked, ♀ flower with campanulate perigonium. Ovaries 4, each 1-ovulate. Fruit nutlets. Aquatic.
6. *Callitriche*.—Flowers axillary, 2-bracteolate. Ovary 4-celled; cells 1-ovulate. Fruit a 4-coccous schizocarp. Aquatic.

II. Diandria.

1. *Pinus*.—Anthers sessile on the bracts of the ♂ amentum. Leaves fasciculate in a sheath. Coniferous.
2. *Pseudotsuga*.—Anthers sessile on the bracts of the ♂ amentum. Leaves distichous. Branchlets smooth, the leaf scars not being prominent. Coniferous.
3. *Fraxinus*.—Flowers paniculate. Calyx obsolete, 4-cleft. Fruit a 1-seeded samara.

III. Triandria.

1. *Typha*.—♂ spike and ♀ spike cylindrical. ♂ spike above ♀ spike. Paleæ 0.
2. *Sparganium*.—♂ spike and ♀ glomerulate. Flowers separated by a paleaceous perigonium. Aquatic.
3. *Carex*.—Glumaceous. Spikes unisexual or androgynous. Spikelets 1-flowered. ♂ spikelet an external palea; ♀ spikelet with 2 paleæ, the interior one transformed into a utricle including the 1-seeded ovary.

4. Phoradendron.—Flowers immersed in the rachis of articulate spikes. Perigonium 3-lobed. Ovary inferior. Fruit a berry. Parasitic.
5. Megarrhiza.—Flowers complete. ♂ flowers in racemes, ♀ flower solitary in the axilla of the ♂ raceme.

IV. Tetrandria.

A. *Flowers complete.*

1. Ptelea.—Calyx 4-parted. Petals 4. Stamens alternate with the petals. Fruit a 2-seeded samara. Shrub. Tree.

B. *Flowers perigoniate in ♂ and ♀.*

1. Urtica.—♀ perigonium 2-sepalous.
2. Hesperocnide.—♀ perigonium gamosepalous.

C. *Perigonium 0, or only present in one sex.*

1. Cupressus.—♂ flowers: anthers attached to a peltate bract. Cone globose. Seeds angulate, narrowly winged. Leaves opposite. Coniferous.
2. Libocedrus.—♂ flowers: anthers attached to a peltate bract. Cone oblong. Wings of the seed unequal. Leaves opposite. Coniferous.

3. Sequoia.—Anthers of the ♂ flowers attached to a peltate bract. Cone ovate. Leaves alternate. Coniferous.
4. Alnus.—♂ flowers with 3-4 cleft perigonium. ♀ flowers: perigonium 0.

V. Pentandria—Polyandria .

A. *Flowers complete, not amentaceous.*

1. Myriophyllum.—♂ flower: stamens 8. ♀ flower: ovary inferior, 4-celled, 4-ovulate. Aquatic.
2. Ptelea.—Calyx 5-parted. Petals 5. Stamens alternating with petals. Style 1. Fruit a 2-seeded samara. Shrub.
3. Rhus.—Calyx 5-parted, persistent. Petals 5. Stamens 5, alternating with the petals. Ovary 1-celled, 1-ovulate. Styles 3. Fruit a drupe.
4. Æsculus.—Calyx irregular, gamosepalous. Petals 5, unguiculate. Filaments ascending. Ovary 3-celled. Style 1. Cells 2-ovulate. Fruit a 1-seeded capsule. Shrub. Tree.
5. Acer.—Stamens 8. Ovary 2-celled, 2-lobed. Fruit a 2-seeded samara. Tree.

B. *Flowers complete. ♂ flowers amentaceous.*

1. Juglans.—♂ perigonium, 2-6 parted. ♀ flowers with superior 4-dentate calyx. Petals 4. Tree.

C. *Flowers incomplete not amentaceous.*

1. *Amarantus*.— Perigonium 3-5-sepalous. Fruit a circumscissile capsule.
2. *Atriplex*.— ♂ perigonium 5-sepalous. ♀ perigonium 2-parted. Stigmas 2. Fruit a compressed utricle.
3. *Ceratophyllum*.— ♂ perigonium ∞ -sepalous. Anthers sessile. Ovary of the ♀ flower 1-celled, 1-ovulate. Fruit a nutlet with persistent style. Aquatic.
4. *Eremocarpus*.— ♂ perigonium 5-6 parted. ♀ perigonium 0. Ovary 1-celled, 1-ovulate. Fruit a 2-valved capsule.
5. *Xanthium*.— ♂ inflorescence: involucre ∞ -phyllous, ∞ -flowered. Perigonium 5-dentate. ♀ inflorescence: involucre gamophyllous, 2-flowered. Fruit a pseudocarp enclosed in the indurated involucre.
6. *Ambrosia*.— ♂ inflorescence: involucre gamophyllus, ∞ -flowered. ♀ inflorescence: involucre gamophyllus, 1-flowered. Spines of the involucre 1-seriate.
7. *Franseria*.— ♂ inflorescence: involucre gamophyllus, ∞ -flowered. ♀ inflorescence: involucre gamoyhyllous, 1-4-flowered. Spines of the involucre ∞ -seriate.

8. Iva.—Involucre gamophyllus. Head androgynous. ♂ flowers funnel-shaped, central; ♀ flowers tubular, peripheral.
9. Platanus.—Flowers without perigonium, densely capitulate on a globose receptacle. Tree.

D. *Flowers incomplete; ♂ flowers amentaceous. Stamens inserted on a perigonium.*

1. Castanopsis.—Involucre of ♀ flowers 1-3-flowered, afterward enclosing the nut in the shape of a prickly burr.
2. Quercus.—Involucre of ♀ flowers 1-flowered, surrounding the base of the nut in the shape of a cup. Trees. Shrubs.

E. *Flowers incomplete. ♂ flowers or both ♀ and ♂, amentaceous. Stamens inserted on bract.*

1. Corylus.—Fruit a nut, inclosed in a 2-cleft, lacinate involucre.
2. Myrica.—♀ flowers amentaceous. Fruit, drupaceous nutlets. Involucre 0. Tree.

DIECIA.

I. Diandria.

1. Salix.—Amentaceous. Perigonium 0. Fruit an ∞-seeded capsule. Shrub.
2. Fraxinus.—Flowers paniculate. Calyx obsolete, 4-cleft. Fruit a 1-seeded samara. Tree.

II. Triandria.

1. *Salix*.—Amentaceous. Perigonium 0.
Shrub. Tree.
2. *Phoradendron*.—Flowers immersed in
the rachis of an articulate spike.
Perigonium 3-lobed. Anthers 2-celled
dehiscent by 2 pores. Berry globose,
1-seeded.
3. *Arceuthobium*.—♂ perigonium 3-lobed.
♀ 2-dentate. Anthers 1-celled, dehis-
cent by a slit. Berry compressed, 1-
seeded.
4. *Atriplex*.—♂ perigonium 3-sepalous.
♀ 2-parted. Fruit a utricle.
5. *Amarantus*.—Perigonium 3-parted.
Fruit a capsule with circumscissile de-
hiscence.
6. *Brizopyrum* (*Distichlis*).—Glumaceous.
Inferior glume ∞ -nerved.
7. *Poa*.—Glumaceous. Inferior glume 1-
nerved.

III. Tetrandria.

1. *Urtica*.—♂ flowers 4-sepalous. ♀
flowers 2-sepalous. Stigma penicillate.
2. *Negundo*.—Perigonium depauperate, 4-
cleft. Ovary 2-lobed, 2-celled. Fruit
a 2-seeded samara. Tree.
3. *Ptelea*.—Calyx 4-5 sepalous. Petals 4.
Fruit a 2-seeded samara. Tree.

4. *Garrya*.—Flowers amentaceous. Bracts decussate, connate in pairs. ♂ flowers 4-parted. Perigonium of ♀ flowers rudimentary. Styles 2. Fruit a berry. Tree.

IV. Pentandria.

1. *Rhus*.—Calyx 5-parted, persistent. Petals 5. Stamens alternating with the petals. Ovary 1-celled, 1-seeded. Styles 3. Fruit a drupe. Tree. Shrub.
2. *Ptelea*.—Calyx 5-parted, petals 5. Stamens alternating with the petals. Style 1. Fruit a 2-seeded samara. Tree.
3. *Rhamnus*.—Calyx 4-cleft. Petals 0. Stamens alternating with the partitions of the calyx. Fruit 2-4-celled, 2-4-seeded. Shrub.
4. *Vitis*.—Calyx obsolete. Petals inserted on a disc, coherent at their apex, and deciduous at their base. Fruit baccate.
5. *Negundo*.—Perigonium depauperate, 5-cleft. Ovary 2-lobed, 2-celled. Fruit a 2-seeded samara. Tree.
6. *Juniperus*.—♂ flowers amentaceous. Stamens adnate to the base of a scale. ♀ flowers amentaceous. Ovules naked, 2 adnate to the base of each bract. Fruit a galbulus.

7. *Atriplex*.—♂ perigonium 5-sepalous. ♀ perigonium 2-parted. Fruit a utricle.
8. *Amarantus*.— Perigonium 5-sepalous. Fruit a capsule with circumscissile dehiscence.

V. Hexandria.

1. *Rumex*.—Ovary 1-celled, 1-seeded. Styles 3. Stigma plumose.

VI. Octandria.

1. *Populus*.—Amentaceous. ♀ flowers cup-shaped. Fruit an ∞-seeded capsule.

VII. Dodecandria,

1. *Nuttallia*.—Flowers complete. Ovaries 5. Fruit 1-5 drupes. Shrub.
2. *Hendecandra* (*Croton*).—♂ perigonium 5-cleft. ♀ perigonium 3-cleft. Ovary 3-celled. Fruit a 3-coccus capsule.

VIII. Icosandria.

1. *Nuttallia*.—Flowers complete. Shrub.
2. *Torreya*.—♂ flowers amentaceous. ♀ flowers single, ovule naked, immersed in an urceolate arillus, in fruit simulating a drupe. Coniferous.

IX. Polyandria.

1. *Thalictrum*.

X. Syngenesia.

1. *Baccharis*. Shrub.

SYNOPSIS OF GENERA AND SPECIES.

Vascular Plants.

CLASS 1. PHANEROGAMÆ.

Sub-Class 1. ANGIOSPERMÆ.

DIVISION 1. DICOTYLEDONES.

SERIES 1. TETRACYCLICÆ.

Plants with a tendency to arrangement of the floral parts in 4 well-defined circles, and with a well established numeric (quinary) law.

Sub-Series 1. GAMOPETALÆ.

(Petals Consolidated.)

Section ANISOCARPÆ. Number of carpidia less than 5.

Sub-Section 1. EPIGYNÆ.

ORDER 1—SYNANDRÆ. Stipules 0. Filaments flattened.

Family 1. COMPOSITÆ.

Inflorescence capitulate. Calyx changed into a pappus. Anthers syngenetic. Ovary 1-celled, 1-seeded; fruit an akene.

Sub-Family 1. *Tubulifloræ*.

Flowers, at least of the disc, tubular.

Tribe 1. ASTEROIDEÆ.

Anthers without tails. Style branches of disc-flowers, flat, with an appendix. Leaves alternate.

a. *Parts of the pappus of definite number.*

1. *Gutierrezia* Lag.

Superflua. Heads few-flowered. Ray-flowers about 3, ligulate; disc-flowers about 5. Scales of the involucre coriaceous, imbricate, their tips green, reflexed. Akenes terete. Pappus of about 9, paleaceous scales.— α . β . Flowers yellow.

1. G. CALIFORNICA. Torr. & Gray. Suffrutescent; leaves linear.—Dry river beds.—Sonoma. Livermore. Summer.

2. *Grindelia* Willd. GUM-PLANT.

Superflua. *Æqualis*. Head ∞ -flowered. Ray-flowers ligulate, 1-seriate. Scales of the hemispherical involucre in several series, their green tips squarrose. Akenes compressed. Pappus aristate.

Resinous herbs; flower buds before opening, bearing a drop of milky-looking resin.— α . β . Flowers yellow.

1. *G. HIRSUTULA*. Hook. Hirsutely pubescent.—Dry hillsides. Common. Summer.

An infusion of the herb has been recommended against asthma by the Californians of Spanish descent.

2. *G. GLUTINOSA* Dunal. Glabrous; scales of the involucre with short tips; awns of the pappus 5 or more.—Livermore. Summer.

3. *G. CUNEIFOLIA* Nutt. Glabrous; scales of the involucre squarrose from the base; awns of the pappus 5 or more; leaves clasping, obtuse.—Salt marshes at Alvarado. Summer.

4. *G. ROBUSTA* Nutt. Glabrous; scales of involucre squarrose from the base; awns of the pappus less than 5; leaves clasping, acute. Salt marshes. Common. Summer. Has been recommended against the eczema caused by *Rhus*—(Poison oak.)

3. *Pentachæta* Nutt.

Superflua. *Æqualis*. Ray-flowers variable, or altogether wanting. Scales of the involucre lanceolate, scarious on the margin and tip, loosely imbricate in about two series. Receptacle convex. Pappus about 5 scabrous bristles shorter than the disc-corolla, sometimes depauperate.—☉. Small vernal herbs.

1. *P. BELLIDIFLORA* Greene. Peduncles glabrous; ray-flowers white.—Corte Madera.

2. *P. EXILIS* Gray. Stem erect; peduncles villous; ray-flowers 0; disc-flowers purple. Marin County.

3. *P. ALSINOIDES* Greene. Stem diffuse, much branched; capitula almost sessile in the axillæ of branches; ray-flowers 0; disc-flowers purple.—Marin County.

b. *Parts of the pappus* ∞.

4. *Lessingia* Cham.

Æqualis. *Superflua*. Ray-flowers with palmate limb. Receptacle flat, naked. Akenes compressed, silky-villous; pappus 1-seriate; bristles ∞, scabrous, rigid.—☉

1. *L. GERMANORUM* Cham. Limb of ray-flowers unequally lobed; stems spreading on the ground; lower leaves spatulate, pinnatifid; flowers yellow.—Sandhills. San Francisco. Summer.

2. *L. RAMULOSA* Gray. Limb of the ray-flowers unequally lobed; stem erect, diffusely branched; leaves not pinnatifid; flowers purple.—Marin County. Summer.

3. *L. LEPTOCLADA* Gray. Limb of ray-flowers equally lobed; stem erect; branches filiform, terminated by the capitula; flowers purple, white.—Nicasio. Summer.

5. *Chrysopsis* Nutt. GOLDEN ASTER.

Superflua. *Æqualis*. Head ∞ -flowered; ray-flowers ligulate. Akenes compressed; pappus 2-seriate; outer row short paleaceous scales, inner row long scabrous bristles.— $\frac{1}{2}$. Flowers yellow.

1. *C. SESSILIFLORA* Nutt. Capitula radiate. Tamalpais. Summer.

2. *C. OREGANA* Gray. Capitula not radiate.—Livermore. Autumn.

6. *Aplopappus* Cass.

Superflua. *Æqualis*. Head ∞ -flowered. Ray-flowers ligulate. Bristles of the pappus ∞ . Flowers yellow.

1. *A. LINEARIFOLIUS* DC. Rays more than 9.— $\frac{1}{2}$. Contra Costa mountains. Summer.

2. *A. ERICOIDES* Hook. Ray-flowers less than 10.— $\frac{1}{2}$. Sand hills. San Francisco. Summer.

7. *Bigelovia* DC.

Æqualis. Superflua by a single ray-flower. Head few-flowered. Receptacle narrow. Pappus 1-seriate; bristles ∞ . Flowers yellow.

1. *B. ARBORESCENS* Gray. Scales of the involucre irregularly imbricate and not in distinct ranks.— $\frac{1}{2}$. Tamalpais. Summer.

8. *Solidago* L. GOLDEN ROD.

Superflua. Head ∞ -flowered. Ray-flowers ligulate, few, distant from each other. Akenes terete, ∞ -costate; pappus of ∞ , capillary bristles.— ζ . Autumn. Suffrutescent plants, with long rod-like branches. Flowers yellow.

1. *S. OCCIDENTALIS* Nutt. Stem branching. ζ . Moist places. Summer.

2. *S. CALIFORNICA*. Nutt. Stem single; the whole plant pubescent.— ζ . Moist sand. Summer.

3. *S. SEMPERVIRENS* L. Stem simple; the plant completely glabrous.— ζ . Salt marshes near San Francisco. Autumn.

All *Solidago* species seem to be possessed of diuretic powers. *S. Virgaurea*, a European species, formerly officinal, is still in use as a domestic remedy.

9. *Corethrogyne* DC.

Frustranea. Head ∞ -flowered; ray-flowers ∞ , 1-seriate; scales of the involucre imbricate; style appendages of disc-flowers with tuft-like bristles; pappus of simple, unequal rigid bristles.— ζ . Ray-flowers blue, purple; habit of aster.

1. *C. OBOVATA* Benth. Contra Costa range. Marin County. Summer.

10. *Aster* (Nees.)

Superflua. Head ∞ -flowered. Ray-flowers 1-seriate, ligulate. Scales of involucre imbricate. Style appendages triangular-lanceolate to subulate. Pappus of copious capillary, scabrous bristles. — 2f. Ray-flowers rose-color, white, blue.

1. *A. RADULINUS* Gray. Pappus rigid; some of the bristles thickened toward the top; rays white.—2f. Berkeley. Autumn.

2. *A. CHAMISSONIS* Gray. Pappus soft; stem glabrous, erect and branching, leafy; involucre imbricate, its scales with green tips; ray-flowers violet.—2f. San Francisco at the cemetery. Autumn.

3. *A. DIVARICATUS* Nutt. Pappus soft; stem glabrous, diffusely branched; branches slender; involucre imbricate, its scales with scarious margins.— \odot . Salt marshes, San Francisco. Autumn.

11. *Erigeron* L. FLEA-BANE.

Superflua. Necessaria. Head ∞ -seriate, ligulate; scales of the involucre imbricate, linear. Pappus scanty of scabrous, capillary bristles.

1. *E. STENOPHYLLUS* Nutt. Stem leafy; leaves linear, scabrous, deep green; rays purple.—2f. Livermore. Autumn.

2. *E. GLAUCUS* Ker. Radical leaves differing from cauline, glaucous, entire, succulent; ray purple.— \mathcal{L} . Seashore. Summer. Autumn.

3. *E. PHILADELPHICUS* L. Leaves irregularly dentate; ray-flowers narrow, numerous, reddish.— \mathcal{L} . Colma. Berkeley. Autumn.

4. *E. CANADENSIS* L. Leaves numerous, small, rays inconspicuous, whitish.— \odot . Waste grounds. Common. Summer. Autumn.

12. *Baccharis* L.

Dioecious. Head without rays. Scales of the involucre ∞ -seriate, imbricate. δ corolla tubular with 5-cleft limb. Pappus capillary, 1-seriate. ♀ corolla filiform, truncate. Pappus copious, capillary.

1. *B. PILULARIS* DC. Leaves sessile cuneiform, sinuately dentate; flowers whitish.— $\frac{1}{2}$. Summer. Autumn. Common.

2. *B. DOUGLASHII* DC. Glutinous; leaves lanceolate, acute, distinctly 3-nerved; capitula in a compound, terminal corymb; receptacle conical.— \mathcal{L} . Moist sand. San Francisco. Summer.

3. *B. VIMINEA* DC. Shrub with the aspect of a willow; leaves lanceolate, acute; capitula in terminal corymbs and racemes, receptacle flat.— $\frac{1}{2}$. River-bed, Niles. Livermore. Summer. Autumn.

Tribe II. INULEÆ.

Anthers with tails, sagittate. Style branches without appendages.

13. *Pluchea* Cass. MARSH-FLEABANE.

Superflua. Necessaria. Head ∞ -flowered. ♀ flowers ∞ -seriate, tubular. Scales of the involucre ∞ -seriate, ovate, imbricate. Receptacle flat, naked. Pappus 1-seriate.

1. *P. CAMPHORATA* DC. Flowers rose-color. —⊙. Salt marshes. Autumn.

14. *Adenocaulon* Hook.

Necessaria. Head few-flowered, all the flowers tubular, Anthers sagittate, not tailed. Scales of the involucre 5, 1-seriate, herbaceous, at last reflexed. Receptacle flat, naked. Akenes exserted, obovate, glandular near the summit; pappus 0.—2. Flowers pale.

1. *A. BICOLOR* Hook.—Redwoods. Summer.

15. *Micropus* L.

Necessaria. Superflua. Head of few flowers, all tubular. ♂ 5, ♀ 5, embraced by the scales of the two-seriate involucre. Receptacle narrow, naked. Akenes falling off with the embracing scales of the involucre. Pappus 0. —⊙. Flowers inconspicuous.

1. *M. CALIFORNICUS* Fisch. & Meyer. Low woolly herb. Common. Spring.

16. Psilocarphus Nutt.

Necessaria. Superflua. Head ∞ -flowered. δ flowers few, tubular, 5-dentate. ♀ flowers ∞ -seriate, filiform, each wrapped in a palea. Scales of the involucre few, scarious. Receptacle convex, paleaceous externally, naked in the center. Akenes enveloped in the paleæ. Pappus 0.— \odot . Flowers inconspicuous.

1. *P. OREGANUS* Nutt. Covered with loose, white wool; akenes cylindrical.—Low ground near creeks. Spring.

2. *P. TENELLUS* Nutt. Covered with appressed grey wool. Akenes fusiform.—Low ground near creeks. Spring.

17. Evax Gærtn.

Necessaria. Superflua. Head ∞ -flowered. δ flowers tubular, few— ♀ flowers ∞ -seriate, filiform. Scales of the involucre few, 1–2-seriate, scarious. Receptacle elongate (like the axis of a spike), paleaceous externally, naked in the center with the δ flowers. Pappus 0.— \odot . Flowers inconspicuous.

1. *E. CAULESCENS* Gray. — Woolly annual. Gravelly alluvium. San Rafael. Spring.

18. Filago L.

Superflua. Head ∞ -flowered. δ flowers tubular, 4-dentate. ♀ flowers ∞ -seriate, filiform; external rows hidden between

the scales of the involucre and the paleæ. Scales of the involucre imbricate, similar to the paleæ. Receptacle elongate (like the axis of a spike). Pappus of ∞ , capillary bristles, depauperate towards the periphery. External row ♀ ; pappus pilose. ☉. Flowers inconspicuous.

1. *F. CALIFORNICA* Nutt. Woolly annual. Common. Spring.

19. *Anaphalis* DC. PEARLY-EVERLASTING.

Necessaria. Superflua. Head ∞ -flowered. ♂ tubular, 5-dentate; style scarcely divided, blunt; ♀ flowers ∞ -seriate, filiform. Scales of the involucre imbricate, ∞ -seriate, radiating; external rows ovate; internal rows longer and narrower. Receptacle flat, naked. Pappus 1-seriate, capillary, its bristles scabrous and distinct at base.—2f.

1. *A. MARGARITACEA* Benth. & Hook.—Common. Summer.

20. *Gnaphalium* L. EVERLASTING.

Superflua. Head ∞ -flowered; ♂ flowers tubular, 5-dentate. Style 2-cleft. ♀ flowers filiform, ∞ -seriate; involucre ovate. Scales imbricate, 8-seriate, scarious; as long as the head. Receptacle flat, naked. Pappus of 1-seriate, capillary bristles. Flowers pale.

1. *G. MICROCEPHALUM* Nutt. Bristles of pappus not united at their base into a ring; involucre mainly scarious, decidedly imbricate; leaves but slightly decurrent; white-woolly; involucre turbinate.—2f. Contra Costa.

2. *G. SPRENGELII* Hook. & Arn. Bristles of pappus not united at their base into a ring; involucre mainly scarious, decidedly imbricate; leaves but slightly decurrent, white-woolly; involucre hemispherical.—○. Common. Summer.

3. *G. DECURRENS* Ives. Bristles of pappus not united at their base into a ring; involucre mainly scarious, decidedly imbricate; leaves decidedly decurrent, glandular, white-woolly only underneath; involucre companulate. Smell of the plant like liquorice.—○. Sandhills. San Francisco. Summer.

4. *G. RAMOSISSIMUM* Nutt. Bristles of pappus not united at their base into a ring; involucre mainly scarious, decidedly imbricate; leaves decidedly decurrent, linear, involucre turbinate. Smell of the plant like liquorice.—○. Sand dunes. Saucelito. Summer.

5. *G. PALUSTRE* Nutt. Bristles of pappus not united at their base in a ring; scales of involucre woolly, only scarious at their tips, all nearly of the same length.—○. Moist grounds. Common. Summer.

6. *G. PURPUREUM* L. Bristles of pappus united at the basè into a ring.— \mathcal{Z} . Near salt marshes. Common. Summer.

Tribe III. AMBROSIEÆ.

Anthers of \mathfrak{z} or \mathfrak{d} flowers distinct, not syngenetic. Style abortive, truncate. Corolla of \mathfrak{z} flowers rudimentary or 0; pappus 0.

21. *Iva* L.

Necessaria. Superflua. Head ∞ -flowered; \mathfrak{z} or \mathfrak{d} flowers ∞ , tubular, 5-lobed; \mathfrak{z} flowers few, 5-dentate. Scales of the involucre 3 to 4, ovate. Paleæ of the receptacle linear.— \mathcal{Z} .

1. *I. AXILLARIS*. Pursh.—Near the seacoast. Summer.

22. *Ambrosia* DC. RAGWEED.

Monœcious. \mathfrak{d} head ∞ -flowered; scales of the involucre united into a cup; receptacle flat, naked; \mathfrak{z} head 1-flowered. Scales of the involucre united into a cup. Corolla 0. Akenes enclosed in the persistent involucre.

1. *A. ARTEMISIÆFOLIA* L. Leaves all bipinnatifid.— \odot . Cultivated grounds. Summer.

2. *A. PSILOSTACHYA* DC. Upper leaves pinnatifid, lower bipinnatifid.— \mathcal{Z} . Cultivated grounds. Summer.

23. Franseria Cav.

Monœcious. ♂ head ∞-flowered. Scales of the involucre united into a cup. Paleæ of the flat receptacle filiform. ♀ head 1-, sometimes 2-4-flowered. Scales of the involucre united into a cup; if the cup be more than 1-flowered, forming as many cells as there are flowers. Involucre armed with ∞ rows of spines. Akenes enclosed in the persistent involucre.—2f.

1. *F. BIPINNATIFIDA* Less. Leaves 2-3-pinnately divided.—Sand dunes, San Francisco. Summer.

2. *F. CHAMISSONIS* Less. Leaves cuneate to ovate, obtusely serrate, the lower sometimes laciniate.—Sand dunes, San Francisco. Summer.

24. Xanthium Tourn. COCKLEBUR.

Monœcious. ♂ head ∞-flowered. Scales of the involucre 1-seriate, distinct. Receptacle cylindrical, paleaceous. ♀ head 2-flowered. Scales of the involucre united into a cup, armed with hooked spines. Corolla filiform. Akenes enclosed in the persistent involucre.—⊙.

1. *X. STRUMARIUM* L. Leaves green on both sides, scabrous, irregularly serrate.—Roadsides. Common. Summer.

2. *X. SPINOSUM* L. Leaves whitish beneath, most of them deeply lobed.—Waste grounds. Common. Summer.

Tribe IV. HELIANTHÆ.

Anthers without tails. Receptacle paleaceous. Pappus not capillary.

a. *Paleæ persistent.*

25. *Balsamorhiza* Hook. Nutt.

Superflua. Head ∞ -flowered. Ray-flowers lanceolate. Receptacle flat, paleaceous. Pappus 0.--2. Flowers yellow.

1. *B. DELTOIDEA* Nutt. Leaves with cordate base, entire to serrate.—Niles. Spring.

2. *B. Hookeri*. Nutt. Leaves pinnately or bipinnately parted.—Hillsides near Lake Chabot. Spring.

26. *Wyethia* Nutt.

Superflua. Head ∞ -flowered; ray-flowers lanceolate. Scales of the involucre 2—3-seriate: external foliaceous; internal paleaceous. Receptacle flat, paleaceous. Pappus cup-shaped or aristate.—2. Flowers yellow.

1. *W. HELENIOIDES* Nutt. Involucre foliaceous, spreading; pappus chaffy, not aristate; young plant tomentose; akenes pubescent towards the apex.—Dry hillsides. Mission Dolores. Spring.

2. *W. GLABRA* Gray. Involucre foliaceous, spreading; pappus chaffy, not aristate; plant glabrous, glutinous; akenes glabrous.—Tamalpais. Spring.

3. *W. ANGUSTIFOLIA* Nutt. Pappus aristate.
Marin County. Spring.

27. *Helianthella* Torr. & Gray.

Frustranea. Head ∞ -flowered. Receptacle flat; paleaceous. Akenes flattened; pappus represented by 2, marginal, aristæ, alternating with 2, caducous paleæ.— \mathcal{M} . Flowers yellow.

1. *H. CALIFORNICA* Gray.—Napa. Spring.

28. *Helianthus* L. SUNFLOWER.

Frustranea. Head ∞ -flowered. Receptacle flat, paleaceous. Akenes slightly compressed; quadrangular. Pappus represented by 2, marginal caducous; paleæ with minute, intermediate ones. Ray-flowers yellow.

1. *H. ANNUUS* L. Receptacle flat; paleæ of receptacle 3-cleft; disc brown; lower leaves cordate, serrate.— \odot Common. Summer.

2. *H. SCABERRIMUS* Benth. Receptacle flat; paleæ of receptacle entire, aristate, the awn as long as the disc-flowers; disc brown. \odot . San Rafael. Summer.

3. *H. EXILIS* Gray. Receptacle flat; paleæ of receptacle entire, aristate, the awn longer than the disc-flowers; disc brown.— \odot . Lake county. Summer.

4. *H. CALIFORNICUS* DC. Receptacle convex; paleæ of receptacle blunt, not aristate; disc-flowers yellow, only their anthers brown.— \mathcal{M} . Berkeley hills. San Rafael. Summer.

b. *Paleæ deciduous*.

29. *Pugiopappus* Gray.

Superflua. Head ∞ -flowered. Ray-flowers broad, deeply crenate. Involucre 2-seriate. Pappus of the disc 0; of the ray 2-aristate. Flowers yellow. Leaves alternate.— \odot .

1. *P. CALLIOPSIDEUS* Gray.—Alma. Summer.

30. *Leptosyne* DC.

Superflua. Head ∞ -flowered. Ray-flowers broad, deeply crenate. Involucre 2-seriate, Pappus 0. Flowers yellow.— \odot .

1. *L. STILLMANI* Gray.—Alma. Summer.

31. *Bidens* L. BEGGAR-TICKS.

Frustranea. *Æqualis*. Head ∞ -flowered. Involucre 2-seriate. Pappus 2-4-aristate. Leaves opposite.

1. *B. CHRYSANTHEMOIDES* Michx. Flowers yellow.— \odot . Swamps near San Francisco (extinct) Marin county. Summer.

32. *Madia* Mol. TAR-WEED.

Superflua. Necessaria. *Æqualis*. Ray-flowers ligulate, 3-dentate, scarcely longer than the involucre. Involucre 1-seriate, herbaceous; scales carinate, complicate, embracing the ray-akenes. Receptacle naked in the center, with 1-2 rows of paleæ between the disc and the ray. Akenes laterally compressed; pappus of the ray 0; of the disc usually 0. Viscid, resinous herbs. Flowers yellow.

1. *M. NUTTALLII* Gray. Ray-flowers exerted and conspicuous; paleæ of the pappus fimbriate.— \mathcal{L} . Redwoods, Marin county. Spring,

2. *M. RADIATA* Kell. Ray-flowers exerted and conspicuous, obtusely 3-dentate; pappus 0.— \odot . Antioch. Spring.

3. *M. ELEGANS* Don. Ray-flowers exerted and conspicuous, acutely 3-lobed; pappus 0. \odot .—Common. Spring and summer.

4. *M. SATIVA* Mol. Ray-flowers short, inconspicuous; pappus 0; akenes of the disc 4-nerved, quadrangular.— \odot . Common. Summer.

5. *M. DISSITIFLORA* Gray. Ray-flowers short, inconspicuous; pappus 0; akenes of the disc without the four nerves and angles.— \odot . Marin county. Summer.

6. *M. FILIPES* Gray. Ray-flowers short, inconspicuous; pappus 0; disc-flower solitary. ☉.—Antioch. Spring.

All the species covered by a resinous exudation of strong, generally disagreeable odor. Recommended by the old settlers in affections of the urinary organs.

33. Hemizonia DC. TAR-WEED.

Necessaria. Superflua. Ray-flowers ligulate, dentate. Involucre 1-seriate, with concave scales embracing the ray-akenes. Receptacle flat, naked in the center, with a row of paleæ between disc and ray or sometimes throughout. Akenes of the ray convex externally, flat internally; pappus of the ray 0; pappus of the disc scaly, aristate, plumose or 0.—☉. Viscid, resinous herbs; flowers yellow, white.

1. *H. LUZULÆFOLIA* DC. Only the ray-akenes developed, these obovate, triangular; terminal area depressed; rays 3-lobed, frequently white.—☉. San Francisco. Summer.

2. *H. MACRADENIA* DC. Ray-flowers numerous with short ligulæ. Ray-akenes turgid, gibbose, the gibbosity pushing the terminal area to the inner angle so that the area appears lateral; receptacle conical; leaves not pungent; flowers always yellow.—☉. Tamalpais. Contra Costa. Autumn.



(99)

3. *H. PARRYI* Greene. Ray-flowers numerous with short ligulæ; ray-akenes turgid, very gibbose, the gibbosity pushing the terminal area to the inner angle so that the area appears lateral, receptacle convex; bracts of involucre pungent; paleæ of receptacle not pungent; upper leaves short, acerose, lower pinnatifid; flowers yellow.—⊙. St. Helena. Summer.

4. *H. PUNGENS* Torr. & Gray. Ray-flowers numerous with short ligulæ; ray-akenes turgid, very gibbose, the gibbosity pushing the terminal area to the inner angle so that the area appears lateral; receptacle convex; upper leaves and bracts of the involucre pungent; paleæ of the receptacle pungent also; lower leaves frequently bipinnatifid; flowers yellow. ⊙. Common. Summer.

5. *H. CORYMBOSA* Torr. & Gray. Ray-flowers cuneate, more than 10. Ray-akenes turgid, very gibbose, the gibbosity pushing the terminal area to the inner angle so that the area appears lateral; akenes 4-5-nerved; receptacle flat, naked on the disc, but the disc flowers separated from the ray-flowers by a cup, formed by connate paleæ; ovary of the disc-flowers not developed; flowers yellow.—⊙. Common. Summer.

6. *H. ANGUSTIFOLIA* DC. Ray-flowers cuneate, more than 10; ray-akenes turgid, very

gibbose, the gibbosity pushing the terminal area to the inner angle so that the area appears lateral; akenes 3-nerved, receptacle flat, naked on the disc, but the disc-flowers separated from the ray by paleæ, connate at their base. Flowers yellow.—☉. Berkeley. Tamalpais. Summer.

7. *H. TRUNCATA* Gray. Ray-flowers few, but with very large 3-lobed yellow ligula. Plant glabrous. ☉. Marin county. Summer.

8. *H. MULTIGLANDULOSA* Gray. Ray-flowers few but with very large, 3-lobed ligula; plant hispid-glandular.—☉. Tamalpais. Summer.

Properties the same as *Madia*.

34. *Holozonia* Greene.

Necessaria. Superflua. Disc flowers ∞ ; ray-flowers 5 to 8, cuneiform, 3-cleft. Scales of the involucre corresponding in number to the ray-flowers, and completely embracing each akene. Receptacle flat, the paleæ united into a cup enclosing the disc-flowers. Pappus of the disc generally 0; of the ray hyaline, spreading—2♂. Flowers white.

1. *H. FILIPES* Greene.—2♂. Sonoma. Autumn.

35. *Lagophylla* Nutt.

Necessaria. Superflua. Ray-flowers 5, cuneiform, 3-lobed. Involucre of 5 scales, acu-

minate, with scarious margins, each completely enclosing the corresponding ray-akene. Receptacle flat, with 5 paleæ between the disc and ray. Pappus 0.—⊙. Flowers yellow, whitish.

1. *L. RAMOSISSIMA* Nutt. Pubescent.—⊙. Tamalpais. Summer.

2. *L. CONGESTA* Greene. Hispid.—⊙. Tamalpais. Summer.

36. *Layia* Hook. & Arn.

Superflua. Head ∞ -flowered; ray-flowers cuneiform, 3-dentate. Involucre 1-seriate, herbaceous; scales acuminate, with scarious margins, completely enclosing the ray-akenes. Receptacle flat, with a row of paleæ between disc and ray, or paleaceous throughout. Ray-akenes linear, attenuate at the base, and with a flat area at the top; pappus of the disc various; of the ray 0.—⊙.

1. *L. CARNOSA* Torr. & Gray. Pappus bristles plumose; ray-flowers inconspicuous, small. Sands of sea shore. Marin County. Spring.

2. *L. HETEROTRICA* Hook. & Arn. Pappus bristles plumose, their hairs erect; ray-flowers large, conspicuous, 3-lobed, white.—Antioch. Livermore. Niles. Spring.

3. *L. ELEGANS* Torr. & Gray. Pappus bristles plumose, their hairs woolly and interlaced; ray-flowers conspicuous, yellow.—Marin County. Spring.

4. *L. HIERACIOIDES* Hook. & Arn. Pappus bristles plumose, their hairs straight and erect; rays conspicuous, yellow, but little longer than the disc-flowers.—Marin County. Spring.

5. *L. GAILLARDIOIDES* Hook. & Arn. Pappus-bristles plumose; their hairs straight and erect; ray-flowers conspicuous, yellow; considerably longer than the disc-flowers. Contra Costa. Saucelito. Spring.

6. *L. PLATYGLOSSA* Gray. Pappus aristate; awns equal; ray-flowers yellow, frequently with white tips.—Saucelito. Spring.

7. *L. CALLIGLOSSA* Gray. Pappus aristate; awns unequal; ray-flowers yellow with white tips.—Berkeley. Spring.

8. *L. CHRYSANTHEMOIDES* Gray. Pappus 0; ray-flowers yellow with white tips.—Alameda. Milbrae. Spring.

37. *Achyrachaena* Schauer.

Superflua. Ray-flowers short and hidden. Involucre 1-seriate; scales with scarious margins, embracing the ray-akenes. Receptacle flat, naked in the centre, paleaceous towards the margin. Akenes clavate, those of the disc truncate, those of the ray with an epigynous disc; pappus of about 10 shining, silvery scales in two series, the outer considerably shorter than the alternate inner ones.—☉. Flowers yellow or pale.

1. *A. MOLLIS* Schauer.—Common. Spring.

Tribe V. HELENIEÆ.

Receptacle not paleaceous. Scales of the involucre herbaceous; pappus not capillary.

38. *Coinogyne* Less. (*Jaumea* Pers.)

Superflua. Involucre 2-seriate, imbricate; scales rounded. Receptacle conical, naked. Pappus 0.—2. Flowers yellow, leaves opposite, fleshy.

1. *C. CARNOSA* Gray.—Salt marshes around San Francisco. Summer.

39. *Burriellia* DC.

Superflua. Ray-flowers few and short. Scales of involucre 5, ovate. Receptacle subulate, naked. Pappus of the ray 2-aristate (sometimes 0), as long as the corolla; of the disc 4-aristate.—⊙. Flowers yellow. Leaves opposite.

1. *B. MICROGLOSSA* DC.—San Francisco, Spring.

40. *Bæria* Fisch. & Mey.

Superflua. Ray-flowers ovate, exserted. Scales of the involucre 10, ovate. Receptacle conical, naked, rough. Akenes angled or nerved; pappus paleaceous or aristate, sometimes 0. Flowers yellow. Leaves opposite.—⊙.

1. *B. MACRANTHA* Gray. Pappus aristate, sometimes 0; akenes not quadrangular; recep-

tacle muricate - roughened; leaves 3 - nerved, ciliate.—⊙. Marin County. Spring.

2. *B. CHRYSOSTOMA* Fisch & Mey. Pappus 0; akenes not quadrangular; receptacle muricate-roughened; leaves linear, not ciliate.—⊙. Common. Spring.

3. *B. GRACILIS* Gray. Akenes quadrangular; pappus uniform, aristate; receptacle muricate-roughened; plant hirsutely pubescent. Common. Spring.

4. *B. CARNOSA* Greene. Pappus uniform, aristate; akenes quadrangular; receptacle muricate-roughened; plant glabrous.—⊙. Salt marshes. Spring.

5. *B. FREMONTII* Gray. Pappus 4 awns, alternating with narrow, small paleæ; receptacle muricate-roughened. Besides the entire leaves, some palmate ones.—⊙. Contra Costa. Spring.

6. *B. ULIGINOSA* Gray. Pappus 2-3 awns, alternating with broad fimbriate paleæ (sometimes pappus altogether 0); receptacle muricate-roughened. Besides the entire leaves, some pinnately or bipinnately lobed ones.—⊙. Common. Spring.

41. *Eriophyllum* Lag (*Bahia* DC.)

Superflua. Ray - flowers ovate, exserted. Scales of the involucre lanceolate, united at the base; receptacle naked, alveolate. Pappus

4, 8, or 12 membranaceous scales. Flowers yellow.

1. *E. STÆCHADIFOLIUM* Lag. Frutescent; heads with short peduncles in loose cymes; receptacle alveolate.— $\frac{1}{2}$. San Francisco. Summer.

2. *E. CONFERTIFLORUM* Gray. Suffrutescent; Head with very short peduncles in compact cymes; receptacle not alveolate.— $\frac{1}{2}$. Common. Summer.

3. *E. CÆSPITOSUM* Dougl. Herbaceous; heads solitary, or few on large peduncles.— $\frac{1}{2}$. Marin County. Summer.

42. *Monolopia* DC.

Superflua. Scales of the involucre united into a dentate cup. Receptacle conical, naked, papillate. Pappus 0. Flowers yellow.

1. *M. MAJOR* DC. Ray-flowers with 3-4 lobed ligulæ and an appendage on the opposite side; bracts of the involucre united.— \odot . Oakland. Saucelito. Summer.

2. *GRACILENS* Gray. Ray-flowers with 3-4-lobed ligulæ and an appendage on the opposite side; bracts of the involucre distinct to the base.— \odot . Santa. Cruz Mountains. Summer.

43. *Lasthenia* Cass.

Superflua. Scales of the involucre united into a dentate cup. Receptacle conical, naked,

papillose. Akenes linear compressed. Pappus 5 to 10 firm scales (sometimes 0).—⊙. Flowers yellow.

1. *L. GLABERRIMA* DC. Pappus paleaceous. Near salt marshes. Spring.

2. *L. GLABRATA* Lindl. Pappus 0. Perfoliate.—Common. Spring.

3. *L. CALIFORNICA* Lindl. Pappus 0. Leaves clasping, not perfoliate.—Berkeley. San Mateo. Spring.

44. Rigiopappus Gray.

Superflua. Scales of the involucre linear, erect, rigid, half embracing the akene. Receptacle flat, naked. Pappus 3-5-aristate.—⊙. Flowers pale. leaves alternate.

45. Chænactis. DC.

Æqualis. Flowers of the ray tubular but frequently enlarged. Involucre hemispherical; its scales narrow. Receptacle flat. Pappus 4 to 12 obtuse, chaffy scales.—⊙.

1. *C. LANOSA* DC. Flowers yellow.—Contra Costa. Spring.

46. Blennosperma Less.

Necessaria. Ray flowers ovate, ligulate, without tube. Scales of the involucre 1-seriate, membranaceous. Receptacle flat, naked.

Akenes pyriform, papillate, gelatinous when wetted; pappus 0.—⊙. Flowers pale yellow. Leaves alternate.

1. *B. CALIFORNICUM* Torr. & Gray.—Common. Spring.

47. *Helenium* L. SNEEZE-WEED.

Superflua. Flowers of the ray palmate. Involucre 2-seriate; external scales ∞ , narrow, foliaceous, spreading, at length reflexed; internal scales few, paleaceous. Receptacle globular, naked. Pappus 5 to 12, membranaceous paleæ. Flowers of the ray yellow; disc purple; leaves alternate.

1. *H. PUBERULUM* DC.—⊙. Moist places. Summer.

Tribe VI. *ANTHEMIDEÆ*.

Receptacle rarely paleaceous. Scales of the involucre scarious. Pappus not capillary.

48. *Anthemis* L. MAY-WEED.

Superflua. Ray flowers ligulate. Scales of the involucre imbricate. Receptacle convex or conical, paleaceous. Akenes ribbed. Pappus 0. Ray-flowers white. Disc yellow.

I. *A. COTULA* L.—⊙. Common on waste grounds, although not indigenous. Native of Europe.

49. *Achillea* L. YARROW.

Superflua. Head ∞ -flowered; ray-flowers 5 to 6, ligulate, ovate. Involucre ovate; scales imbricate, those of the receptacle hyaline. Akenes marginate; pappus 0. γ .

1. *A. MILLEFOLIUM* L. Ray-flowers and disc pale.—Common. Summer.

50. *Matricaria* L. CHAMOMILE.

Superflua. *Æqualis*. Scales of the involucre imbricate. Receptacle conical, naked. Akenes angulate. Pappus coroniform or 0. Disc yellow; flowers of the ray white.— \odot

1. *M. DISCOIDEA* DC. Pappus 0; areola of akene merely surrounded by a rim; ray-flowers white, depauperate.—Summer. Common.

2. *M. OCCIDENTALIS* Greene. Pappus coroniform, unequal, the side toward the ray considerably developed, the inner side reduced to a rim.—Contra Costa. Summer.

51. *Chrysanthemum* L.

Superflua. Involucre hemispherical, spreading. Scales ∞ -seriate, imbricate, appressed. Receptacle not conical, naked. Akenes short, somewhat terete, costate, truncate at the apex; pappus 0.

1. *C. SEGETUM* L. Ray-flowers yellow.— \odot . Has been found near Oakland. Summer.

52. Soliva Ruiz & Pavon.

Necessaria. Superflua. Ray-flowers ∞ , without corolla. Style scarcely bifid, persistent. Receptacle flat, naked. Akenes obcompressed, winged, crowned by the style, sessile, pappus 0.— \odot . Flowers inconspicuous.

1. *S. SESSILIS* Ruiz & Pavon.—Moist grounds near the coast. All the year round.

53. Cotula L. BRASS-BUTTONS.

Superflua. Corolla of ray-flowers wanting. Receptacle flat, naked, papillose. Akenes compressed, winged, those of the ray stipitate; pappus 0.— \odot . Flowers yellow.

1. *C. CORONOPIFOLIA* L. Leaves pinnatifid. Aquatic. All the year round.

2. *C. AUSTRALIS* Hook. Leaves bipinnatifid; lobes linear.—Waste places. All the year round.

54. Tanacetum L. TANSY.

Æqualis. Superflua. Ray-flowers not ligulate. Receptacle convex, naked. Akenes angulate, with large epigynous disc. Pappus coroniform, or 0.— \mathcal{L} . Flowers yellow.

1. *T. CAMPHORATUM* Less. Suffrutescent. Sand dunes near San Francisco. Summer.

55. Artemisia L. WORMWOOD. SAGE-BRUSH.

Æqualis. Superflua. Necessaria. Ray flowers tubular. Scales of involucre dry, with

scarious margins. Receptacle naked or villous. Akenes obovate, with small epigynous disc; pappus 0. Heads small; flowers yellowish.

1. *A. PYCNOCEPHALA* DC. Superflua; flowers of the disc ♂, but sterile; their style not bifid; plant silky-villous; leaves pinnately parted.— $\frac{1}{2}$. Seashore. Summer.

2. *A. DRACUNCULOIDES* Pursh. Superflua. Flowers of the disc ♂, but sterile; their style not bifid; plant glabrous; leaves linear.— $\frac{1}{2}$. Lake Chabot. Summer.

3. *A. CALIFORNICA* Less. Superflua. Flowers of the disc ♂, and fertile; their style bifid; frutescent, paniculately branched, canescent; leaves pinnately parted; lobes filiform.— $\frac{1}{2}$. Common. Summer.

Branches and leaves insecticide of considerable power.

4. *A. LUDOVICIANA* Nutt. Style of the flowers of the disc bifid; involucre tomentose; leaves lanceolate; the lower frequently parted into 3-5 lobes.— $\frac{1}{2}$. Common. Summer.

Tribe VII. SENECTIONIDÆ.

Receptacle naked. Pappus capillary.

56. *Nardosmia* Cass. (*Pelasites*, Tourn.)

Monœcia, Necessaria. Heads dimorphous, ♂ or ♀ contains ♂ flowers ∞ , ♀ flowers few. ♀ or ♂ head contains ♀ or ♂ flowers few, ♀ flowers ∞ .

Involucre 1-seriate. bracteolate. Receptacle naked. Akenes glabrous, ribbed; pappus of copious capillary bristles — 2. Flowers pale, appearing before the leaves.

1. *N. PALMATA* Gray.—Taylorville. Santa Cruz Mountains. Spring.

57. *Arnica* L.

Superflua. Ray flowers elongate, with distinct, but sterile anthers. Involucre 2-seriate. Pappus capillary.—2. Flowers yellow.

1. *A. DISCOIDEA* Benth.—Santa Cruz Mountains. Summer.

58. *Senecio* L. GROUNDSEL.

Superflua. *Æqualis*. Involucre 1-seriate, bracteolate. Pappus capillary. Flowers yellow.

1. *S. HYDROPHILUS* Nutt. Glabrous; heads erect; rays few, sometimes 0; leaves almost entire; the radical, large, with long petioles; upper cauline leaves sessile, clasping.—2. Berkeley. Summer.

2. *S. ARONICOIDES* DC. Heads erect; ray-flowers few, frequently 0; leaves repand-denticulate; cauline variable, uppermost reduced to bracts.—2. Presidio. Spring.

3. *S. EURYCEPHALUS* Torr. & Gray. Rays elongated; leaves pinnately parted; lobes cuneate and acutely incised.—2. San Mateo. Summer.

4. *S. DOUGLASHI* DC. Ray-flowers elongated; leaves linear or pinnately parted into linear segments.— \mathcal{L} . Contra Costa hills. Summer.

5. *S. VULGARIS* L. Rays 0.— \odot . Waste grounds. All the year round. Weed introduced from Europe.

Tribe VIII. EUPATORIACEÆ.

Style branches club-shaped. *Æqualis*.

59. *Bulbostylis* DC. (*Brickellia* Ell.)

Scales of involucre imbricate, 2-3-seriate. Receptacle flat, naked. Margin of the tubular corolla minutely 5-dentate. Style bulbous at base. Akenes 10-striate; pappus ∞ -seriate, bristly, scabrous. Leaves opposite.

1. *B. CALIFORNICA* Gray. Suffrutescent; flowers pale.— \mathcal{L} . Niles Station. Summer.

Tribe IX. CYNAROIDEÆ.

Style branches concreted. Corolla deeply 5-cleft. Receptacle bristly.

60. *Cnicus* L. THISTLE.

Æqualis. Scales of the involucre ∞ -seriate, imbricate, ending in a spine. Pappus deciduous, ∞ -seriate; bristles plumose, connected at their base into a ring. Filaments syngenetic. $\odot\odot$.

1. *C. AMERICANUS* Gray. Involucre ovoid; bracts appressed, imbricate, with loose, scarious tips; flowers ochroleucous.—⊙⊙. Marin County. Summer.

2. *C. EDULIS* Gray. Bracts of the involucre not appressed, not rigid, but loose, tapering from a narrow base gradually into a short flaccid spine; corolla lobes filiform, with a thickened tip; flowers purple or whitish.—⊙⊙. Common. Summer.

3. *C. HALLII* Gray. Bracts of the involucre not appressed, not rigid, but loose, tapering from a narrow base gradually into a flaccid, short spine; corolla lobes linear, flat; flowers rose-color or whitish.—⊙⊙. Marin County, Summer.

4. *C. OCCIDENTALIS* Gray. Bracts of the involucre with short coriaceous base and squarrose, subulate tips; heads solitary, terminal, very large; flowers crimson. Common. Summer.

5. *C. FONTINALIS* Greene. Bracts of the involucre herbaceous, broad, reflexed, with a short spinose tip; heads nodding; flowers whitish.—Crystal Springs; very local. Summer.

6. *C. QUERCETORUM* Gray. Bracts of the involucre appressed, coriaceous, only the outer mucronate, the inner unarmed; all plane without

glandular viscid spot on the dorsal side; flowers rose, purple or whitish.—Presidio. Oakland. Summer.

7. *C. BREWERI* Gray. Bracts of the involucre bearing on their dorsal side, near the tip, an oval, viscid, glandular spot; bracts much appressed, coriaceous, outer bracts tipped with a squarrose, slender prickle; flowers pale, purple or whitish.—Tamalpais. Summer.

61. *Silybum* Vaill. SPOTTED THISTLE.

Æqualis. Pappus ∞ -seriate. Bristles rigid. Filaments monadelphous.— \odot . Flowers purple.

1. *S. MARIANUM* Gærtn.—Common. Summer.

Native of the Mediterranean region. Seeds of the plant used as an antispasmodic.

62. *Centaurea* L. STAR-THISTLE.

Frustranea. *Æqualis*. Involucre imbricate. Ray-flowers funnel-shaped. Pappus of ∞ , scabrous, filiform bristles.

1. *C. BENDICTA* L. Akenes terete; flowers yellow.— \odot .

Ballast-weed. Introduced from the Mediterranean region. Summer.

2. *C. MELITENSIS* L. Akenes costate. Flowers yellow.— \odot .

Ballast-weed. Introduced from the Mediterranean region. Summer.

Sub-Family 2. *Ligulifloræ*.

All flowers of the head ligulate.

Tribe I. CICHORIACEÆ.

63. *Stephanomeria* Nutt.

Head few-flowered. Involucre 1-seriate, calyculate. Receptacle flat, naked. Akenes 5-costate, with a callosity at their place of insertion. Pappus 1-seriate, bristly, plumose, white, coalescent.—⊙.

1. *S. VIRGATA* Benth. Flowers matutinal. Common. Summer.

64. *Rafinesquia* Nutt.

Head ∞-flowered. Involucre 1-seriate, calyculate. Receptacle flat, naked. Akenes rostrate, with a callosity at their place of insertion; pappus white, plumose, capillary, coalescent.—⊙. Flowers pale.

1. *R. CALIFORNICA* Nutt. Common. Summer.

The smell of the plant resembles that of opium.

65. *Hypochæris* L.

Involucre imbricate. Receptacle paleaceous. Paleæ deciduous. Akenes rostrate; pappus plumose. Flowers yellow.

1. *GLABRA* L.—⊙. Ballast weed, introduced from Europe. Summer.

66. (*Microseris*) *Scorzonella* Nutt.

Involucre campanulate, imbricate in several ranks. Receptacle flat, naked, foveolate. Akenes more than 5-costate; pappus paleaceous; paleæ entire, lanceolate, tipped with a barbellate awn. Heads nodding when in bud. Plant caulescent. Flowers yellow.

1. *S. PROCERA* Gray. Head more than 100-flowered; awns simply barbellate, 3 times longer than the paleæ.—2f. Sonoma. Summer.

2. *S. PALUDOSA* Greene. Head less than 75 flowered; awns simply barbellate, twice as long as the paleæ.—2f. Marshy grounds. Marin County near Corte Madera. Summer.

3. *S. SYLVATICA* Gray. Awns nearly plumose, shorter than the paleæ.—2f. Contra Costa. Summer.

67. (*Microseris*) *Calais* DC.

Involucre conical, imbricate. Receptacle flat, alveolate. Akenes more than 5-costate; paleæ of the pappus 5, flat, bifid, with short awns. Head always erect on hollow, scape-like peduncles, that swell toward the head. Flowers yellow.

1. *C. KELLOGGII* Greene. Awn of the pappus longer than the palea; palea emarginate (not deeply cleft).—☉. San Bruno Mountains; Marin County. Spring.

2. *C. LINDLEYI* DC. Awn of the pappus somewhat shorter than the palea; palea brownish, persistent.—☉. Common. Spring.

3. *C. LINEARIFOLIA* DC. Awn of the pappus not longer than the palea; palea white, persistent. ☉.—Common. Spring.

68. (*Microseris*) **Microseris** Don.

Involucral bracts imbricate, the outer rank very short, calyculate. Receptacle flat, slightly alveolate. Paleæ of the pappus entire, tapering into a scabrous awn. Capitula nodding on slender scapes, which are not thickened towards the head. Flowers yellow.—☉.

1. *M. DOUGLASSII* Gray. Pappus paleaceous; paleæ 5, boat-shaped, tapering abruptly into an awn of the length of the akene.—Common. Spring.

2. *M. ATTENUATA* Greene. Pappus paleaceous; paleæ 5, boat-shaped, tipped with an awn shorter than the akene.—Contra Costa. Spring.

3. *M. ACUMINATA* Greene. Pappus paleaceous; paleæ 5, flat and straight, gradually tapering into an awn, shorter than the palea. Contra Costa. Spring.

4. *M. BIGELOVII* Gray. Pappus paleaceous; paleæ 5, flat and straight, passing into an awn more than twice the length of the palea.—Common. Spring.

5. *M. ELEGANS* Greene. Pappus paleaceous; paleæ deltoid, very short, with a slender awn twice their length.—Niles. Spring.

6. *M. APHANTOCARPHA* Gray. Pappus paleaceous; paleaceous part nearly obsolete, aristate part long but setaceous, slender, and very fragile.—San Francisco. Spring.

69. *Malacothrix* DC.

Receptacle, flat, naked. Akenes truncate; pappus 1-seriate, bristles scabrous, silvery, deciduous in a ring.—⊙.

1. *M. CALIFORNICA* DC. Scapose; involucre bracts in more than 2 ranks; flowers yellow. Contra Costa. Spring.

2. *M. CLEVELANDI* Gray. Branching; involucre bracts in 2 ranks of equal length; besides the regular caducous pappus an external accessory pappus of 1 persistent bristle and a dentate crown; flowers yellow.—Contra Costa. Spring.

3. *M. OBTUSA* Benth. Branching; involucre bracts in two ranks of unequal length; persistent bristles 0; flowers white.—Marin County. Spring.

70. Hieracium FOURN. HAWKWEED.

Involucre ∞ -seriate, imbricate. Akenes terete, with a thin, crenulated ring on the apex, pappus 1-seriate, capillary.— \mathcal{L} .

1. *H. ALBIFLORUM* Hook. Involucre ∞ -flowered; akenes linear; pappus sordid; flowers white.—Saucelito. Summer.

71. Troximon Nutt.

Involucre imbricate. Akenes linear. Apex contracted. Pappus ∞ -seriate. Bristles silky, not deciduous in a ring.

1. *T. APARGIOIDES* Less. Beak of the akene about the length of body; flowers yellow.— \mathcal{L} . San Francisco. Summer.

2. *T. HUMILE* Gray. Beak of the akene several times the length of body; pappus as long as the beak; flowers yellow.— \mathcal{L} . Common. Summer.

3. *T. LACINIATUM* Gray. Beak of the akene several times the length of the body; pappus much shorter than the beak; leaves laciniate—when more deeply parted and pinnatifid, the lobes linear; flowers yellow.— \mathcal{L} . Marin County. Summer.

4. *T. GRANDIFLORUM* Gray. Beak of the akene several times the length of body; pappus much shorter than the beak; leaves runcinate

—when more deeply parted and pinnatifid, the lobes not linear; flowers yellow.— \mathcal{L} . Common. Summer.

5. *T. HETEROPHYLLUM* Greene. Beak of the akene but little longer than body; flowers yellow.— \odot . Common. Spring.

72. *Taraxacum* Haller. DANDELION.

Involucre imbricate. Akenes contracted into a long beak; pappus ∞ -seriate; bristles white, capillary nearly persistent.— \mathcal{L} . Flowers yellow.

1. *T. OFFICINALE* Weber. Ballast weed introduced from Europe; rare. Summer. Official under the old Linnean name. *Leontodon Taraxacum*.

73. *Sonchus* L. SOW-THISTLE.

Involucre ∞ -seriate, imbricate. Akenes compressed, truncate. Pappus ∞ -seriate, capillary. Hairs in fascicles, soft. Flowers yellow.

1. *S. OLERACEUS* L. Auricles of the cauline leaves acute.— \odot . Common. All the year round. Ballast weed from Europe.

2. *S. ASPER* Vill. Auricles of the clasping cauline leaves rounded.— \odot . Ballast weed from Europe.

Family 2. CAMPANULACEÆ.

Calyx superior, 5-cleft. Stamens 5, alternating with the divisions of the gamapetalous corolla, and inserted between the corolla and the ovary. Anthers introrse, filaments flattened. Ovary 2-5 celled; cells ∞ -ovulate. Placentæ central. Style 1. Stigmas 2-5. Fruit capsular. Leaves alternate.—Lactescent herbs.

1. Githopsis Nutt.

Calyx tube clavate, 10-costate; with 5 narrow, persistent lobes. Corolla campanulate, 5-lobed. Filaments 5, membranaceous. Ovary 3-celled. Stigmas 3. Capsule clavate, 10-costate, enclosed by the calyx, opening by a round hole at the apex. Seeds ∞ .— \odot . Dwarfish.

1. *G. SPECULARIODES* Nutt. Flowers blue. Marin County. Contra Costa. San Francisco. Spring.

2. Specularia Heister. VENUS'S LOOKING-GLASS.

Calyx-tube elongated. Lobes narrow. Corolla rotate, 5-lobed. Filaments membranaceous. Capsule elongated, opening by parietal valves near the apex. Seeds ∞ .— \odot . Flowers blue.

1. *S. BIFLORA* Gray.—Marin County. Contra Costa. Spring.

3. **Campanula** Tourn. BELL-FLOWER.

Calyx tube turbinate. Corolla campanulate. Filaments membranaceous at their base. Capsule turbinate, opening by parietal valves. Flowers blue.

1. *C. EXIGUA* Rattan. Stem-leaves linear.—⊙. Tamalpais. Spring.

2. *C. PRENANTHOIDES* Durand. Stem-leaves ovate, serrate.—2f. Marin County redwoods. Spring.

4. **Heterocodon** Nutt.

Early flowers cleistogamic; later ones expanding. Calyx turbinate. Corolla campanulate. Capsule membranaceous, turbinate, opening irregularly.—⊙. Flowers blue.

1. *H. RARIFLORUS* Nutt.—Marin County. Spring.

Family 3. **CUCURBITACEÆ.**

Monœcious or diœcious. ♂ : Stamens 5, 3-adelphous. Cells of anthers contorted, flexuous. ♀ : ovary 2-5-celled. Placentæ parietal. Style 1. Stigmas 3.

1. **Megarrhiza** Torr. (*Echinocystis*. Torr. & Gray.)
BIG-ROOT.

Monœcious. ♂ flowers racemose. ♀ flowers single from the axil of the ♂ raceme. Limb of calyx minutely 5-dentate, Corolla rotate,

deeply 5-lobed. Fruit a berry. Seeds large, globose.— \mathcal{L} . Root large; stem climbing by tendrils. Flowers greenish white.

1. *M. FABACEA* Naud. (*Californica* Torr.) Ovary globose, densely echinate; fruit globose, covered with pungent spines; seeds globose.—San Francisco. Spring.

2. *M. MARAH* Kellogg. Ovary ovate, covered with soft spines; fruit ovate, muricate; seeds lenticular.—San Francisco. Spring.

The enormous rhizomes of both species abound in an acrid, drastic juice, which renders the fecula, contained in a fair proportion, poisonous.

Order 2. RUBIALES. Filaments terete.

Family 1. DIPSACEÆ.

Leaves opposite. Stipules 0. Flowers in involucrate heads; single flowers protected by a double (involucellate) calyx, tetrandrous. Ovary 1-celled, 1-ovulate. Fruit utricular.

1. *Dipsacus* Tourn. TEASEL.

Head oblong. Leaflets of the involucre radiate. External calyx (involucellum) quadrangular; internal cyathiform.— $\odot\odot$. Spring.

1. *D. FULLONUM* L. Flowers pale.—Cultivated grounds. Summer. Escaped from cultivation, Originally Mediterranean.

Family 2. VALERIANACEÆ.

Leaves opposite. Stipules 0. Inflorescence cymose. Calyx pappus-like. Ovary 3-celled; 2 cells sterile, and the fertile one 1-ovulate. Fruit an akene.

1. Plectritis. DC.

Limb of calyx straight, entire, obsolete. Tube of the corolla gibbous; limb 5-cleft, 2-labiate; triandrous. Akene winged by the persistent, sterile cell.—☉.

1. *P. MACROCERA* Gray. Corolla almost regular; akene semilunar; dorsal angle of akene obsolete; flowers white or rose-color. Common. Spring.

2. *P. CONGESTA* Lindl. Corolla distinctly 2-labiate; dorsal angle of akene distinctly carinate; flowers rose-color.—Marin County. Spring.

Family 3. CAPRIFOLIACEÆ.

Leaves opposite. Stipules 0. Fruit baccate.

1. Sambucus Tourn. ELDER.

Limb of Calyx 5-dentate. Corolla rotate, regular. Ovary 3-5-celled; cells 1-ovulate: styles 9; stigmas 3-5. Fruit a 3-5-seeded berry.

1. *S. GLAUCA* Nutt. Flowers cream-colored; berries blue, pruinose.—½. Berries eatable; root a powerful sudorific,

2. *Symphoricarpus* Cass. SNOWBERRY.

Calyx-limb 5-4-dentate, persistent. Corolla funnel-shaped, 5-4-lobed. Ovary 4-celled; 2 cells ∞ -ovulate but abortive, alternating with 1-ovulate and fertile. Style filiform. Berry 4-celled, 2-seeded.— $\frac{1}{2}$.

1. *S. RACEMOSUS* Michx. Glabrous; leaves oval; corolla narrowed at base; flowers pinkish; berries white.—Common. Summer.

2. *S. MOLLIS* Nutt. Pubescent; leaves orbicular; corolla with broad base; flowers pinkish; berries white.—Common. Summer.

3. *Lonicera* L. HONEYSUCKLE. TWIN-BERRY.

Calyx-limb small, 4-dentate. Corolla tubular or campanulate. Limb irregular 5-cleft. Style filiform. Berry 3-celled; cells few-seeded.

1. *L. INVOLUCRATA* Banks. Flowers in pairs; bracts foliaceous, bractlets conspicuous and accrescent in fruit; flowers orange-colored; berries shining black, enveloped in dark red, involucrate bracts. Stem erect.— $\frac{1}{2}$. Common. Summer. "Twin-berry"

2. *L. HISPIDULA* Dougl. Flowers in whorls round an axis, forming a loose spike; uppermost leaves connate; flowers in different shades of red and yellow; berries orange-red. Stem not erect.— $\frac{1}{2}$. Common. Summer.

Family 4. RUBIACEÆ.

Leaves opposite or verticillate. If opposite, stipulate.

1. Cephalanthus L. BUTTON-BUSH.

Flowers capitate. Limb of the calyx 4-dentate. Corolla tubular, slender, with 4-cleft limb. Stamens 4, inserted in the throat of the corolla. Style 1, exserted. Stigma capitate. Ovary 2-4 celled. Cells 1-ovulate.— $\frac{1}{2}$.

1. *C. OCCIDENTALIS* L. Flowers white; rare. Niles Station. Summer.

2. Galium L. BED-STRAW. CLEAVERS.

Limb of the calyx entire, obsolete. Corolla 4-3 cleft, rotate. Ovary 2-celled. Cells 1-ovulate. Styles 2, connate at base, capitate at apex. Fruit sometimes fleshy. Stems quadrangular. Leaves verticillate.

1. *G. APARINE* L.—♀ Fruit a dicocous schizocarp, granulate and setaceous; leaves 6-8 in a whorl; stem retrorsely bristly; flowers white.—☉. Common. Summer.

2. *G. BOREALE* L.—♀. Fruit a dicocous schizocarp; leaves 4 in a whorl, 3-nerved, not cuspidate; flowers white.—♂. Tamalpais. Summer.

3. *G. TRIFIDUM* L.—♀. Fruit a dicocous schizocarp; leaves 4-6 in a whorl, 1-nerved, not cuspidate; flowers small, white.—♂. Marin County. Summer.

4. *G. ASPERRIMUM* Gray.—♀ Fruit dicocious, a scabrous schizocarp; leaves 4–6 in a whorl, cuspidate; cymes dichotomous; flowers white.—℥. Common. Summer.

5. *G. TRIFLORUM* Michx.—♀ Fruit dicocious, an uncinatè hispid schizocarp; leaves 4–6 in a whorl, cuspidate; flowers in 3-florous or biternate cymes, whitish.—℥.

In shady, moist places; Marin County. San Mateo. Summer. This plant developes, in withering, the fragrance peculiar to the European *Asperula odorata*, the *Waldmeister* of the Germans. It is used frequently for the so-called “May-drink.”

6. *G. CALIFORNICUM* Hook & Arn. Dioecious. Fruit baccate; leaves thinnish, ovate, acuminate, 4 in a whorl; flowers yellowish.—℥. San Francisco. San Mateo. Contra Costa. Summer.

7. *G. NUTTALLII* Gray. Dioecious. Fruit baccate; leaves coriaceous, small, oval, obtuse, mucronulate, 4 in a whorl; flowers yellowish; ♀.—Common. Summer.

8. *G. ANDREWSII* Gray. Dioecious. ♂ flowers in a few flowered cyme; ♀ flowers solitary; fruit baccate; leaves crowded, acerose subulate; 4 in a whorl.—℥. Common. Summer.

Sub-Section 2. HYPOGYNÆ.

ORDER 1. GENTIANALES. Tendency to develop an apocarpus ovary either by separating the two cells entirely or by making the capsule septicidous.

Family 1. ASCLEPIADACEÆ.

Monadelphous. Gynandrous. Anthers ex-trose; their pollen changed into pollinaria. Ovaries 2. Styles 2, united into a pentagonal stigma. Fruit a pair of follicles, or by abortion a single one. Lactescent plants.

1. *Asclepias* L. MILK-WEED.

Corolla rotate, 5-parted. Stamineal tube short, expanding into a corona of 5 cucullate processes, from the cavity of each of which rises a corniform process. Cells of anthers separated, each connected to the cell of the neighboring one by an apical appendage, the pollinia becoming suspended over a gland, five of which protrude from the margin of the stigma. Seeds ∞ , with a silky arillus.— γ .

1. *A. SPECIOSA* Torr. Follicles echinate; flowers purplish.—Marin County. Contra Costa. Laundry Farm. Summer.

2. *A. VESTITA* Hook. & Arn. Follicles glabrous; erect on a deflexed pedicel; umbels ∞ -flowered, terminal umbel pedunculate; lateral

ones sessile; flowers whitish or purplish.—Niles Station. Summer.

3. *A. MEXICANA* Cav. Follicles glabrous; erect on erect pedicels; umbels ∞ -flowered; peduncles longer than the pedicels; flowers whitish.—Alameda on marshy places. Summer.

Family 2. APOCYNACEÆ.

Stamens not monadelphous. Anthers introse. Corolla contorted in æstivation and its lobes oblique. Lactescent.

1. *Apocynum* Tourn. DOG-BANE. INDIAN HEMP.

Corolla campanulate, 5-cleft, with 5 scales opposite the lobes and near their base. Stamens inserted on the base of the corolla. Filaments very short. Anthers sagitate, conniving into a ring. Ovaries 2, with 5, hypogynous scales. Fruit 2-follicular. Seeds with a hairy arillus.—2f.

1. *A. ANDROSÆMIFOLIUM* L. Cymes loose; corolla campanulate; tube longer than the calyx-lobes; corolla lobes revolute; flowers flesh-color.—Livermore. Summer. Rare. (Dog-bane.)

2. *A. CANNABINUM* L. Cymes dense, compact; corolla tube not longer than the calyx-lobes; corolla lobes almost erect; flowers pale. Alvarado, on marshy places; very local. Sum-

mer. (Indian hemp.) The strong, silky fiber of the stem has been recommended as a textile substance, comparable to hemp.

Family 3. GENTIANACEÆ.

Anthers introrse. Lobes of the corolla not oblique. Ovary ∞ -ovulate, septicidal. Not lactescent.

1. *Microcala* Link. (*Cicendia* Adans.)

Calyx costate, 4-dentate. Corolla salver-shaped. Withering anthers not spirally contorted. Style filiform. Stigma peltate or 2-lobed.—☉. Dwarfish herbs. Flowers yellow.

1. *M. QUADRANGULARIS* Gries. Saucelito. Spring.

2. *Erythræa* Pers. CANCHALAGUA.

Calyx tubular, angular, 5-cleft. Corolla funnel-shaped, withering, but not deciduous. Withering anthers spirally contorted. Style filiform, deciduous. Capsule 1-celled, half divided by the prominent parietal placentæ. ☉. Flowers pink.

1. *E. FLORIBUNDA* Benth. Anthers oblong; lobes of corolla lanceolate; seeds globular. San Bruno range. Spring.

2. *E. MUHLENBERGII* Gries. Anthers oblong; lobes of corolla very obtuse; seeds oval. San Miguel. Spring.

3. *E. DOUGLASHII* Gray. Anthers oblong; lobes of corolla obtuse; seeds globular.—San Francisco. Spring.

4. *E. TRICHANTHA* Gries. Anthers linear; lobes of corolla narrow, lanceolate.—Marin County. Spring.

5. *E. VENUSTA* Gray. Anthers linear; lobes of corolla oval.—San Leandro. Spring.

3. *Gentiana* L. GENTIAN.

Corolla campanulate 4-5 lobed, withering but not deciduous. Style short or 0. Stigma 2-parted, persistent. Capsule septicidal.—♀.

1. *G. OREGANA* Engelm. Flowers blue.—Tamalpais. Very local. Autumn.

4. *Menyanthes* TOURN. BUCK-BEAN.

Corolla funnel-shaped, induplicate in æstivation, deciduous. Limb 5-lobed, bearded on the upper surface.—♀. Aquatic. Flowers white.

1. *M. TRIFOLIATA* L. San Francisco. Summer. (Extinct since 1859.)

ORDER 2—DIANDRÆ. Corolla regular. Gamopetalous, 4-5 cleft or 0. Stamens 2. Ovary 2-celled; ovules in each cell 1-3. Leaves opposite.

Family 1. OLEACEÆ.

Æstivation valvate.

1. *Fraxinus* Tourn. ASH.

Flowers diclinous. Calyx 2-4 parted or 0. Corolla 2-4 parted; divisions 2-4 or 0. Ovary 2-celled; cells 2-ovulate. Stigma sessile, 2-cleft. Fruit a samara.— $\frac{1}{2}$.

1. *F. DIPETALA* Hook. & Arn. Petals 2.
 $\frac{1}{2}$.—Livermore. Niles. Spring.

2. *F. OREGANA* Nutt. Dioecious, apetalous.
 $\frac{1}{2}$.—Near creeks. Marin County. Menlo Park. San Mateo. Niles. Spring.

Infusion of leaves has formerly been used at the Russian settlements against rheumatism.

ORDER 3—LAMIALES. Corolla 2-labiate. Number of stamens less than lobes of the corolla. Ovaries 2-4 or ovary 2-4 celled. Cells 1-2 or 4-ovulate. Fruit an akene, drupe or capsule. Leaves opposite, Stipules 0.

Family 1. LABIATÆ.

Ovary 4-cleft, 4-ovulate, inserted on a hypogynous disc. Style 1. Fruit 4 akenes.

1. *Trichostema* Gronov. BLUE CURLS.

Tube of the corolla slender, limb 5-cleft. Lobes oblong, declined. Stamens didynamous, long-exserted. Filaments spiral in the bud.

1. *T. LANCEOLATUM* Benth. Flowers blue.
⊙.—Niles Station. Livermore. Dry river beds. Summer.

2. *Mentha* L. MINT.

Calyx 5-dentate. Tube of corolla included. limb exerted, 4-lobed; upper lobe the largest. Stamens 4, scarcely didynamous, erect, distant. Anthers 2-celled. Cells parallel.—♂.

1. *M. VIRIDIS* L. Inflorescence terminal; dense glomerules, crowded around narrow spikes; spikes leafless; leaves almost sessile. Irrigated grounds. Summer. Autumn. Escaped from cultivation. Native of Europe. (Spearmint.)

2. *M. PIPERITA* L. Inflorescence terminal; glomerules on uninterrupted narrow spikes; spikes leafless; leaves distinctly petioled; flowers pale.—Irrigated grounds. Summer. Autumn. Escaped from cultivation. Native of Europe. (Peppermint.)

3. *M. CANADENSIS* L. Inflorescence axillary; uppermost axils of leaves without flowers.—Wet places. Native species. Summer.

3. *Lycopus* TOURN. WATER HOREHOUND.

Diandrous. Posterior pair of stamens 0 or sterile. Otherwise as *Mentha*.—♂.

1. *L. LUCIDUS* Turcz. Flowers pale—Marin County. Aquatic. Summer.

4. *Pycnanthemum* Benth. MOUNTAIN MINT.

Corolla tube as long as the calyx, limb 2-labiate, upper lip nearly entire. Lower lip

3-lobed. Lobes obtuse. Stamens 4, straight, divergent. Cells of anthers parallel.— γ .

1. *P. CALIFORNICUM* Torr. Flowers pale. Contra Costa. Summer.

5. *Monardella* Benth.

Limb of corolla 2-labiate. Upper lip 2-cleft. Lower lip 3-cleft. All the lobes narrow. Stamens 4, straight, divergent. Cells of anthers at last divergent or divaricate.

1. *M. VILLOSA* Benth. Flowers capitate; leaves ovate, pinnately veined. Bracts ovate, foliaceous; flowers purplish.— γ . Marin County. Summer.

2. *M. DOUGLASII* Benth. Flowers capitate; leaves lanceolate; bracts ovate, pinnately veined, membranaceous, tapering gradually into a cusp; flowers purplish.— \odot . Marin County. Summer.

3. *M. BREWERI* Gray. Flowers capitate. leaves ovate, pinnately veined, abruptly cuspidate; flowers purplish.— \odot . Contra Costa. Summer.

4. *M. LANCEOLATA* Gray. Flowers capitate; leaves lanceolate; bracts nerved, with veinlets crossing from nerve to nerve; flowers rose-color.— \odot . Contra Costa. Summer.

5. *M. UNDULATA* Benth. Flowers capitate; leaves spatulate, obtuse, undulate-margined; bracts ovate, nerved, without veinlets crossing

from nerve to nerve; flowers purplish.—⊙.
Common. Summer.

6. *Micromeria* Benth. YERBA BUENA.

Calyx tubular, 13-nerved, 5-dentate. Corolla 2-labiate. Stamens didynamous, ascending.—2f.

1. *M. DOUGLASII* Benth. Flowers white. Common. Summer.

Formerly much in use as a carminative.

7. *Pogogyne* Benth.

Calyx campanulate, 15-nerved, 5-cleft. 2 lower teeth much longer than the three upper ones. Corolla 2-labiate. Stamens didynamous, ascending, convergent in pairs. Style villous.—⊙.

1. *P. DOUGLASII* Benth. All stamens fertile; inflorescence taking the shape of a continuous spike; lower calyx lobes twice the length of the calyx-tube; flowers violet.—Common. Summer.

2. *P. PARVIFLORA* Benth. All stamens fertile; inflorescence taking the shape of a continuous spike; lower calyx lobes about the length of the calyx-tube; flowers violet.—Berkeley hills. Summer.

3. *P. SERPYLLOIDES* Gray. Posterior pair of stamens sterile; inflorescence taking the shape of an interrupted spike; flowers violet. Berkeley. Summer.

8. *Acanthomintha* Gray.

Calyx tubular 13-nerved, naked in the throat. Upper lip 3-dentate. Lower lip 2-parted, shorter. Teeth spinulose. Corolla tube exceeding the calyx, naked. Upper lip curved, entire or 2-lobed. Lower lip 3-lobed. Upper pair of stamens sterile and shorter. Lower pair ascending. Anthers 2-celled. Cells divaricate. Bracts thorny.—⊙.

1. *A. LANCEOLATA* Curran. Flowers rose-color.—Niles. Summer.

9. *Sphacele* Benth.

Calyx campanulate, enlarged in fruit, 10-nerved, reticulate-veined, 5-dentate. Tube of the corolla wide, with a hairy ring near its base, the 5 lobes of the limb obtuse. Lower lobes the longer ones. Stamens ascending. Cells of the anthers divergent.—½. Flowers large, pale.

1. *S. CALYCINA* Benth.—San Mateo. Summer.

10. *Salvia* L. SAGE.

Calyx 2-labiate. Corolla ringent. Stamens 2, parallel. Anthers separated by a connective, shaped like a branch of the filament; only the posterior end of it bearing an anther cell.

1. *S. CARDUACEA* Benth. White-woolly with cobwebby hair; radical leaves sinuate to

pinnatifid, spinulose; upper lip of calyx 3-dentate; flower lavender-color.—⊙. Livermore. Spring.

2. *S. COLUMBARIA* Benth. Puberulent; radical leaves pinnately parted; partitions crenate, rugose; upper lip of calyx 2-dentate, teeth connivent; flowers blue.—⊙. Niles station. Spring.

11. *Audibertia* Benth.

Calyx 2-labiate. Corolla 2-labiate; upper lip 2-cleft, spreading. Stamens ascending. Anthers dimidiate, (upper half of the apparent filament showing by an articulation or spur its being originally a connective between the 2 cells of an anther.)

1. *A. GRANDIFLORA* Benth. Leaves hastate, sticky; flowers crimson.— $\frac{1}{4}$. $\frac{1}{2}$. Bernal Hights. Summer.

2. *A. STACHYOIDES* Benth. Leaves lanceolate; flowers violet.— $\frac{1}{2}$. Lake Chabot. Summer.

12. *Lophanthus* Benth. GIANT HYSSOP.

Calyx subregular, 15-nerved, 5-dentate. Upper pair of stamens longer and declined; lower pair shorter and ascending. Anther 2-celled; cells parallel.— $\frac{1}{4}$. Flowers purplish.

1. *L. URTICIFOLIUS* Benth.—Marin County. Summer.

13. *Scutellaria* L. SKULLCAP.

Calyx 2-labiate, closed in fruit, at last splitting and the upper lip deciduous. Tube of corolla exerted; upper lip crest-shaped, with 2 lateral lobes; lower lip reduced to a single lobe. Stamens didynamous, ascending, parallel. Anthers of the lower pair dimidiate.

1. *S. TUBEROSA* Benth. Root stock filiform, bearing tubers; none of the floral leaves shorter than the flowers; flowers blue.— \mathcal{L} . San Francisco. Contra Costa. Spring.

2. *S. CALIFORNICA* Gray. Root stock filiform, but without tubers; uppermost floral leaves shorter than the flowers; flowers pale.— \mathcal{L} . Marin County. Spring.

14. *Prunella* L. SELF-HEAL.

Calyx 2-labiate, closed in fruit and flattened. Corolla with a scaly or hairy ring near its base; upper lip concave; lower lip 3-lobed. Stamens didynamous, ascending. Anthers approximate in pairs.— \mathcal{L} .

1. *P. VULGARIS* L. Flowers blue, — San Francisco. Summer.

15. *Marrubium* L. HOREHOUND.

Calyx tubular, 5-10 dentate. In fruit the teeth spreading. Tube of the corolla included. Limb 2-labiate. Stamens didynamous; included. Cells of anthers divaricate.— \mathcal{L} .

1. *M. VULGARE* L. Flowers white.—San Francisco. Contra Costa. Summer.

Native of Europe. Perhaps in older times used as a medicinal herb.

16. *Stachys*. HEDGE-NETTLE.

Limb of the corolla 2-labiate, upper lip erect, concave, lower lip spreading, 3-lobed, middle lobe longest. Stamens ascending, approximate in pairs, parallel, contorted in withering.

1. *S. AJUGOIDES* Benth. Corolla pale, tube about the length of the calyx; leaves oblong with rounded base; lower leaves petioled; upper sessile; none of the floral leaves shorter than the clusters, which are arranged in an interrupted spike.— \mathcal{L} . Common. Summer.

2. *S. ALBENS* Gray. Corolla pale, tube about the length of the calyx; leaves oblong with cordate base; lower leaves short-petioled; upper nearly sessile; most of the floral leaves shorter than the inflorescences, which are arranged in an interrupted spike.— \mathcal{L} . Santa Clara County. Summer.

3. *S. PYCNANTHA* Benth. Corolla pale; tube about the length of the calyx; floral leaves all reduced to small bracts.— \mathcal{L} . Common. Summer.

4. *S. BULLATA* Benth. Corolla purple; tube somewhat longer than the calyx.— ζ . Common. Summer.

5. *S. CHAMISSONIS* Benth. Corolla red; tube twice as long as the calyx.— ζ . Presidio. Saucelito. Wild Wood Glen. Point Bonita. Summer.

Family 2. **VERBENACEÆ.**

Didynamous. Ovary undivided. Fruit drupaceous or baccate, or splitting at length into its component parts (nutlets.)

1. *Verbena* L.

Calyx 5-cleft. Corolla salver-shaped. Fruit at length splitting into four 1-seeded nutlets.

1. *V. OFFICINALIS* L. Flowers pale.— \odot . Not common. Summer.

2. *Lippia* L.

Fruit 2-celled, splitting into two 1-seeded nutlets.

1. *L. NODIFLORA* Michx. Flowers rose-color. \odot . Niles. Summer.

Family 3. **PLANTAGINEÆ.**

Calyx 4-sepalous, persistent. Corolla gamopetalous, 4-cleft, scarious. Stamens 4, alternate with the lobes of the corolla. Anthers versatile. Style.1.

1. *Plantago* L. PLANTAIN.

Flowers bracteate, spicate. Ovary 2-celled (spuriously 4-lobed.) Fruit a capsule, transversally dehiscent.

1. *P. MAJOR* L. Corolla glabrous; leaves 5-9 nerved; cells of ovary containing several ovules.—☿. Everywhere. Summer.

Popular remedy with the old inhabitants. Leaves used as external application; supposed to cool when put to a sore with their under-side, and to draw when put there by their upper side.

2. *P. MARITIMA* L. Corolla tube externally pubescent; leaves linear, fleshy.—☿. On rocks near the sea. Summer.

3. *P. LANCEOLATA* L. Corolla glabrous; leaves 3-5 nerved; cells of ovary each containing 1 ovule.—⊙⊙. San Mateo. Summer.

4. *P. PATAGONICA* Jacq. Entire inflorescence silky pubescent; leaves 1-3-nerved; cells of ovary each containing 1 ovule.—⊙. Everywhere. Spring.

5. *P. HIRTELLA* HBK. Dioecious; scape and inflorescence hirsute; leaves 5-7 nerved; cells of ovary each containing 2 ovules.—☿. Ocean Lake. Berkeley. Summer.

An infusion of the leaves is much recommended by old inhabitants as a gargle in diphtheria.

6. *P. BIGELOVII* Gray. Polygamous, diandrous; spikes short, thick, dense; leaves filiform, fleshy.—☉. Salt marshes. Summer.

7. *P. HETEROPHYLLA* Nutt. Dioecious; spike slender; leaves fleshy.—☉. Sandy ground. Spring.

ORDER 4. PERSONALES. Corolla gamopetalous, irregular, its lobes not corresponding in number to the number of stamens which are inserted into its tube. Ovary superior, consolidated from two seed-leaves. Ovules ∞ . Fruit a capsule or berry. Stipules 0.

Family 1. OROBANCHEÆ.

Ovary 1-celled; placentæ parietal. Parasites without chlorophyll. Leaves reduced to scales.

1. *Anoplanthus* Endl. (*Aphyllon*).

Flowers without bractlets. Calyx 5-cleft. Corolla tubular, curved; limb 5-cleft. Stamens didynamous; cells of anthers separated, mucronate at their base. Placentæ 4. Capsule 1-celled, dehiscent by 2 valves; each valve with a placenta on each margin.—24.

1. *A. UNIFLORUS* L. Rhizome bearing only a few scapes; calyx lobes longer than the calyx-tube; flowers dull yellow, more or less tinged with violet; fragrant.—Saucelito. Rare. Spring.

2. *A. FASCICULATUS* Nutt. Rhizome bearing fasciculate scapes (peduncles); calyx lobes not longer than the calyx tube and much shorter than the corolla; flowers dull yellow, more or less tinged with purple.—Saucelito. Rare. Summer.

2. *Aphyllon* Mitchell.

Flowers with bractlets. Otherwise nearly as the preceding.—2f.

1. *A. COMOSUM* Gray. Paniculate-racemose; bractlets remote from calyx; calyx half the length of corolla; flowers pink.—Rare. Summer.

Family 2. **SCROPHULARIACEÆ.**

Ovary 2-celled. Placentæ axillary or central. Capsule 2-celled, with 2 or 4 valves,

Tribe 1. **RHINANTHEÆ.**

Lower half of corolla in æstivation covering the upper half. Corolla tubular; limb 2-labiate.

1. *Pedicularis* Tourn.

Didynamous. Corolla ringent; upper lip laterally compressed. Anthers of equal size and insertion. Capsule loculicidal,—2f.

1. *P. DENSIFLORA* Benth. Flowers crimson.—Hill-sides, Spring.

2. *Cordylanthus* Nutt.

Calyx spathaceous; fissure, if present, lateral; lips of corolla short; upper one laterally compressed. Stamens didynamous or diandrous. Anthers 2-celled; cells separated and of different shape and insertion. Capsule loculicidal.—☉.

1. *C. MARITIMUS* Nutt. Calyx 1-phyllous; flowers sessile, axillary; anthers of the two longer stamens 2-celled; anthers of the two shorter stamens dimidiate; leaves and bracts entire, pale green; corolla purplish.—Salt marshes, San Francisco. Summer.

2. *C. MOLLIS* Gray. Calyx 1-phyllous; flowers sessile, axillary, diandrous; anthers 2-celled; leaves and bracts linear, hirsute, the upper ones sometimes dentate, laciniate; flowers pale.—Salt marshes. Vallejo. Summer.

3. *C. PILOSUS* Gray. Didynamous. Calyx 2-phyllous; all anthers 2-celled; leaves and bracts soft, villous; flowers yellowish or purplish.—Santa Clara County, on dry grounds. Spring.

4. *C. FILIFOLIUS* Nutt. Didynamous. Calyx 2-phyllous, all anthers 2-celled; floral leaves hispid, flowers purplish.—Santa Cruz mountains. Spring.

3. *Orthocarpus* Nutt.

Calyx spathaceous, cleft vertically. Corolla personate, upper lip the smaller one, lower lip saccate. Didynamous; cells separated and of different shape and insertion. Capsule loculicidal.—☉.

1. *O. LITHOSPERMOIDES* Benth. Lower lip 3-saccate; sacs ample, ventricose, much larger than the upper lip; upper lip straight, slender, subulate, anther 2-celled; corolla white, tinged with purple.—Ocean Lake. Colma. Spring.

2. *O. FAUCIBARBATUS* Gray. Lower lip 3-saccate, much larger than the straight slender upper lip; anthers 1-celled; stamens wrapt in the involute upper lip; plant glabrous, only bracts puberulent; corolla pale.—Contra Costa. Spring.

3. *O. ERIANTHUS* Benth. Lower lip 3-saccate, much larger than the straight, slender upper lip; anthers 1-celled; stamens wrapt in the involute upper lip; plant pubescent; corolla yellow; in a variety (hybrid?) white or rose-color, but upper lip always dark.—Marin County. Spring.

4. *O. FLORIBUNDUS* Benth. Lower lip 3-saccate, much larger than the straight slender upper lip; anthers 1-celled; stamens not wrapt in the lanceolate upper lip; corolla white, tube twice the length of the calyx.—Marin County. Millbrae. Spring.

5. *O. PUSILLUS* Benth. Lower lip 3-saccate, larger than the slender, straight upper lip; anthers 1-celled; stamens not wrapt in the lanceolate upper lip; corolla purplish, its tube not surpassing the calyx.—Common. Spring.

6. *O. PURPURASCENS* Benth. Lower lip simply saccate, broader but not longer than the upper lip, the tip of which is incurved and back bearded; bracts and corolla white, rose or crimson.—Common. Spring.

7. *O. CASTILLEIODES* Benth. Lower lip ventricose; upper lip straight, its back not bearded; leaves lanceolate, commonly laciniate; upper leaves not attenuate, but cuneate-dilated and deeply cleft; spike dense; bracts and corolla pale, sometimes sprinkled with red.—Marsh near Tamalpais. Spring.

8. *O. DENSIFLORUS* Benth. Lower lip ventricose; upper lip straight, its back not bearded; leaves linear; upper leaves attenuate; bracts 3-cleft, as long as the flowers; spike dense; bracts and corolla purple and white.—Common. Spring.

9. *O. ATTENUATUS* Gray. Lower lip ventricose; upper lip straight, its back not bearded; leaves linear; bracts with slender lobes, herbaceous and barely white-tipped; flowers pale; lower lip purple-spotted; spike virgate.—Common. Spring.

4. *Castilleia* Nutt. PAINTED-CUP.

Calyx spathaceous, cleft vertically. Corolla ringent, upper lip the larger. Stamens didynamous. Cells of anthers separated and of different shape and insertion. Capsule loculicidal.

1. *C. FOLIOLOSA* Hook. & Arn. Dorsal and ventral side of calyx equally cleft; plant tomentose; leaves linear; bracts crimson.—2f. Common. Summer.

2. *C. PARVIFLORA* Bong. Dorsal and ventral side of calyx equally cleft; plant villous, viscid; leaves laciniate, lobes linear; bracts crimson, yellow or white.—2f. San Mateo. Summer.

3. *C. LATIFOLIA* Hook. & Arn. Dorsal and ventral side of calyx equally cleft; plant villous, viscid; leaves oval, obtuse, usually entire; when lobed, lobe obtuse, oval; bracts crimson, yellow or white.—2f. Common. Summer.

4. *C. AFFINIS* Hook & Arn. Ventral side of calyx more deeply cleft than dorsal side. Calyx and upper part of the bracts petaloid, crimson, sometimes orange.—2f. Common. Summer.

Tribe II. VERONICEÆ.

Lower half of the corolla in æstivation covering the upper half. Corolla not tubular nor 2-labiate.

1. *Veronica* L. SPEEDWELL. BROOKLIME.

Calyx 4-5-cleft. Corolla rotate, 4-lobed, upper lobe the largest. Ovary few-seeded. Capsule compressed, emarginate.

1. *V. PEREGRINA* L. Upper leaves alternate, sessile; lowest leaves opposite, petiolate; flowers single, axillary bluish.—☉. Low grounds. Spring.

2. *V. AMERICANA* Schweinitz. All leaves opposite; flowers in axillary racemes; corolla bluish.—♂. Aquatic, Not common. Summer.

2. *Synthyris* Benth. (*Wulfenia* Jacq.)

Calyx 4-cleft. Corolla campanulate. Limb of corolla 4-lobed; posterior lobe emarginate, diandrous. Capsule 2-celled. Ovules ∞.

1. *S. ROTUNDIFOLIA* Gray. Flowers purplish.—♂. Lagunitas Creek. Spring.

3. *Limosella* L. MUDWORT.

Calyx 5-dentate. Didynamous. Limb of corolla 5-cleft. Cells of anthers confluent, transversely dehiscent. Capsule 1-celled, ∞-seeded.—☉.

1. *L. AQUATICA* L. Flowers pale.—Salt marshes. Summer.

Tribe III. GRATIOLEÆ.

Upper lip in æstivation covering the lower. Stigma flat, 2-lobed.

1. *Gratiola* L.

Calyx 5-parted, with two bractlets. Corolla 2-labiate. Filaments 4; two of them fertile, two sterile (sometimes 0). Ovary ∞ -ovulate. Capsule loculicidal, at length septifragal.

1. *G. EBRACTEATA* Benth. Corolla pale. \odot .—Marin County. Spring.

2. *Mimetanthe* Greene.

Calyx campanulate, not angulate. Capsule dehiscent only by the dorsal suture.

1. *M. PILOSA* Greene. Corolla yellow.— \odot . Dry river beds. Summer. The whole plant has a nauseous smell, resembling that of *Datura stramonium*.

3. *Mimulus* L. MONKEY-FLOWER.

Calyx angular; angles carinate. Placentæ of the capsule remain united and only separate at last near the apex. Valves of the capsule membranaceous.

1. *M. INCONSPICUUS* Gray. Glabrous; leaves ovate, entire, 3-5 nerved; all cauline leaves sessile; calyx ventricose, its teeth about equal and very short; corolla yellow.— \odot . Livermore. Summer.

2. *M. NASUTUS* Greene. Glabrous; stem 4-angular; leaves dentate, 3-5 nerved, mostly petioled; calyx ventricose, in fruit conspicuously pointed by the projecting upper tooth;

corolla yellow, commonly with a purple spot on the lower lip.—☉ Common in marshy places. Summer.

3. *M. LUTEUS* L. Glabrous; leaves variable in form, dentate; upper leaves sessile, lower leaves petiolate; calyx ventricose, inflated; teeth unequal, but in fruit without the projection of the upper largest tooth; corolla yellow, marked with purple.—☿. Common in marshy places. Summer.

4. *M. MOSCHATUS* Dougl. Villous and viscid; leaves pinnately veined; corolla yellow.—☿. Wet places in the mountains. Summer.

5. *M. CARDINALIS* Dougl. Villous and viscid; leaves ovate, the upper connate; limb of corolla very oblique; corolla scarlet.—☿. Water-courses. Contra Costa. Summer.

4. *Diplacus* Nutt. STICKY MONKEY-FLOWER.

Calyx angular; angles carinate. Placentæ meeting, but in dehiscence separate through their whole length. Valves of the capsule coriaceous. Tube of corolla funnel-shaped.

1. *D. GLUTINOSUS* Wendt. Flowers orange-color, but running sometimes into different shades of yellow and red.—♁. Common. Summer.

5. **Eunanus** Benth.

Calyx angular; angles carinate. Placentæ of the capsule separate in dehiscence for their whole length. Tube of corolla slender, filiform.

1. **E. DOUGLASHII** Benth. Corolla red, sometimes spotted.—☉. Marin County. Spring.

Tribe IV. DIGITALEÆ.

Upper lip of corolla in æstivation covering the lower lip. Stigma minute, not flat. Capsule septicidal.

1. **Pentstemon** L'Herit.

Didynamous, with a fifth sterile filament. Calyx 5-parted. Limb of corolla labiate.—♂.

1. **P. AZUREUS** Benth. Anthers sagittate; their cells confluent at the apex, dehiscent by a common split, that does not extend beyond the middle of the cell; corolla blue. Niles. Summer.

2. **P. CENTRANTHIFOLIUS** Benth. Cells of the anthers divaricate; dehiscent for their whole length; corolla scarlet. Niles. Summer.

Tribe V. ANTIRRHINEÆ.

Upper lip in æstivation covering the lower lip. Tube of corolla gibbous or calcarate. Capsule neither loculicidal nor septicidal.

1. *Collinsia* Nutt.

Calyx 5-parted. Corolla personate, upper lip erect, lower lip 3-lobed, middle lobe laterally compressed, hiding the didynamous stamens. Capsule few-seeded, septifragal.—☉.

1. *C. PARVIFLORA* Dougl. Flowers with long pedicels, mostly solitary in the axils of the leaves; throat of corolla longer than the limb; corolla small, blue; upper lip sometimes white. Saucelito. Spring.

2. *C. SPARSIFLORA* Fisch & Meyer. Flowers with long pedicels; mostly solitary in the axils of leaves; throat of corolla shorter than the limb; corolla pale-purple lower lip violet.—Marin County. Ocean Lake. Spring.

3. *C. BARTSLÆFOLIA* Benth. Flowers with short pedicels, crowded in the axils of leaves or bracts; throat of corolla longer than broad; corolla pale-purplish. — Mission Dolores. Spring.

4. *C. BICOLOR* Benth. Flowers with short pedicels, crowded in the axils of leaves or bracts; throat of corolla as broad as long, inflated and saccate; upper lip of corolla pale, lower lip violet.—San Rafael. Spring.

2. *Antirrhinum* L. SNAPDRAGON.

Didynamous. Calyx 5-cleft. Corolla saccate, 2-labiate, personate. Capsule dehiscent by pores at the apex.

1. *A. VAGANS* Gray. Corolla violet.—☉.
Contra Costa. San Mateo. Spring.

3. *Linaria* TOURN. TOAD-FLAX.

Didynamous. Calyx 5-cleft. Corolla with a spur, 2-labiate, personate. Capsule separating at the apex into two valves..

1. *L. CANADENSIS* Dumont. Corolla blue.—☉.
Common. Spring.

Tribe VI. VERBASCEÆ.

Upper lip in æstivation covering the lower lip. Corolla neither ringent nor personate. Capsule septicidal.

1. *Scrophularia* TOURN. FIGWORT.

Corolla short, globular. Limb narrow, 5-lobed. Middle lobe of lower lip reflexed. Didynamous (rudiment of a fifth stamen).

1. *S. CALIFORNICA* Cham. Corolla dark purple.—☿. Common. Summer.

Order 5. POLEMONIALES. Stamens as many as the lobes of the corolla, and more than the cells of the ovary. Stipules 0.

Family 1. SOLANACEÆ.

Corolla regular. Ovary 2-4 celled; ∞-ovulate. Fruit capsule or berry.

1. *Solanum* L. POTATO. NIGHTSHADE.

Calyx persistent in fruit. Corolla rotate; in æstivation valvate, induplicate. Anthers con-

nivent, opening by apical pores. Fruit an ∞ -seeded berry.

1. *S. NIGRUM* L. Corolla deeply 5-cleft; white or blue; berry black or orange.— \odot . Common throughout the year.

Very variable species. The berries of most, perhaps of all our Californian variations are not poisonous, and are frequently eaten by children.

2. *S. UMBELLIFERUM* Esch. Corolla merely 5-angled, rotate, blue; berries red.— $\frac{1}{2}$. $\frac{1}{2}$. Common. Flowering at all seasons. The berries of this species are at least suspicious.

2. *Datura* L. THORN-APPLE.

Calyx tubular, circumscissile. Corolla funnel-shaped, plicate and convolute in æstivation. Anthers dehiscent longitudinally. Ovary spuriously 4-celled. Capsule ovate, septifragal. Seeds reniform.

1. *D. STRAMONIUM* L. Corolla white, tinged with pale blue.— \odot . Sunnyside. Summer.

Poisonous ballast-weed; differs from the European and Asiatic type by the bluish tinge of its corolla.

3. *Nicotiana* L. TOBACCO.

Calyx tubular, persistent. Corolla funnel or salver-shaped. Anthers longitudinally dehiscent. Ovary 2-celled, ∞ -ovulate. Capsule septicidal. Seeds ∞ , minute.

1. *N. BIGELOVII* Watson. Flowers white.
⊙. Buena Vista, Sonoma. Summer.

Family 2. **BORRAGINEÆ.**

Ovary 4-celled, 4-lobed. Cells 1-seeded.
Fruit 4 nutlets. Style 1.

1. *Amsinckia* Lehm.

Calyx 5-parted. Corolla salver- or funnel-shaped. Limb 5-lobed. Ovary 4-lobed. Style central. Fruit 4 nutlets, attached to a conical disc above its base.—⊙. Inflorescence circinate. Flowers yellow.

1. *A. VERNICOSA* Hook & Arn. Nutlets triquetrous, straight, smooth, shining; attached at the lower part of the inner angle.—Contra Costa. Spring.

2. *A. TESSELATA* Gray. Nutlets testaceous, tessellate, rugose; calyx lobes obtuse; corolla orange-color.—Contra Costa. Spring.

3. *A. INTERMEDIA* Fisch. & Mey. Nutlets scabrous, convex and carinate on the back; calyx-lobes narrow, acute; corolla tube scarcely exceeding the calyx.—Paper Mill Creek. Spring.

4. *A. SPECTABILIS* Fisch. & Mey. Nutlets granulate, convex and carinate on the back; calyx lobes linear; corolla orange, tube at least twice the length of the calyx lobes.—Contra Costa. Spring.

6. *A. LYCOPSOIDES* Lehm. Nutlets reticulate, rugose; calyx lobes lanceolate.—Common. Spring.

2. *Krynitzkia* Fisch. & Mey.
(*Eritrichium* Schrad).

Nutlets ventrally attached to the conical or columnar disc from near their base.

1. *K. MURICULATA* Gray. Nutlets attached to the disc from the base nearly up to the apex; calyx in fruit somewhat longer than the nutlets, its sepals lanceolate, armed with pungent bristles; mid-rib of sepals of usual shape; nutlets muricate; corolla white.—☉. Common. Spring.

2. *K. AMBIGUA* Gray. Nutlets attached to the disc from the base up to nearly the apex; calyx in fruit considerably longer than the nutlets, its sepals narrow, armed with pungent bristles; mid-rib of sepals prominent; nutlets scabrous; corolla white.—☉. Common. Spring.

3. *K. TORREYANA* Gray. Nutlets attached to the pyramidal disc only up to their middle; nutlets smooth, all four maturing; corolla white.—☉. Saucelito. Spring.

4. *K. LEIOCARPA* Fisch. & Mey. Nutlets attached to the subulate disc nearly their whole length, smooth, all four maturing; corolla white.—☉. Saucelito. Spring.

5. *K. OXYCARYA* Gray. Nutlets attached only by their lower third to the conical gyno base; only one of the four maturing; corolla white.—⊙. Marin County. Spring.

6. *K. CHORISIANA* DC. Nutlets attached to the disc only by their base; corolla nearly rotate; scales near the throat conspicuous, puberulent; corolla white.—⊙. Wet ground, San Francisco. Spring.

7. *K. CALIFORNICA* DC. Nutlets attached to the disc only by their base; corolla inconspicuous; scales near the throat not puberulent; corolla white.—⊙. Wet grounds. Common. Spring.

3. *Plagiobothrys* Fisch. & Mey.

(*Eritrichium* Schrad).

Nutlets attached near the middle of their ventral face to the conical disc.

1. *P. NOTHOFULVUS* Gray. Calyx cleft only to its middle; corolla white.—⊙. Common. Spring.

2. *P. CANESCENS* Benth. Calyx cleft nearly to the base; corolla white.—⊙. Livermore. Spring.

4. *Pectocarya* DC.

Corolla funnel-shaped; throat closed. Style central, very short. Fruit four nutlets, marginate and in pairs,—⊙. Flowers axillary, sessile, white.

1. *P. PENICILLATA* A. DC.—Marin County. Spring.

5. *Cynoglossum* L. HOUND'S-TONGUE.

Corolla funnel-shaped. Throat closed by five scales. Ovary 4-lobed. Style central. Fruit 4, depressed, echinate nutlets.— 2ζ . Inflorescence paniculate, ebracteate, circinate.

1. *C. GRANDE* Dougl. Corolla blue.—Common in woods. Spring.

6. *Heliotropium* L. HELIOTROPE.

Corolla salver-shaped. Limb 5-lobed; sinuses plicate. Ovary 4-celled. Cells 1-seeded. Style terminal, short. Stigma peltate. Fruit a 4-pyrenous drupe. Inflorescence circinate.

1. *H. CURASSAVICUM* L. Corolla pale.— 2ζ . Seashore. Summer.

Family 3. **HYDROPHYLLÆ.**

Ovary incompletely 2-celled. Style 2-cleft or 2 styles. Fruit a capsule.

1. *Eriodictyon* Benth. YERBA SANTA.

Corolla campanulate. Ovary 2-celled; cells ∞ -ovulate: styles 2; stigmas clavate. Fruit a capsule, 2-celled, few-seeded, at first loculicidal, at last septical.— $\frac{1}{2}$. Inflorescence circinate, arranged in a panicle. Flowers blue or white.

1. *E. GLUTINOSUM* Benth. Corolla pale or bluish.—Chaparral and chemisal. Summer.

An exceedingly variable species. The resinous exudation of the young branches and leaves has a bitter, somewhat aromatic taste. The plant is called "Yerba Santa." It acts as a mild diuretic, and was used as such by the old inhabitants of California. At present considerable quantities are exported, partly for medicinal purposes, partly as a harmless and agreeable substitute for hops in brewing certain varieties of beer, especially porter.

2. *Romanzoffia* Cham.

Corolla salver-shaped. Style entire, Stigma capitate. Ovary incompletely 2-celled, ∞ -ovulate. Capsule ovate, loculicidal. Seeds ∞ , very small.— σ . Inflorescence loosely circinate.

1. *R. SITCHENSIS* Bongard. Corolla pale.—Sunnyside. Paper Mill Creek. Spring.

3. *Emmenanthe* Benth.

Calyx 5-parted. Sinuses naked; lobes equal. Corolla campanulate, persistent. Ovary incompletely 2-celled, ∞ -ovulate. Stigma 2-cleft.— \odot . Inflorescence circinate. Flowers yellow.

1. *E. PENDULIFLORA* Benth. Corolla cream-colored.—Livermore. Summer.

4. *Phacelia* Juss.

Calyx 5-parted. Sinuses naked. Lobes equal. Corolla campanulate, deciduous. Ovary incompletely 2-celled or 1-celled, with two parietal placentæ. Number of ovules variable. Stigma 2-cleft. Capsule incompletely 2-celled or 1-celled, loculicidal. Inflorescence circinate, cymose.

1. *P. CIRCINATA* Jacq. Ovary contains four ovules, inflorescence thyrsoïd, circinate, with very short pedicels; leaves entire, the lowermost tapering into a petiole with one or two pairs of leaflets; corolla blue or white.—♀. Common. Summer.

2. *P. MALVÆFOLIA* Cham. Ovary contains four ovules; circinate spikes solitary or geminate, not collected into a thyrsoïd inflorescence; all leaves petiolate, cordate, lobed, dentate; corolla pale.—⊙. San Francisco. Summer. Leaves sting somewhat like nettles.

3. *P. HISPIDA* Gray. Ovary contains four ovules; inflorescence thyrsoïd, circinate; flowers with short pedicels; sepals very narrow, much longer than the capsule; leaves about 5-parted; corolla red.—⊙. Livermore. Spring.

4. *P. DISTANS* Benth. Ovary contains four ovules; inflorescence thyrsoïd, circinate; flowers sessile; sepals unequal and much longer than the capsule; leaves finely and decom-

poundly dissected; stamens scarcely exerted; corolla whitish, ochroleucous or violet.—⊙. Alameda. Marin County. Summer.

5. *P. TANACETIFOLIA* Benth. Ovary containing four ovules; inflorescence thyrsoïd, circinate; flowers with very short pedicels; sepals but little longer than the capsule; leaves finely and decom poundly dissected; stamens much exerted; corolla bluish.—⊙. Strawberry Valley. Contra Costa. Summer.

6. *P. CILIATA* Benth. Ovary containing four ovules; inflorescence cymose, circinate; the circinna rather short; sepals in fruit accrescent and transversely veined; leaves pinnately parted, the partitions pinnatifid; stamens somewhat shorter than the corolla; white or blue. ⊙.—Belmont. Spring.

7. *P. DIVARICATA* Gray. Ovary containing more than four ovules; inflorescence in racemes; pedicels short; leaves ovate, entire; corolla blue.—⊙. Tiburon. Spring;

5. *Ellisia* L.

Calyx 5-parted, sinuses naked. Corolla campanulate, stamens included. Ovary 1-celled with two parietal placentaë. Stigma 2-cleft. Capsule membranaceous, loculicidal, the placentaë separating from the capsular valves, simulating a second internal capsule. ⊙.—Flowers pale.

1. *E. CHRYSANTHEMIFOLIA* Benth. Corolla whitish.—Oakland hills. Spring.

6. *Nemophila* Nutt.

Calyx with reflexed, appendiculate sinuses. Corolla campanulate. Ovary 1-celled, with two parietal placentæ. Stigma 2-cleft. Capsule membranaceous, loculicidal, the placentæ separating from the capsular valves, simulating a second internal capsule.—○.

1. *N. INSIGNIS* Dougl. Ovary containing more than four ovules; scales on the base of corolla rounded and partly free; corolla blue. Common. Spring.

2. *N. MENZIESII* Hook. & Arn. Ovary containing more than four ovules; scales at the base of corolla narrow and wholly adnate; corolla blue or white, more or less marked by dots.—Common. Spring.

3. *N. AURITA* Lindl. Ovary containing but four ovules; scales on the base of corolla in pairs, broad and partly free; corolla violet. San Mateo. Spring,

4. *N. PARVIFLORA* Dougl. Ovary contains but 4 ovules; scales on the base of corolla oblong, wholly adnate by one edge; corolla light blue or white.—Common in shady places. Spring.

Family 4. **POLEMONIACEÆ.**

Æstivation imbricate. Ovary 3-celled. Style 3-cleft. Embryo straight. Cotyledons foliaceous.

1. *Polemonium* Tourn. JACOB'S-LADDER.

Corolla rotate. Stamens ascending. Filaments dilated at their base. Seeds ∞ .

1. *P. CARNEUM* Gray. Corolla blue.— γ . Point Bonita, near the lighthouse. Summer.

2. *Gillia* Ruiz & Pavon.

Corolla funnel or salver-shaped. Stamens straight, equally inserted.

1. *G. PUSILLA* Benth. Leaves opposite, uppermost sometimes alternate, palmately divided; divisions filiform; corolla short, funnel-shaped; pedicels capillary; corolla purplish, with yellowish throat, sometimes pale.— \odot . Marin County. Spring.

2. *G. DICHOTOMA* Benth. Leaves all opposite; corolla salver-shaped, its tube shorter than the calyx; flowers almost sessile, but not capitate; corolla white.— \odot . Livermore. Spring.

4. *G. DENSIFLORA* Benth. Leaves opposite, palmately parted, fascicled in the axils; corolla salver-shaped, its tube about the length of the obovate lobes; flowers in glomerules; bracts herbaceous; corolla red or white.— \odot , Alameda. San Rafael. Spring.

4. *G. ANDROSCAEA* Steud. Leaves opposite, palmately parted; corolla salver-shaped, its tube more than twice the length of the lobes;

flowers in glomerules; bracts herbaceous, hirsute, much shorter than the corolla tube; corolla red or white, with yellowish throat.—⊙. Common. Spring.

5. *G. MICRANTHA* Steud. Leaves opposite, palmately parted; corolla salver-shaped, its tube more than four times the length of the lobes; flowers in glomerules; bracts herbaceous, pubescent, shorter than the flower-tube; corolla red, white, lilac or yellow.—⊙.

6. *G. TENELLA* Benth. Leaves opposite, palmately parted; corolla salver-shaped, its tube more than four times the length of the corolla lobes; flowers in glomerules; bracts herbaceous, hispidulous, shorter than the corolla tube; corolla purple or pink, with yellow throat.—⊙. Common. Spring.

7. *G. CILIATA* Benth. Leaves opposite, palmately parted; corolla salver-shaped, its tube more than four times the length of the lobes; flowers in glomerules; bracts very hirsute, about the length of the corolla tube; calyx lobes acerose; corolla rose, violet or pale.—⊙. Marin County. Spring.

8. *G. SQUARROSA* Hook. & Arn. Leaves alternate, pinnately parted, partitions parted or incised; corolla funnel-shaped; flowers in glomerules, densely bracteate; bracts rigid, pungent; stamens included in the tube of the corolla; corolla blue.—⊙. Common. Summer.

The disagreeable smell of this plant is the cause of its California name, "Skunkweed."

9. *G. COTULÆFOLIA* Steud. Leaves alternate, 2-pinnately parted; corolla funnel-shaped; flowers in glomerules, densely bracteate; bracts spinescent; stamens exserted; ovules no more than two in each cell; corolla bluish or pale.—☉. Common. Summer.

Smell somewhat like Chamomile.

10. *G. INTERTEXTA* Steud. Leaves alternate, pinnately parted, partitions but little, if at all, divided; flowers in glomerules, densely bracteate; bracts with very villous base; corolla funnel-shaped, not exceeding the calyx lobes; stamens exserted; ovules more than two in each cell; corolla small and white.—☉. Common. Summer.

11. *G. LEUCOCEPHALA* Gray. Leaves alternate, pinnately parted, soft; partitions slender and frequently entire; flowers in glomerules; bracts barely pungent; corolla funnel-shaped, exceeding the calyx-lobes; stamens considerably exserted: ovules two in each cell; corolla white.—☉. Common. Summer.

12. *G. VISCIDULA* Gray. Leaves alternate, pinnately parted; partitions entire, subulate: flowers in glomerules; bracts dilated at their base; corolla funnel-shaped, about twice the length of the spinescent calyx-lobes; plant

viscid, pubescent; corolla violet.—⊙. Tamalpais. Summer.

13. *G. DENSIFOLIA* Benth. Leaves alternate, pinnately laciniate, lobes spinulose; corolla salver-shaped; stamens exserted; flowers in glomerules; bracts foliaceous; stems leafy to their tops; corolla blue.—24. Livermore. Summer.

14. *G. VIRGATA* Steud. Leaves alternate, filiform, simple or 3-parted, the partitions filiform; corolla salver-shaped; stamens exserted; flowers in glomerules; bracts foliaceous; stems with but few leaves; virgate; corolla blue.—⊙. Livermore. Summer.

15. *G. CAPITATA* Dougl. Leaves alternate, 2-3-pinnately divided into slender lobes; corolla funnel-shaped, its throat being very slightly dilated; flowers crowded but not sessile, the cluster itself on a long stalk; bracts inconspicuous; calyx glabrous; stamens inserted into the sinuses of the light blue corolla. ⊙.—Marin County. Summer.

16. *G. ACHILLEÆFOLIA* Benth. Leaves alternate, 2-3-pinnately divided into slender lobes; corolla somewhat funnel-shaped, but its throat abruptly and amply dilated; flowers in clusters but not sessile; bracts inconspicuous; calyx woolly; corolla blue.—⊙. Sand hills. San Francisco. Summer.

17. *G. MULTICAULIS* Benth. Leaves alternate, 2-pinnately parted into linear lobes; corolla funnel-shaped, its tube shorter than the calyx and about the length of the ovate corolla lobes; flowers in clusters, pedicels in fruit about equaling the calyx; bracts inconspicuous; corolla violet.—☉. Oakland hills. Summer.

18. *G. TRICOLOR* Benth. Leaves alternate, 2-pinnately parted into linear lobes; corolla much longer than the calyx with very short tube, but ample, funnel-shaped throat; lobes of the corolla longer than the stamens; flowers in short peduncled clusters; pedicels very short; bracts inconspicuous; tube of the corolla yellowish, throat marked with purple; lobes violet.—☉. Livermore. Summer.

19. *G. INCONSPICUA* Dougl. Leaves alternate, pinnatifid; lobes short, mucronate; corolla narrowly funnel-shaped, about the length of the calyx; bracts inconspicuous; corolla violet.—☉. Common. Summer.

3. *Collomia* Nutt.

Corolla salver-shaped, with long tube. Stamens unequally inserted.

1. *C. GRACILIS* Dougl. All leaves sessile, entire; corolla violet; the tube yellowish.—☉. Tamalpais, Spring.

The seeds when moistened develop a mucilage, which in Mexico is called "Chia" and used in the preparation of a cooling drink.

2. *C. GILIOIDES* Benth. Lower leaves petiolate and most of them pinnately incised; calyx rounded at base; corolla pink.—☉. Tamalpais. Spring.

3. *C. HETEROPHYLLA* Hook. Lower leaves petiolate, 2-pinnatifid; calyx acute at its base; corolla pink.—☉. Lagunitas Creek. Spring.

Family 5. CONVULVACEÆ.

Æstivation contorted. Ovary 2-3 or 4-celled. Embryo curved. Cotyledons foliaceous, conduplicate, corrugate.

1. *Dichondra* Forst.

Ovary divided into two carpidia, each 2-ovulate. Creeping herb. Flowers inconspicuous.

1. *D. REPENS* Forst. Flowers pale.—♀. Telegraph hill. Spring.

2. *Convolvulus* L.

BIND-WEED. MORNING-GLORY.

Calyx 5-cleft, persistent. Corolla funnel-form, 5-angulate, 5-plicate. Cells of the capsule 2-ovulate.—♀.

1. *C. SOLDANELLA* L. Margin of corolla entire; stigmas ovate, thickish; corolla pink. Seashore. Summer.

2. *C. CALIFORNICUS* Choisy. Margin of corolla entire; stigmas linear-oblong, flat; bracts at the base of the calyx; corolla white or reddish.—Berkeley. Summer.

3. *C. LUTEOLUS* Gray. Margin of corolla entire; stigmas linear, flat; bracts distant from the calyx; corolla pale-yellow, sometimes reddish.—Common. Summer.

4. *C. ARVENSIS* L. Margin of corolla entire; stigmas filiform; bracts at the base of the pedicel and minute; corolla white, sometimes tinged with rose.—Common. Summer.

5. *C. PENTAPETALOIDES* L. Corolla deeply 5-cleft, purplish.—☉. Not common. Contra Costa. Summer.

3. *Cressa* L.

Corolla funnel-shaped, 5-parted, not plicate. Styles 2. Cells of the ovary 2-ovulate. Fruit a 1-seeded capsule.—2f.

1. *C. CRETICA* L. Corolla white, silky outside.—Seashore. Summer.

4. *Cuscuta* Tourn. DODDER.

Aphyllous. Destitute of chlorophyll. Corolla urceolate. Limb 4-5-lobed. Ovary 2-celled; cells 2-ovulate. Capsule indehiscent, sometimes circumscissile. Embryo spiral. Cotyledons 0.

1. *C. SALINA* Engelm. Calyx lobes as long as the tube of the white corolla.—Common. Summer. Parasite of *Salicornia* and other salt-marsh herbs.

2. *C. SUBINCLUSA* Durand & Hilgard. Calyx lobes much shorter than the tube of corolla; corolla white, calyx usually reddish.—Parasites on *Ceanothus* and other shrubs.—Not common. Summer.

Section II. ISOCARPICÆ (Number of carpodia corresponding to number of floral parts).

Order 1. PRIMULALES.

Stamens opposite to the lobes of corolla. Ovary 1-celled. Placenta central.

Family 1. LENTIBULARIÆ.

Diandrous. Corolla irregular, spurred.

1. *Utricularia* L. BLADDERWORT.

1. *U. VULGARIS* L. Corolla yellow. Aquatic.—Olema. Summer.

Family 2. PLUMBAGINEÆ.

All parts in fives except the ovule, which is single.

1. *Armeria* Willd. THRIFT.

Flowers in an involucrate head. Corolla 5-parted or five distinct petals. Styles 5.—2.

1. *A. VULGARIS* Willd. Corolla rose-color. Common. Spring.

2. *Statice* L. SEA-LAVENDER.

Flowers bracteate in one-sided spikes.—2f.

1. *S. LIMONIUM* L. Corolla violet.—Salt marshes. Summer.

Family 3. PRIMULACEÆ.

Pentandrous. Placenta central, ∞ -ovulate.

1. *Dodecatheon* L. SHOOTING STAR.

Flowers umbellate. Filaments shorter than the anthers, connivent in a cone. Acaulescent. Flowers purple, pink or white.—2f.

2. *D. MEADIA* L.—Common. Spring.

2. *Androsace* Tourn.

Flowers umbellate. Filaments shorter than the anthers, included in the salver-shaped corolla.

1. *A. OCCIDENTALIS* Pursh. Flowers white.—
⊙. Contra Costa Range. Santa Cruz mountains. Spring.

3. *Glaux* L. MILKWORT.

Calyx campanulate, 5-cleft, colored. Corolla 0. Stamens 5, alternate with the lobes of the calyx. Capsule 5-valved, few-seeded.—2f. Leaves decusate; flowers white.

1. *G. MARITIMA* L.—Salt marshes. Summer.

4. *Trientalis* L. STAR-FLOWER.

Flowers several, terminal. Floral parts 5 to 7. Capsule few-seeded, longitudinally dehiscent.—2f.

1. *LATIFOLIA* Torrey. Corolla pink or pale.—
Shady woods. Spring.

5. *Anagallis* Tourn. PIMPERNEL.

Flowers axillary. Lobes of the corolla broad.
Capsule transversely dehiscent. Leaves oppo-
site.

1. *ARVENSIS* L. Corolla red, sometimes pale,
purple or blue.—☉. Common all the year
round. An infusion of this herb was used by
the old Californians for headache.

6. *Samolus* Tourn. BROOKWEED.

Calyx half-superior. Corolla campanulate.
Stamens 10; the five fertile opposite to the lobes
of the corolla; the five sterile, alternate. Cap-
sule 5-valved, ∞ -seeded. Leaves alternate;
flowers white.— \mathcal{L} .

1. *L. VALERANDI* L. Rare. Formerly in a
marsh near San Francisco; at present found
occasionally in Marin County.

ORDER 2. *ERICALES*. Stamens alternate
with the lobes of corolla or twice as many.
Placentæ axillary.

Family 1. *PYROLACEÆ*.

Corolla split into petals. Hypogynous disc
0. Seeds minute.

1. *Pyrola* Tourn.

Calyx 5-parted. Corolla 5-petalous. Stamens 10. Style filiform.— $\frac{1}{2}$. In forest shades.

1. *P. APHYLLA* Smith. Parasitic without chlorophyll; petals white.—Lagunitas Creek. Summer.

Family 2. RHODORACEÆ.

Ovary free. Fruit a capsule, septicidal.

1. *Rhododendron* L.

Corolla funnel-shaped, 5-lobed. Stamens ascending. Cells of anthers dehiscent by an apical pore. Seeds ∞ , minute.— $\frac{1}{2}$.

1. *R. OCCIDENTALE* Gray. Corolla white, upper lobes yellow.—Point Bonita. Taylorville. Sonoma. Summer. (Azalea.) The root of this shrub contains a powerful narcotic.

Family 3. ERICACEÆ.

Gamopetalous. Fruit either baccate or a loculicidal capsule.

1. *Gaultheria* L. WINTERGREEN.

Calyx 5-cleft, petaloid. Corolla urceolate. Stamens 10. Cells of anthers dehiscent by an apical pore. Fruit a spurious berry, i. e., a capsule, 5-celled, ∞ -seeded, enclosed by the enlarged and fleshy calyx.—Leaves evergreen. Flowers white, rosy.— $\frac{1}{2}$.

1. *G. SHALLON* Pursh. Berry red, at last black. Edible. Saucelito. Tamalpais.

2. *Arctostaphylos* Adans. MANZANITA.

Corolla urceolate. Stamens 10. Disc hypogynous. Cells of ovary 1-ovulate. Fruit a berry. Cells 1-seeded; seeds sometimes coherent by their covering. Leaves evergreen. Flowers white, rosy.— $\frac{1}{2}$. Fruit edible.

1. *A. PUNGENS* HBK. Almost glabrous; leaves entire, coriaceous, mucronate; petioles slender; pedicels glabrous; drupe glabrous; corolla white.— $\frac{1}{2}$. Common. Spring.

2. *A. TOMENTOSA* Dougl. Young branchlets tomentose, old ones bristly; leaves almost entire, coriaceous, petioled; ovary hirsute; drupe puberulent, but becoming glabrous at last; corolla white.—Hillsides. Spring.

3. *A. ANDERSONII* Gray. Branchlets puberulent and bristly; leaves generally conspicuously serrulate, thin, almost sessile; drupes depressed, densely covered with viscous bristles. Livermore. Spring.

3. *Arbutus* TOURN. MADROÑA.

Corolla urceolate. Stamens 10. Disc hypogynous. Cells of ovary ∞ -ovulate. Fruit a berry. Cells several-seeded. Leaves evergreen. Flowers white.— $\frac{1}{2}$.

1. *A. MENZIESII* Pursh.—Common. Spring.
Fruit edible.

Family 4. VACCINIÆ.

Ovary inferior.

1. *Vaccinium* L. HUCKLEBERRY.

Anther cells separate, elongated at the apex into a tube. Fruit a berry, ∞ -seeded, crowned by the persistent calyx.

1. *V. OVATUM* Pursh. Corolla pink.— $\frac{1}{2}$.
Redwoods. Spring.

Sub-Series 2. POLYPETALÆ.

Section 1. **DISCOPHORÆ.** Ovary inferior and crowned by a well developed disc.

ORDER 1. UMBELLALES. Cells of ovary 1-ovulate. Stamens alternate.

Family 1. UMBELLIFERÆ.

Stamens 5. Ovary 2-celled. Styles 2.
Fruit a schizocarp. (Cremocarp.)

1. *Caucalis* L.

Umbels regularly compound. Secondary ribs of the cremocarp more prominent than the primary. Margins of the endosperm inflexed \odot .—Flowers white.

1. *C. NODOSA* Hudson. Decumbent; umbels naked.—Marin County. Livermore. Spring.

2. *C. MICROCARPA* Hook. & Arn. Erect; umbels involucrate.—Common. Spring.

The chewing of this herb, called by the Spanish “Yerba de vibora,” or an infusion of it, is recommended by them as an antidote against the bite of the rattlesnake.

2. *Daucus* Tourn. CARROT.

Umbels regularly compound. Secondary ribs of the cremocarp more prominent than the primary. Intervals 1-vittate. Face of the endosperm flat.

1. *D. PUSILLUS* Michx. Flowers white.—☉. Common. Spring.

3. *Ferula* Tourn.

Umbels regularly compound. Cremocarp dorsally compressed. Marginal ribs winged. Wings coherent. Dorsal ribs filiform. Vittæ ∞. Flowers yellow.—2♂.

1. *F. CALIFORNICA* Gray.—Sunnyside. Summer.

4. *Heracleum* L. COW-PARSNIP.

Umbels regularly compound. Cremocarp dorsally compressed. Marginal ribs winged. Wings coherent. Vittæ shorter than the mericarp.—2♂. Flowers white.

1. *H. LANATUM* Michx.—Common. Summer. Herb of suspicious qualities.

5. *Peucedanum* L.

Umbels regularly compound. Cremocarp dorsally compressed. Marginal ribs winged. Wings coherent. Vittæ as long as the mericarp.—2f.

1. *P. DASYCARPUM* Torr. & Gray. Leaves much dissected; cremocarp tomentose; flowers white.—Common. Summer.

Smell of entire plant like celery.

2. *P. UTRICULATUM* Nutt. Leaves much dissected; cremocarp glabrous, distinctly ribbed; flowers yellow.—Common. Summer.

3. *P. CARUIFOLIUM* Torr & Gray. Leaves much dissected; cremocarp glabrous; dorsal ribs obsolete, only the marginal ones developed; flowers yellow.—Common. Summer.

4. *P. TRITERNATUM* Nutt. Leaves 2-ternate to 3-quinate; umbellules involuclate; flowers yellow.—Contra Costa. Summer.

5. *P. LEIOCARPUM* Nutt. Leaves 2-ternate to 3-quinate; umbellules naked; flowers yellow.—Livermore. Summer.

The roots of several species of *Peucedanum* have been used as food, but as those used in this way are not sufficiently identified it is safer to abstain from using them till the different species have been subjected to an examination, not merely in regard to their botanical

characters, but also in regard to their effects on the human organism.

6. Angelica L.

Umbels regularly compound. Cremocarp dorsally compressed. Marginal ribs winged. Wings distinct; intervals 1-vittate. Flowers white.—2f.

1. A. TOMENTOSA Watson.—Common. Summer. Properties like those of the officinal *Angelica Archangelica*.

7. Selinum L.

Umbels regularly compound. Cremocarp dorsally compressed, 4-winged by very distinct marginal ribs, and only coherent by a carinate commissure.—2f. Flowers white.

1. S. PACIFICUM Watson.—Saucelito. Summer.

8. Ligusticum L. LOVAGE.

Umbels regularly compound. Cremocarp dorsally compressed, 4-winged by the very distinct lateral ribs. Mericarps coherent by a convex commissure.—2f.

1. L. APIIFOLIUM Benth. & Hook. Flowers white.—Tamalpais. Summer.

9. Œnanthe L. FOOL'S PARSLEY.

Umbels regularly compound. Cremocarp terete, ovate. Ribs obtuse. Intervals 1-vittate. Aquatic. Flowers white.—2f.

1. *Æ. CALIFORNICA* Watson.—Common. Aquatic. Summer.

The European congeners of this plant, *Æ. fistulosa* and *Æ. crocata*, are undoubtedly poisonous. Our species is eagerly eaten by cattle, a fact that becomes very evident in a swamp near the Presidio, where it grows in company with *Cicuta Californica*, the latter not being touched by the animals when *Ænanthe* is eaten up to the roots. It would be desirable to investigate the properties of the species.

10. *Osmorrhiza* Raf. SWEET CICELY.

Umbels regularly compound. Cremocarp terete, elongate, angulate, sulcate, hispid. Vittæ 0. Flowers white.— \mathcal{L} .

1. *O. BRACHYPODA* Torr. Involucral bracts linear and acuminate, equaling the flowers. Saucelito. Summer.

2. *O. NUDA* Torr. Involucre and involucels small and caducous.—Common. Summer.

11. *Sium* L. WATER-PARSNIP.

Umbels regularly compound. Cremocarp laterally compressed, oblong. Mericarp dorsally convex, plain on the face, 5-ribbed. Ribs filiform. Intervals more than 1-vittate. Flowers white.— \mathcal{L} .

1. *S. CICUTÆFOLIUM* Gmelin.—Aquatic. Baden Station. Summer. Probably poisonous.

12. *Cicuta* L. WATER-HEMLOCK.

Umbels regularly compound. Cremocarp laterally contracted. Ribs flattened, intervals 1-vittate. Mericarp terete. Aquatic. Flowers white.— \mathcal{L} .

1. *C. CALIFORNICA* Gray. Leaves pinnate, the lower sometimes 2-pinnate at base; leaflets serrate, the veinlets running into the teeth. Presidio. Summer.

2. *C. BOLANDERI* Watson. All leaves 2-pinnate; leaflets serrate, the veinlets running into the sinuses; involucre bracts linear.—Suisun. Alvarado marshes. Summer.

3. *C. MACULATA* L. Lower leaves 2-pinnate; leaflets serrate, the veinlets running into the sinuses; involucre obsolete.—Tamalpais. Summer. Poisonous aquatics.

13. *Berula* Koch.

Umbels regularly compound. Cremocarp laterally contracted, ovate. Mericarp terete, 5-ribbed, lateral ribs not contiguous. Epicarp thick, corky. Ribs filiform. Intervals ∞ -vittate.

1. *B. ANGUSTIFOLIA* Koch.— \mathcal{L} . Flowers white. Baden Station. Summer.

14. *Pimpinella* L.

Umbels regularly compound. Cremocarp laterally contracted, ovate. Mericarp convex,

dorsally, plain on the ventral side, 5-ribbed. Ribs filiform. Lateral ribs contiguous. Intervals ∞ -vittate.—24.

1. *P. APIODORA* Gray. Flowers pale.—Contra Costa. Belmont. San Mateo. Summer.

Odor pleasant, like celery. An infusion of the root is used in cases of chronic catarrh, like the tincture of its European congener. *P. Saxifraga*.

15. *Carum* L. CARAWAY.

Umbels regularly compound. Cremocarp laterally compressed, oblong. Mericarp equally 5-ribbed. Ribs filiform. Lateral ribs contiguous. Commissure plain. Intervals 1-vittate. Flowers white.—⊙ ⊙.

1. *C. KELLOGGII* Gray.—Presidio. Alameda. Sunnyside. Summer. Seeds very fragrant.

16. *Apiastrum* Nutt.

Umbels regularly compound. Cremocarp very much contracted at the commissure, cordate. Mericarp incurved at base and apex, 5-ribbed; ribs little elevated. Intervals 1-vittate. Flowers white.—⊙.

1. *A. ANGUSTIFOLIUM* Nutt.—Niles. Spring.

17. *Conium* L. HEMLOCK.

Umbels regularly compound. Cremocarp laterally compressed. Endosperm deeply sul-

iate on the ventral side. Mericarp 5-ribbed; ribs undulate, crenate. Vittæ 0. Flowers white.—⊙⊙.

1. *C. MACULATUM* L.—San Mateo. Sunny-side. Summer.

A well known medicinal and poisonous herb.

18. *Deweya* Torr. & Gray.

Umbels regularly compound. Cremocarp laterally compressed. Mericarp reniform in transverse section, 5-ribbed. Intervals more than 1-vittate. Flowers yellow.—2f.

1. *D. HARTWEGI* Gray. Ribs of mericarp prominent.—Oakland hills. Wild-Cat Creek. Strawberry Valley. Spring.

2. *D. KELLOGGII* Gray. Ribs of mericarp filiform.—Tamalpais. Spring.

19. *Sanicula* Tourn. SANICLE.

Umbels not regularly compound. Lobes of calyx limb foliaceous. Flowers polygamous. Cremocarp subglobose, aculeate. Ribs 0. Vittæ ∞.—2f.

1. *S. TUBEROSA* Torr. Leaves 2-3-pinnate; flowers yellow.—Tiburon. Spring.

2. *S. BIPINNATIFIDA* Dougl. Ripe cremocarp sessile; leaves long petiolate, triangular in outline, pinnately 3-5-lobed; segments distant, incisely lobed and decurrent on their rachis; flowers purple, in some localities yellow. Common. Spring.

3. *S. LACINIATA* Hook. & Arn. Ripe cremocarp sessile; leaves triangular in outline, 3-parted, the partitions pinnatifid to 2-pinnatifid. flowers yellow.—Tamalpais. Spring,

4. *S. MENZIESII* Hook. & Arn. Ripe cremocarp pedicillate; leaves deeply 3-5-lobed; the lobes broad, dentate; flowers yellow.—Common. Spring.

5. *S. ARCTOPOIDES* Hook. & Arn. Ripe cremocarp pedicillate; leaves deeply 3-lobed, the lobes laciniately cleft; flowers yellow.—Common. Spring.

6. *S. MARITIMA* Kellogg. Lower leaves entire, or slightly 3-lobed; upper leaves palmately parted, partitions cuneate and somewhat lobed; flowers greenish.—Alameda marshes. Summer.

22. *Eryngium* Tourn. ERYNGO.

Umbels not regularly compound. Lobes of the calyx-limb rigid. Cremocarp tuberculate. Ribs 0. Vittæ 0.—Spinose plants.

1. *E. PETIOLATUM* Hook. Flowers bluish. ☉. Marshes. Summer.

21. *Bowlesia* Ruiz & Pavon.

Umbels simple. Cremocarp ovate, much contracted on the commissure, plane on the dorsal sides of the mericarps. Ribs 0. vittæ 0. Leaves opposite.

1. *B. LOBATA* Ruiz & Pavon. Flowers white.—⊙. Golden Gate Park. San Mateo. Spring.

22. *Hydrocotyle* TOURN. WATER PENNYWORT.

Umbels simple. Cremocarp laterally compressed, c a r i n a t e. Ribs filiform. Vittæ 0. Aquatic.—24.

1. *H. PROLIFERA* Kellogg. Leaves peltate, emarginate at base, crenate. Flowers greenish.—San Francisco. Summer.

2. *H. RANUNCULOIDES* L. Leaves orbicular, not peltate, lobate; lobes crenate; flowers greenish.—San Francisco. Summer.

Family 2. *ARALIACEÆ*.

Fruit a berry or drupe. Leaves alternate.

1. *Panax* L. (*Fatsia* Benth. & Hook).

Polygamous. Petals 5. Stamens 5. Ovary 2-celled. Styles 2. Fruit a 2-celled berry.

1. *P. HORRIDUM* Benth. & Hook. Flowers greenish.—½. Tamalpais. Summer.

2. *Aralia* L. SPIKENARD.

Ovary 5-celled. Styles 5. Fruit a drupe, with five pyrenæ.

1. *A. CALIFORNICA* Watson. Flowers greenish; berry dark purple or black.—24. Shady gulches. Summer.

Family 3. CORNACEÆ.

Stamens 4. Leaves opposite.

1. *Cornus* L. CORNEL. DOGWOOD.

Flowers ♂, Petals 4. Ovary 2-celled. Style 1. Fruit a drupe, with two pyrenæ.— $\frac{1}{2}$.

1. *C. NUTTALLII* Audubon. Flowers in a dense glomerule, supported by a conspicuous petaloid involucre; flowers greenish; fruit red.—Formerly in the vicinity of San Francisco, where it is now extinct. Still to be found on Bolinas Heights. Common in the Sierra.

2. *C. CALIFORNICA* C. A. Meyer. Flowers in a cyme, not involucre white; fruit white, sometimes blue.—Crystal Springs. Marin County. Summer.

2. *Garrya* Dougl. TASSEL TREE.

Diœcious. Flowers amentaceous, ternate between decussate bracts. Petals 0. Ovary 1-celled; styles two, persistent. Ovules two. Fruit a berry.— $\frac{1}{2}$.

1. *G. ELLIPTICA* Dougl. Leaves sessile, undulate; fruit purple.—Shady ravines around the bay. Spring.

2. *G. FREMONTII* Torr. Leaves petiolate, not undulate; fruit purple.—Wright's Station. Santa Cruz mountains. Spring. "Quinine

tree" of the settlers. All parts intensely bitter; even the berries, which in *G. elliptica* are sweet and edible, although the bark and leaves partake of the bitterness of the other species.

ORDER 2. CORNICULATÆ. Calyx gamosepalous; stamens equaling or double the number of the petals: always correspondent. Ovaries compound of several carpidia, each carpidium ∞ -ovulate.

Family 1. SAXIFRAGACEÆ.

Number of carpidia less than that of the petals. Fruit a capsule.

1. *Saxifraga* L. SAXIFRAGE.

Stamens 10. Pistils two. Ovary 2-celled. Fruit a loculicidal capsule.— \mathcal{L} .

1. *S. VIRGINIANA* Michx. Petals white.—Lagunitas Creek. Sunnyside. Spring.

2. *Boykinia* Nutt.

Stamens five. Pistils two. Ovary 2-celled. Fruit a 2-celled capsule.— \mathcal{L} .

1. *B. OCCIDENTALIS* Torr. & Gray. Petals white.—Lagunitas Creek. Summer.

3. *Tellima* R. Brown.

Stamens 10. Pistils 2-3. Petals lobed. Ovary 1-celled; styles short; stigmas capitate. Capsule valvularly dehiscent near the apex.— \mathcal{L} .

1. *T. GRANDIFLORA* R. Br. Petals laciniately pinnatifid, purple.—Saucelito. Tamalpais. San Miguel. Spring.

2. *T. BOLANDERI* Gray. Petals almost entire; styles smooth; calyx almost hypogynous; petals white.—Tamalpais. Spring.

3. *T. HETEROPHYLLA* Hook. & Arn. Petals obtusely 3-lobed; styles smooth; calyx almost hypogynous; petals white.—Common. Spring.

4. *T. AFFINIS* Bolander. Petals 3-dentate; styles granulose; calyx perigynous; petals white.—San Rafael. Spring.

6. *Tiarella* L.

Stamens 10. Pistils 2. Petals entire. Ovary 1-celled; styles long; stigmas simple. Capsule valvularly dehiscent to the base. Valves unequal. Placentæ parietal. Flowers white. 2♂.

1. *T. UNIFOLIATA* Hook.—Marin County. Spring.

5. *Heuchera* L. ALUM-ROOT.

Stamens 5. Pistils 2. Petals entire. Ovary 1-celled; styles long. Capsule valvularly dehiscent. Valves equal.—2♂.

1. *H. MICRANTHA* Dougl. Calyx pointed at the base, shorter than its pedicel.—Camp Taylor. Summer.

2. *H. PILOSISSIMA* Fisch. & Mey. Calyx rounded at the base; about the length of its pedicel.—Camp Taylor. Tamalpais. Crystal Springs. Summer.

Family 2. **RIBESIACEÆ.**

Stamens 5. Pistils 2. Fruit a berry. Leaves alternate.

1. *Ribes* L.

GOOSEBERRY. CURRANT.

Ovary inferior, 1-celled, with 2 parietal, ∞ -ovulate placentæ.— $\frac{1}{2}$.

1. *R. MENZIESII* Pursh. Thorny; vernation plicate; anthers sagittate; calyx tube campanulate, purple; petals white; berry prickly, yellowish.—Common. Spring. Fruit scarcely edible. “Prickly Gooseberry.”

2. *R. DIVARICATUM* Dougl. Thorny; vernation plicate; anthers not sagittate; calyx tube campanulate, dull colored; petals white; berry not prickly, dark purple.—Common. Spring. Fruit edible.

3. *R. SANGUINEUM* Pursh. Not thorny; vernation plicate; calyx tube cylindrical, rose-red; berry glandular, dark.—Common. Spring. “Wild Currant.” Fruit scarcely edible.

4. *R. AUREUM* Pursh. Not thorny; vernation convolute; flowers yellow.—Rare. Strawberry Valley. Wild-Cat Creek. Spring. “Golden Currant.”

Family 3. PHILADELPHEÆ.

Ovary inferior or half inferior. Number of carpidia disposed to correspond with calyx lobes. Fruit a capsule. Leaves opposite.

1. Whipplea Torr.

Stamens 5. Pistils 3-5. Ovary 3-5-celled. Cells 1-ovulate. Fruit a septicidal capsule. Flowers white.— $\frac{1}{2}$.

1. W. MODESTA Torr.—Redwoods. Summer.

Family 4. CRASSULACEÆ.

Calyx, corolla and stamens alternating. If the stamens form two circles the inner circle is opposite to the petals. Ovaries opposite the petals. Fruit follicles with central dehiscence.

1. Cotyledon L.

(*Echeveria* DC.)

Calyx 5-parted. Petals 5, coherent by their claws. Stamens 10.— $2\frac{1}{2}$. Fleshy herb.

1. C. CÆSPITOSA Haworth. Flowers yellow. Rocky places. Summer.

2. Sedum L. STONE CROP.

Petals 5, entirely free. Stamens 10.— $2\frac{1}{2}$. Fleshy herbs.

1. S. SPATHULIFOLIUM Hook. Leaves glaucous, obtuse, narrowed toward the base; flowers yellow. Rocks near the Presidio. Summer.

2. *S. STENOPETALUM* Pursh. Leaves acute, lanceolate; flowers yellow.—Camp Taylor. Summer.

3. *Tillæa* L.

Stamens as many as petals.—☉. Minute vernal aquatics.

1. *T. MINIMA* Miers. Flowers clustered, white.—San Miguel.

2. *T. ANGUSTIFOLIA* Nutt. Flowers solitary, white.—Mission Dolores.

Section 2. *CENTROSPERMÆ*. Central placentation from the base of the ovary.

ORDER 1. *FICOIDALES*. Anomalous, transitional types.

Family 1. *FICOIDEÆ*.

Calyx superior. Petals ∞ . Stamens ∞ -seriate, inserted with the petals. Ovary 4-20-celled, ∞ -ovulate. Fruit a capsule, ∞ -seeded. Stipules 0.

1. *Mesembrianthemum* L. FIG-MARYGOLD.

Characters of the family.

1. *M. ÆQUILATERALE* Haworth. Flowers red.—2f. Rocks near the seashore. Summer. Fruit edible.

ORDER 2. *CARYOPHYLLALES*. Flowers regular. Calyx inferior. Sepals as many as

petals. Stamens as many or twice as many. Ovary 1-celled, placenta central.

Family 1. SILENEÆ.

Calyx gamosepalous. Petals and stamens inserted into a carpophore. Stamens if equal to the petals alternating with them. Ovary ∞ -ovulate. Styles several. Fruit a capsule. Leaves opposite. Stipules 0.

1. Silene L.

CAMPION. CATCHFLY.

Calyx 5-dentate without scales at the base. Petals unguiculate. Stamens 10. Styles 3. Capsule dehiscent by teeth; seeds reniform.

1. *S. GALLICA* L. Villous; leaves spatulate; flowers nearly sessile, racemose; petals pale, entire, scarcely exceeding the calyx.—☉. Common. Summer.

2. *S. ANTIRRHINA* L. Glabrous; leaves lanceolate; flowers in a leafless, dichotomous panicle on long pedicels; petals obovate, equaling the calyx.—☉. Livermore. Summer.

3. *S. CALIFORNICA* Durand. Flowers few on the ends of branches; petals deeply parted, with bifid segments; scarlet.—☉. Oakland waterworks. Summer.

4. *S. LACINIATA* Cav. Flowers paniculate; petals deeply 4-cleft, with linear, acute lobes; bright scarlet.—☉. Contra Costa range. Summer.

5. *S. VERECUNDA* Watson. Petals bifid, rose-colored.— \mathcal{L} . Cemetery of San Francisco. Summer.

Family 2. **ALSINEÆ.**

Sepals distinct to the base. Carpophore 0. Stamens, if equaling the petals, alternating with them. Ovary ∞ -ovulate; styles several. Fruit a capsule. Leaves opposite; stipules 0.

1. ***Cerastium* L.**

MOUSE-EAR CHICKWEED.

Sepals 5. Petals 5, emarginate. Stamens 10. Stigmas 5. Capsule cylindrical, dehiscent by 10 teeth. Flowers white.

1. *C. ARVENSE* L. Petals about twice the length of the sepals.— \mathcal{L} . Common. All the year round.

This weed has, in Europe, the reputation of being poisonous to sheep.

2. *C. PILOSUM* Ledebour. Petals but little exceeding the sepals.— \mathcal{L} . Punta de los Reyes. Summer.

2. ***Stellaria* L.** CHICKWEED.

Petals 2-lobed. Stamens 10. Stigmas 3-5. Capsule globose, dehiscent by 6-10 valves. Flowers white.

1. *S. MEDIA* L. Flowers on slender pedicels.— \odot . Common. All the year round.

Ballast weed, introduced from Europe.

2. *S. NITENS* Nutt. Flowers on short pedicels.—☉. Tamalpais. Spring.

3. *Arenaria* L. SANDWORT.

Sepals unchanged in fruit. Petals entire. Stamens 5 or 10. Stigmas 3. Capsule ovoid, dehiscent by 3 valves, these valves sometimes 2-parted. Flowers white.

1. *A. DOUGLASHII* Torr. & Gray. Valves of the capsule entire; leaves filiform; sepals acute, 3-nerved.—☉. Tamalpais. Summer.

2. *A. CALIFORNICA* Brewer. Valves of the capsule entire; leaves lanceolate; sepals acute, 3-nerved.—☉. Oakland Hills. Spring.

3. *A. PALUSTRIS* Watson. Valves of the capsule entire; leaves linear; sepals obtuse, without nerves. Formerly in a marsh near San Francisco, where at present it is extinct. Has not been found in any other locality.

4. *A. MACROPHYLLA* Hook. Valves of the capsule 2-cleft; sepals acuminate, 1-nerved. ♀.—Saucelito. Summer.

4. *Sagina* L. PEARLWORT.

Sepals 4 or 5. Petals 4 or 5. Stamens twice 4 or twice 5. Stigmas 4 or 5, alternate with the sepals. Capsule dehiscent in 4 or 5 valves, opposite to the sepals.—☉.

1. *S. OCCIDENTALIS* Watson. Petals white. San Francisco. Spring.

Family 3. **PARONYCHIEÆ.**

Like *Alsineæ*, but the parts frequently defective or reduced in numbers. Leaves with scarious stipules.

1. *Spergula* L. CORN-SPURREY.

Stamens 5 or 10. Ovary ∞ -ovulate; styles 5, alternate with the sepals. Capsule 5-valved. Valves opposite to the sepals.— \odot .

1. *L. ARVENSIS* L. Flowers white.—San Francisco. Summer. Ballast weed, introduced from Europe.

1. *Spergularia* Pers. (*Lepigonum* Fries).

SAND-SPURREY.

Ovary ∞ -ovulate; style 3-5-cleft. Capsule 3-5-valved.

1. *S. MACROTHECA* Fisch. & Meyer. Leaves with large ovate stipules; flowers rose-colored.— \mathcal{L} . Margins of salt marshes. Summer.

2. *S. MEDIA* Fries. Leaves with short stipules; flowers white.— \odot . Contra Costa. Spring.

3. *Pentacæna* Bartl.

Divisions of the calyx unequal, persistent in fruit; the three external with cucullate apex, ending in a spine; two internal mucronate. Petals 5, minute. Stamens 3-5. Ovary 1-ovulate; style 2-cleft. Fruit a utricle.— \odot .

1. *P. RAMOSISSIMA* Hook. & Arn. Prostrate; pungent; everlasting.— $\frac{1}{4}$. Presidio. Cemetery. Spring.

Family 4. PORTULACACEÆ.

Flowers regular, but parts not corresponding in numbers.

1. *Portulaca* Tourn. PURSLANE.

Calyx tube connate with ovary. Limb 2-parted, free and circumscissile. Petals 4-6. Stamens 8 to ∞ . Capsule circumscissile.

1. *P. OLERACEA* L. Fleshy herb; petals yellow.— \odot . San Francisco. Spring. Summer. Escaped from cultivation.

2. *Calandrinia* HBK.

Sepals 2, persistent. Petals 5 to ∞ , equal. Stamens opposite to the petals, variable in number. Ovary ∞ -ovulate; style 3-cleft. Capsule 3-valved, ∞ -seeded.

1. *C. MENZIESII* Hook. Petals red.— \odot . Common. Spring.

3. *Claytonia* L MINER'S LETTUCE.

Sepals 2, persistent. Petals 5, equal. Stamens 5, opposite to the petals. Ovary 3 or 6-ovulate; style 3-cleft. Capsule 3-valved, 3-seeded.

1. *C. SIBIRICA* L. Inflorescence in loose, simple racemes. Herbaceous bracts with most

of the pedicels; single pair of cauline leaves distinct; leaves thin; petals rose-color.— \mathcal{C} . Saucelito. Spring.

2. *C. PERFOLIATA* Don. Inflorescence fasciculate; bracts few and minute; leaves thick, succulent; single pair of cauline leaves entirely connate; petals white.— \odot . Common. Spring. The whole plant edible as a salad.

3. *C. PARVIFLORA* Dougl. Inflorescence becoming at last racemose; bracts few and minute; leaves thick, succulent; single pair of cauline leaves imperfectly connate, sometimes on one side entirely distinct; petals pale rose-color.— \odot . Berkeley. Spring.

4. *C. SPATHULATA* Dougl. Inflorescence in a loose raceme; bracts few and minute; leaves thick, succulent; single pair of cauline leaves lanceolate but little connate, frequently entirely distinct; petals bright rose-color.— \odot . Tiburon. Tamalpais. Spring.

5. *C. CHAMISSONIS* Eschsch. Several pairs of cauline leaves; petals white.— \odot . Tiburon. Spring.

6. *C. LINEARIS* Dougl. Leaves alternate. Petals white.— \odot . Guerneville. Spring.

4. *Montia* L.

Sepals 2, persistent. External petals 3; internal petals 2, connate. Stamens 3 or 5. Ovary 3-ovulate; style 3-cleft. Capsule 3-valved, 3-seeded. Flowers white.— \odot .

1. *M. FONTANA* L. Flowers inconspicuous.—Sunnyside. Spring.

ORDER 3. CHENOPODIALES. Corolla 0, Perigonium inferior. Stamens opposite to the sepals. Ovary 1-celled, centrospermous. Stipules 0.

Family 1. AMARANTACEÆ.

Perigonium without tube, 3-bracteate; anterior bract longer than the two lateral ones. Style simple.

1. *Amarantus* L.

Polygamous, monœcious. Stamens distinct. Style short; stigmas 2 or 3. Fruit a circumscissile utricle.—⊙.

1. *A. RETROFLEXUS* L. Flowers green.—Cultivated grounds. Summer.

Family 2. CHENOPODIACEÆ.

Perigonium without tube, not more than 1-bracteate.

1. *Chenopodium* L. FIG WEED.

Bracts 0. Perigonium 5-cleft. Lobes dorsally carinate. Stamens 5. Ovary 1-celled, 1-ovulate; stigmas 2. Fruit a depressed utricle enclosed in the persistent perigonium.

1. *C. ALBUM* L. Leaves smooth, pruinose, rhombic, sinuate; the uppermost lanceolate,

entire; flowers densely clustered in dense spikes, forming a strict, close panicle.—⊙. Cultivated grounds. Summer.

Ballast weed, introduced from Europe.

2. *C. MURALE* L. Leaves shining, rhombic, dentate; flowers clustered in loose spicate panicles.—⊙. Cultivated grounds. Summer.

Ballast weed, introduced from Europe.

3. *C. AMBROSIODES* L. Leaves lanceolate, dentate; flowers clustered in slender axillary spikes.—⊙. Cultivated grounds. Summer.

Introduced from South America. Possesses anthelmintic properties.

4. *C. CALIFORNICUM* Watson. Leaves triangular, hastate; flowers densely clustered in simple terminal spikes.—2f. Cultivated grounds. Summer.

2. *Atriplex* L. ORACHE.

Flowers polygamous. ♂ and ♀ perigonium 3 or 5-parted. ♀ perigonium 2-parted. Stigmas 2. Fruit a compressed utricle.

1. *A. PATULUM* L. Leaves 3-angular, hastate; lower leaves opposite; bracts large, rhombic to hastate.—⊙. Cultivated grounds. Summer. Ballastweed, introduced from Europe.

2. *A. CORONATUM* Watson. All leaves alternate; bracts orbicular, surrounded by a herbaceous, dentate margin.—⊙. Contra Costa. Summer.

3. *A. LEUCOPHYLLUM* Dietr. All leaves alternate; fruiting bracts spongy, rhombic to ovate. ☉.—On the seashore. Summer.

4. *A. CALIFORNICUM* Mag. Lower leaves opposite, sessile; stem and leaves furfuraceous; fruiting bracts membranaceous; rhombic.—☿. On the seashore. Summer.

3. *Salicornia* TOURN. SAMPHIRE.

Flowers in threes, immersed in a rachis, decussately arranged and forming a spike. Perigonium gamophyllous, saccate. Stamens 1 or 2. Fleshy, articulate, aphyllous, saline plants.

1. *S. AMBIGUA* Michx.—☿. Salt marshes. Summer.

4. *Suaeda* FORSK. SEA BLITE.

Flowers axillary, with minute, scale-like bracts. Perigonium urceolate, 5-cleft. Stamens 5. Fruit a utricle, enclosed in the inflexed perigonium. Leaves terete, fleshy.

1. *S. CALIFORNICA* Watson—☿. Salt marshes on an island near Alameda. Summer.

Family 3. *NYCTAGINEÆ*.

Flowers involucrate. Perigonium gamophyllous, corolla-like, its persistent tube enclosing the akene.

1. *Abronia* Juss.

Involucre 5-leaved, ∞ -flowered, persistent. Perigonium salver-shaped. Stamens 5, included. Stigma clavate. Leaves opposite.

1. *A. UMBELLATA* Lam. Leaves attenuate into a slender petiole; perigonium rose-colored.— φ . Seashore. Summer.

2. *A. LATIFOLIA* Esch. Petioles distinct, but short; perigonium yellow.— φ . Seashore. Summer.

Section 3. EUCYCLICÆ. Floral parts distinct in well defined circles.

ORDER 1. TRICOCCÆ. Flowers diclinous. Ovary superior. Ovules 1-2 collateral, pendulous from the summit of cells, which separate at last from a central axis.

Family 1. EUPHORBIACEÆ.

Cells 1-ovulate.

1. *Eremocarpus* Benth.

Monœcious. Flowers cymose. δ : perigonium 5-parted; stamens 6 or 7, central, inflexed in æstivation. φ : perigonium 0; ovary with 5 glands at the base, 1-celled, 1-ovulate; style simple; capsule 2-valved.— \odot .

1. *E. SETIGER* Benth.— \odot . Marin County. Niles. Summer.

The herb has the smell of strawberries, but is poisonous. The crushed leaves are used by the Indians to catch fish by stupefying them.

2. **Hendecandra** Esch.

(*Croton* L.)

Diœcious. ♂ : flowers in racemes; perigonium 5-cleft, with 5 glands opposite to the lobes; stamens more than 5. ♀ flowers: perigonium, 5-cleft; glands 0; ovary 3-celled; styles 3, each 4-parted; capsule 3-coccus, each coccus 2-valved.

1. *H. PROCUMBENS* Esch. (*Croton Californicus* Mull. Arg.)—2f. Common in sandy soil. Summer. Drastic poison.

3. **Euphorbia** L. SPURGE.

Monœcious. Androgynous. Flowers umbellate. Involucre campanulate. ♂ : flowers, perigonium 0; monandrous, stipitate; bracteate. ♀ : flower solitary in the center of the umbel; perigonium dentate or 0; ovary 3-celled; styles 3, each 2-cleft.

1. *E. OCELLATA* Dur. & Hilg. Glands of the involucre petaloid; leaves all opposite, entire.—⊙. Sonoma. Millbrae. Summer.

2. *E. SERPYLLIFOLIA* Pers. Glands of the involucre with a narrow, white margin; leaves all opposite; base oblique; apex denticulate. ⊙.—Marin County. Summer.

3. *E. LEPTOCERA* Engelm. Glands of the involucre crescent-shaped, without colored margin; cauline leaves scattered, those of the branchlets opposite.—⊙. Common. Summer.

4. *E. LATHYRIS* L. Glands of the involucre crescent-shaped, without colored margin; leaves all opposite.—⊙⊙. Common. Summer.

Poisonous plant introduced from Europe and found everywhere. The seeds contain an oil analogous to that of croton. This plant is said to drive away or kill rats, as also the castor-oil plant. *Ricinus communis*.

ORDER 2. MALVALES. Flowers regular. Calyx free, 5-parted, valvate in æstivation. Corolla 5-parted, contorted in æstivation. Stamens often monadelphous. Carpidia several, free, or connate with the central axis into an ∞-celled ovary. Leaves alternate, stipulate.

Family 1. MALVACEÆ.

Claws of petals united with each other, and with the column of ∞, monadelphous stamens. Anthers reniform, 1-celled.

1. *Lavatera* L. TREE MALLOW.

Involucre 3-6-cleft. Ovaries ∞, verticillate; 1-ovulate. Style 1, springing from the central axis; stigmas ∞, filiform. Fruit a schizocarp; seeds ascending.

1. *L. ASSURGENTIFLORA* Kellogg. Petals rose-colored. — ½. Near cultivated places. Summer.

Native of the island of Anacapa, but having escaped cultivation, now spontaneous in different localities.

2. *Malva* L. MALLOW.

Involucre 3-leaved. Ovary ∞ -celled; cells 1-ovulate. Styles as many as cells, united at base; stigmas obtuse. Fruit a depressed capsule; seed ascending.

1. *M. BOREALIS* Wallman. Flowers pale.—
⊙. Cultivated grounds. Summer. Ballast weed from Europe.

3. *Sidalcea* Gray.

Involucre 0. Ovary ∞ -celled; cells 1-ovulate. Styles as many as cells, united at base. Fruit 5-10-coccous schizocarp; seed ascending.

1. *S. HUMILIS* Gray (*malvæflora?* Gray). Raceme long and loose; petals rose or purple. ¼. Common. Summer.

2. *S. DIPLOSCYPHA* Gray. Flowers in umbellate clusters; pedicels subtended by persistent, hispid bractlets; petals rose or purple. ⊙. Suñol. Summer.

3. *S. MALACHROIDES* Gray. Flowers in close, racemose, nearly spicate clusters; petals small, white.—⊙. Marin County. Summer.

4. **Malvastrum** Gray (*Sphæralcea* St. Hilaire).

Ovary ∞ -celled; cells 2-ovulate. Styles as many as cells, united at base; stigmas capitate. Fruit a loculicidal capsule, at length also septicidal.

1. **M. THURBERI** Gray, Petals pale.— $\frac{1}{2}$. San Mateo. Niles. Summer.

5. **Sida** Kunth.

Ovary ∞ -celled; cells 1-ovulate. Styles as many as cells, united at base; stigmas capitate. Fruit a ∞ -seeded capsule. Seeds pendulous from the apex.— $2\frac{1}{2}$.

1. **S. HEDERACEA** Torr. Flowers pale. Common. Summer.

Order 3. **GERANIALES**. Calyx free, imbricated in æstivation. Petals 5, contorted or convolute in æstivation; stamens hypogynous; definite. Ovary the compound of a definite number of carpidia.

Family 1. **GERANIACEÆ**.

Sepals 5. Petals 5; hypogynous; alternate with the sepals; stamens 10. Carpidia 5, verticillate round a columnar axis (gynophore). Styles distinct at base; connate towards their apex. Carpidia 5, 2-ovulate, 1-seeded, ventrally dehiscent. Leaves stipulate.

1. *Geranium* L.

Stamens monadelphous, all fertile. Styles persistent.

1. *G. CAROLINIANUM* L. Corolla rose-color. —⊙. Common. Summer.

2. *Erodium* L'Her.

STORKSBILL. PIN-CLOVER.

Stamens monadelphous, alternately sterile or depauperate. Styles persistent.

1. *E. CICUTARIUM* L'Her. Leaves pinnate; leaflets pinnatifid, with narrow acute lobes; flowers rose-colored. —⊙. Common. Summer. Fodder plant, called *Alfilerilla* (*alfiler*, in Spanish, needle) from the shape of the fruit. If the plant is not indigenous, it must have been introduced for a considerable time, or else it would not have a distinct name from the old settlers.

2. *E. MOSCHATUM* L'Her. Leaves pinnate; leaflets ovate, doubly serrate; stipules conspicuous; flowers rose-color. —⊙. Cultivated grounds. Summer. Ballast weed, introduced from Europe.

3. *E. BOTRYS* Bertol. Leaves pinnatifid, lobes dentate, stipules small; petals pale. —⊙. Cultivated grounds. Summer.

4. *E. MACROPHYLLUM* Hook. & Arn. Leaves reniform, lobed. —⊙. Niles. Summer.

Family 2. OXALIDEÆ.

Sepals 5. Petals hypogynous. Ovary 5 carpidia, opposite the petals, attached to the axis by their central angle, 2-to- ∞ -ovuled; ovules vertically arranged. styles 5, persistent. Leaves alternate.

1. Oxalis L. WOOD-SORREL.

Petals 5. Stamens 10, monadelphous, the 5 opposite the petals shorter. Capsule oblong.

1. *O. OREGANA* Nutt. Peduncles 1-flowered; petals white or rose-color.— \mathcal{L} . Crystal Springs. Camp Taylor. Spring.

2. *O. CORNICULATA* L. Peduncles two to several-flowered; petals yellow.— \mathcal{L} . Common. Spring.

Family 3. LINEÆ.

Calyx persistent. Petals hypogynous, unguiculate. Perfect stamens 5, sometimes 4. Ovary 5, sometimes 4, 3 or 2-celled; cells 2-ovulate. Ovules collateral, more or less separated by a spurious septum. Styles equalling the cells in number. Capsule globose.

1. Linum. FLAX.

Sepals 5. Petals 5.

1. *L. PERENNE* L. Pentagynous; petals blue.— \mathcal{L} . Colma. Millbrae. Summer.

2. *L. BREWERI* Gray. Trigynous; pedicels short; stipular glands conspicuous; flowers almost solitary; petals yellow.—⊙. Contra Costa hills. Summer.

3. *L. CONGESTUM* Gray. Trigynous; pedicels short; stipular glands very small; sepals pubescent; flowers in terminal fascicles; petals dark-red.—⊙. Tamalpais. Summer. Rare.

4. *L. CALIFORNICUM* Benth. Trigynous; pedicels short; stipular glands conspicuous; upper flowers cymose, lower flowers solitary; petals rose-colored.—⊙. San Francisco cemetery. San Rafael. Summer.

5. *L. SPERGULINUM* Gray. Trigynous; pedicels conspicuous; stipular glands 0; flowers rose-colored or white.—⊙. Marin County. Summer.

Family 4. *LIMNANTHÆ*.

Calyx persistent, valvate in æstivation. Petals alternating with calyx lobes, inserted into a perigynous disc. Stamens twice as many as petals. *Carpidia* verticillate, free, 1-ovulate; style central on the apex, 5, sometimes 3-cleft. Fruit 5, sometimes 3 akenes. Stipules 0.

1. *Limnanthes* R. Br.

Calyx 5-parted. Petals 5, cuneiform. Ovaries 5. Fruit 5, rugose akenes.—⊙.

1. *L. DOUGLASHII* R. Br. Sepals glabrous; petals white or rose-colored, claws always yellow. Sunnyside. Summer.

2. *L. ALBA* Hartweg. Sepals villous; petals white, claws yellow.—Millbrae. Summer.

Order 4. **TEREBINTHALES.** Flowers regular. Calyx free. Petals 5, imbricate or valvate in æstivation, not contorted, nor truly convolute, inserted into a disc. Stamens 5, or a multiple. Ovary 1-5 carpidia, syncarpous or apocarpous; carpidia 1-2-ovulate.

Family 1. RUTACEÆ.

Stamens inserted on the external margin of the disc. Carpidia more than 1-ovulate. Stipules 0.

1. *Ptelea* L. HOP-TREE.

Polygamous. Calyx 4-5-parted. Petals 4 or 5. Stamens 4 or 5, alternate with the petals. Ovary on a convex disc, 2-celled; cells 2-ovulate; style 1. Fruit a 2-seeded samara. Flowers white or greenish.— $\frac{1}{2}$.

1. *P. ANGUSTIFOLIA* Benth.—Niles. Spring.

Family 2. TEREBINTHACEÆ. (*Anacardiaceæ*).

Stamens inserted on the inner margin of the disc. Ovary 1, or if more, only one fertile; 1-ovulate. Fruit indehiscent.

1. *Rhus* L.

POISON-OAK. YEDRA.

Polygamous. Calyx 5-parted, persistent. Stamens 5. Ovary one, 1-celled, 1-ovulate; styles 3. Fruit a dry drupe.

1. *R. DIVERSILOBA* Torr. & Gray. Flowers paniculate; petals greenish. Fruit pale.— $\frac{1}{2}$. Common. Summer.

2. *P. AROMATICA* Ait. Flowers spicate; petals yellow; fruit red.— $\frac{1}{2}$. Livermore. Spring.

Juice and exhalation of this shrub unlike those of *R. diversiloba* do not produce eczema, but are entirely harmless.

Family 3. JUGLANDEÆ.

(Connecting link between *Terebinthales* and *Cupuliferae*.)

Flowers diclinous. ♂ : amentaceous; perigonium adnate to the bract, and imbricate in æstivation; petals 3, or its multiple. ♀ : aggregate or racemose; perianth connate with the imperfectly 2-4-celled, 1-ovulate ovary; fruit a 1-pyrenous drupe; pyrena of valvular dehiscence. Leaves imparipinnate, stipules 0.

Juglans L. WALNUT.

♂ : stamens more than six. ♀ flowers: few, terminal; calyx 4-parted; petals 4; style 2.— $\frac{1}{2}$.

1. *J. CALIFORNICA* Watson. — Very local. Walnut Creek. Spring.

ORDER 5. SAPINDALES. Parts of the androecium not symmetrical. Calyx free, imbricate in æstivation. Petals inserted into a hypogynous disc. Stamens generally more than petals, but not their multiple. Carpidia 3, sometimes 2, more or less connate into an ovary. Ovules of definite number.

Family 1. POLYGALACEÆ.

Flowers irregular, often imitating *Papilionaceæ*. Sepals 5, the two lateral petaloid (wings). Petals 5, sometimes 3, connate with the staminal tube. Anterior petal concave (carina). Stamens 8, sometimes 4 or less, usually monadelphous. Ovary 2-celled; style 1. Stipules 0.

1. *Polygala* L.

Calyx persistent. Stamens 8, ascending. Filaments united at base into an anteriorly cleft tube. Ovary 2-celled; cells 1-ovulate. Capsule loculicidal.

1. *P. CUCULLATA* Benth. Flowers rose-color.—24. Tamalpais. Crystal Springs. Summer.

Family 2. SAPINDACEÆ.

Sepals 5, often irregular and more or less connate. Disc fleshy. Petals alternate with sepals, and appendiculate at their claws—sometimes one or all of them wanting. Sta-

mens 1-seriate, more than 5. Ovary generally 3-celled; cells 1-2 ovulate.

1. *Æsculus* L. BUCKEYE.

Polygamous. Calyx irregular, gamosepalous. Petals unguiculate, 5, or by abortion of the anterior one, 4. Stamens 6-8, filaments ascending. Cells of ovary 2-ovulate. Fruit a capsule, generally by abortion 1-celled, 1-seeded. Leaves palmate, deciduous.— $\frac{1}{2}$.

1. *Æ. CALIFORNICA* Nutt. Petals white, tinged with red and yellow.—Common. Summer.

The seeds contain an active principle not yet sufficiently examined.

2. *Acer* Mœnch. MAPLE.

Polygamous. Petals as many as sepals, inserted into the margin of the lobate, hypogynous disc. Stamens 8, inserted with the petals. Ovary 2-celled, 2-lobed; cells 2-ovulate. Fruit a 2-seeded samara.— $\frac{1}{2}$. Leaves opposite, palmately lobed, deciduous.

1. *A. MACROPHYLLUM* Pursh. Flowers yellow.—San Mateo. Marin County. Spring.

3. *Negundo* Mœnch. BOX ELDER.

Polygamous. Dioecious. Petals 0. ♂ : stamens 4, sometimes 5, opposite to the calyx teeth. ♀ : ovary 2-celled, 2-lobed; cells 2-ovu-

late; fruit a 2-seeded samara. Leaves opposite, imparipinnate, deciduous.— $\frac{1}{2}$.

1. *N. CALIFORNICUM* Torr. & Gray.—San Mateo. Marin County. Spring.

ORDER 6. CELASTRALES. Parts of the flower correspondent in number. Disc tumid, adnate to the base of the calyx. Stamens inserted with the petals on the margin of the disc. Ovules definite.

Family 1. RHAMNACEÆ.

Calyx conspicuous, gamosepalous, valvate in æstivation. Stamens inserted with the petals and alternate with the calyx lobes. Ovary 2-3-celled; cells 1-ovulate. Styles 2-3, more or less connate.

1. *Rhamnus* Juss.

Calyx-tube urceolate, margin 4-5-cleft. Petals minute, with short claws, or 0. Ovary free. Fruit a 2-3-pyrenous drupe. Leaves alternate, stipulate.— $\frac{1}{2}$.

1. *R. CROCEA* Nutt. Flowers 4-merous, apetalous; fruit red.—Lake Chabot. San Rafael. Taylorville. Spring.

2. *R. CALIFORNICA* Esch. Flowers 5-merous; petals small, ovate, emarginate; ripe fruit black.—Common. Summer.

“*Cascara sagrada*” of the old Californians. An infusion of the bark of this species and of *R.*

Purshiana of Northern California was in use among them as a purge. There also was a belief that the infusion of the bark, if the branch was scraped towards the top, would act as an emetic.

An *infuso-decoct.* of the leaves is one of the most reliable remedies against fresh cases of the eczema caused by poison oak.

2. *Ceanothus* L.

CALIFORNIA LILAC. CHAPPARAL.

Calyx-tube hemispherical, concave; margin petaloid. Petals unguiculate; margin cucullate. Ovary half immersed into the disc; style 3-cleft. Fruit a 3-seeded capsule with base embraced by the calyx tube, dehiscent at the inner angle of the cells. Leaves simple.— $\frac{1}{2}$.

1. *C. THYRSIFLORUS* Eschsch. Leaves all alternate, 3-nerved from their base; flowers in dense, compound racemes, blue, fragrant.—Common. Summer.

2. *C. DENTATUS* Torr. & Gray. Leaves all alternate, frequently fascicled, 3-nerved from the base; flowers in small fascicles on naked, terminal peduncles, blue.—Tamalpais. Summer.

3. *C. DIVARICATUS* Nutt. Leaves all alternate, 3-nerved from their base; flowers in racemes, pale; branches spinose.—Niles. Livermore. Summer.

4. *C. PAPILLOSUS* Torr. & Gray. Leaves all alternate, pinnately veined, glandular; flowers blue.—Redwood City. Alma. Summer.

5. *C. CRASSIFOLIUS* Torr. Leaves often opposite, with numerous straight veins, oblong; young branches tomentose; flowers in dense fascicles, pale.—Wright's Station. Summer.

6. *C. CUNEATUS* Nutt. Leaves often opposite, cuneate, with numerous straight veins; young branches almost smooth; flowers in loose fascicles, pale.—Marin County. Summer.

Family 2. CELASTRACEÆ.

Calyx conspicuous, gamosepalous. Stamens alternate with the petals and opposite to the calyx lobes.

1. *Euonymus* Tourn. SPINDLE-TREE.

Sepals and petals 4 or 5, spreading. Ovary 3-4-celled, half immersed into the disc. Capsule loculicidal.

1. *E. OCCIDENTALIS* Nutt. Flowers dark brown.— $\frac{1}{2}$. Crystal Springs. Taylorville. Summer.

Family 3. AMPELIDEÆ.

Calyx inconspicuous. Stamens inserted with the petals, opposite and of equal number.

Ovary 3-celled; cells 2-ovulate; ovules collateral. Fruit a berry. Lower leaves opposite; upper by the transformation of one of the pair into a tendril, apparently alternate.

1. *Vitis* Tourn. GRAPE.

Petals coherent at their apex, caducous.

1. *V. CALIFORNICA* Benth.— $\frac{1}{2}$.

ORDER 7. GUTTIFERALES. Flowers regular. Calyx free, sepals 4 or 5, imbricate or 2-seriate, never regularly valvate in æstivation. Petals convolute in æstivation, sometimes 0. Stamens ∞ , more or less connate at base, sometimes reduced in number. Carpidia several, connate into an ovary.

Family 1. HYPERICINEÆ.

Sepals 2-seriate. Petals as many as sepals and alternate. Carpidia 3 or 5. Ovules ∞ , 2-seriate in each. Styles 3 or 5, more or less united. Fruit a septicidal capsule. Leaves opposite. Stipules 0.

1. *Hypericum* L. ST. JOHN'S-WORT.

Stamens polyadelphous; stigmas capitate. Flowers yellow.— $\frac{1}{4}$.

1. *H. CONCINNUM* Benth. Leaves not clasping; stamens decidedly polyadelphous in three phalanges; petals longer than the sepals.—Tamalpais. Camp Taylor. Crystal Springs. Summer.

2. *H. ANAGALLOIDES* Cham. & Schlecht. Leaves clasping; stamens rather polyandrous; petals somewhat shorter than the sepals.— Presidio. Saucelito. Lagunitas. Summer.

Family 2. SALICINEÆ

Dioecious. Amentaceous. ♂ : perigonium 0; stamens central. ♀ : perigonium 0; ovary 2 carpodia, connate into one; ovules ∞; styles 2; capsule 1-celled, 2-valved, ∞-seeded; seeds with hairy arillus. Leaves deciduous, alternate, stipulate.

1. *Salix* Tourn. WILLOW.

Bracts entire.

1. *S. LÆVIGATA* Bebb. Stamens more than 2; petioles downy; aments coeval with leaves.

½. Alms House. Spring.

2. *S. LASIANDRA* Benth. Stamens more than 2; petioles glandular at their upper ends; aments coeval with leaves.— ½. Alameda. Spring.

3. *S. LONGIFOLIA* Muhl. Diandrous; scales deciduous; aments coeval with leaves.— ½. Common. Spring.

4. *S. LASIOLEPIS* Benth. Diandrous; scales persistent, darker at the apex; capsules acute, smooth, dark green; aments precocious.— ½. Common. Spring.

5. *S. COULTERI* Anders. Monandrous; aments in the axils of persistent leaves.— $\frac{1}{2}$. Marin County. Spring.

2. *Populus* Tourn.

POPLAR. COTTONWOOD.

Bracts fimbriate, laciniate.— $\frac{1}{2}$.

1. *P. TRICHOCARPA* Torr. & Gray.— $\frac{1}{2}$. Marin County. Niles. Spring.

ORDER 8. PARIETALES. Ovary 1-celled; placentation parietal; ovules ∞ .

Family 1. CISTINEÆ.

Calyx 2-seriate; 2 external sepals smaller than the 3 internal, contorted to the left in æstivation, sometimes 0. Petals 5, caducous, contorted to the right in æstivation. Stamens ∞ . Ovary free, formed by several connate carpidia; style 1. Fruit a capsule; placentation sometimes apparently axillary, by placentæ protruding towards the center.

1. *Helianthemum* Tourn. ROCK-ROSE.

Ovary 3-valved. Placentæ in the middle of the valves; each valve more than 2-seeded. Flowers yellow.

1. *H. SCOPARIUM* Nutt.— $\frac{1}{4}$. Tamalpais. Summer.

Family 2. LOASACEÆ.

Calyx-tube more or less connate with the

ovary, costate; margin 4 or 5-parted. Petals 4 or 5, and their multiples. Stamens ∞ , some of the external sometimes petaloid. Ovary inferior, 1-celled; ovules ∞ ; style 1. Fruit a capsule immersed in the calyx tube.

1. **Mentzelia** L.

Calyx limb 5-parted, persistent. Petals not cucullate. Stamens ∞ , inserted in the throat of the calyx below the petals. Style 3-cleft. Capsule dehiscent near the apex.

1. *M. LINDLEYI* Torr. & Gray. Petals obovate, abruptly acuminate, yellow.— \odot . Niles. Suñol. San Leandro. Summer.

2. *M. LÆVICAULIS* Torr. & Gray. Petals lanceolate, cream-colored.— $\odot \odot$. Arroyo del Monte, near Livermore. Summer.

Family 3. **DATISCEÆ.**

Calyx tube connate with ovary. Petals 0. Ovary 1-celled; placentæ on the middle nerve of the carpidia; ovules ∞ ; styles on the margins of the carpidia, 2 for each placenta, sometimes united into a contiguous style. Fruit a capsule, crowned by the calyx margin and style.

1. **Tricerastes** Presl. (*Datisca* L).

Flowers ♀ . Calyx-tube 3-angular; margin 3-dentate. Stamens 3, alternate with the lobes of the calyx; anthers extrorse. Ovary with

3 placentæ; styles 3, 2-parted, opposite the calyx-teeth. — 2f.

1. *T. GLOMERATA* Benth & Hook.—Sunny-side. Niles. Summer.

Family 4. **FRANKENIACEÆ.**

Flowers regular. Calyx gamophyllus free, tubular, costate, persistent. Petals alternate with the lobes of calyx, unguiculate. Stamens not correspondent in number to petals; anthers extrorse. Ovary 1-celled. Fruit a capsule 2-4-valved, included in the persistent calyx-tube.

1. *Frankenia* L.

Leaves opposite. Stipules 0.—2f.

1. *F. GRANDIFLORA* Cham. & Schlecht. Petals pink.—Seashore. Summer.

Family 5. **VIOLACEÆ.**

Sepals 5, persistent. Petals 5, unguiculate. Stamens inserted into a hypogynous disc, filaments flat; anthers short, introrse, pressed against the ovary, the connective elongated beyond the anthers. Ovary free, 1-celled, formed by 3 carpodia; ovules ∞ , 2-seriate; style 1, persistent; fruit a 3-valved capsule; valves bearing the seeds on the middle nerve. Leaves stipulate.

1. *Viola* L. VIOLET.

Sepals unequal, appendiculate at base. Petals unequal, the inferior longest and calcarate. Stamens 5; anthers cohering into a tube, the two inferior ones appendiculate at their base.

1. *V. ODORATA* L. Stemless; petals violet.
 2. Throckmorton Ranch. Spring.

Sweet violet; probably escaped from cultivation.

2. *V. CANINA* L. Stem leafy; spur as long as the sepals; petals violet.—2. Cemetery, Saucelito. Spring.

3. *V. OCELLATA* Torr. & Gray. Stem erect, leafy. Leaves cordate; stipules small, scarious; spur merely saccate; petals gradually changing color from purple to white.—2. Tamalpais. Santa Cruz Mountains. Summer.

4. *V. PEDUNCULATA* Torr. & Gray. Stem leafy, ascending; leaves rhombic; stipules narrowly lanceolate, foliaceous; sepals lanceolate; spur merely saccate; petals yellow, the outside tinged with purple.—2. Common. Spring.

5. *V. AUREA* Kellogg. Stem leafy, ascending; leaves lanceolate; stipules lanceolate, foliaceous; sepals linear, acuminate; spur merely saccate; petals yellow.—2. Contra Costa mountains. Summer.

6. *V. SARMENTOSA* Dougl. Stem prostrate, leafy; leaves cordate, finely crenate; petals yellow.— ζ . Tamalpais. Taylorville. Summer.

7. *V. LOBATA* Benth. Erect leafy stem; cauline leaves palmate; petals yellow, externally tinged with purple.— ζ . Marin County. Summer.

Series 2. *APHANOCYCLICÆ*. Floral parts showing a tendency to spiral arrangement, and generally distinct. Parts of the gynœcium only occasionally connate. Calyx and corolla often of the same or similar structure. Numerical law not yet established. Stamens generally more numerous than sepals and petals.

ORDER 1. *CRUCIFLORÆ*. Calyx and corolla formed from 2 and its multiples. Circles of stamens more than 1. Carpidia 2 and its multiples.

Family 1. *CRUCIFERÆ*.

Sepals 2 pairs. Petals 4. Stamens tetradynamous. Ovary free, compound of 2 lateral carpidia; placentæ parietal, on the margins of a spurious septum (replum); style 1. Fruit a siliqua, silicula or lomentum. Leaves alternate; stipules 0.

1. *Raphanus* TOURN. RADISH.

Indehiscent siliqua irregularly moniliform, several-seeded.

1. *R. SATIVUS* L.—⊙. Common all the year round.

Escaped from gardens. Originally a native of China, now in cultivation over the whole earth.

2. *Thysanocarpus* Hook.

Indehiscent silicula (samara) orbicular, winged, 1-seeded.—⊙. Spring.

1. *T. PUSILLUS* Hook. Pubescent throughout; cauline leaves lanceolate, sessile, but not clasping; silicula orbicular, hirsute.—Niles.

2. *T. LACINIATUS* Nutt. Cauline leaves linear, scarcely auricled at their clasping base; silicula obovate, slightly pubescent.—Niles.

3. *T. CURVIPES* Hook. Glabrous; cauline leaves conspicuously auricled at the clasping base; silicula orbicular.—Common.

3. *Senebiera* Poir.

Indehiscent silicula (schizocarp) 2-seeded. Seeds globose, rugose. Flowers minute, white.

1. *S. DIDYMA* Pers.—⊙. Common. Summer.

4. *Lepidium* R. Br. PEPPERGRASS.

Silicula dorsally compressed, ovate, with carinate valves and narrow replum. Seeds solitary in each cell, pendulous from the apex of the replum.

1. *L. NITIDUM* Nutt. Silicula shining, glabrous; petals whitish.—⊙. Common. Spring.

2. *L. LATIPES* Hook. Silicula strongly reticulated; petals greenish.—⊙. Marin County. Summer.

5. *Capsella* Vent. SHEPHERD'S-PURSE.

Silicula dorsally compressed, with carinate, 1-nerved valves and narrow replum. Seeds several. Flowers white.—⊙.

1. *C. BURSA-PASTORIS* L.—Common all the year round.

Ballast weed, introduced from Europe.

6. *Tropidocarpum* Hook.

Siliqua dorsally compressed, linear, with 1-nerved, carinate valves, and very narrow replum. Seeds ∞. Flowers yellow.—⊙

1. *T. GRACILE* Hook.—Marin county. Summer.

7. *Nasturtium* R. Br. WATER-CRESS.

Siliqua linear, with concave valves, and only rudimentary nerve. Seeds ∞, 2-seriate in each valve.—Aquatic.

1. *N. OFFICINALE* R. Br. Petals white.—2f. Common all the year round.

Introduced from Europe, at present perfectly naturalized.

8. *Sisymbrium* L. HEDGE-MUSTARD.

Siliqua linear, with concave, 3-nerved valves. Seeds ∞ , 1-seriate in each valve.

1. *S. OFFICINALE* Scop. Branches divaricate; siliqua closely appressed to the axis of the raceme; petals yellow.— \odot . Common all the year round.

Ballast weed from Europe.

2. *S. ACUTANGULUM* DC. Branches ascending; siliqua erect; petals yellow.— \odot . Mission Dolores. San Jose. Summer.

Ballast weed, native of the Mediterranean region.

3. *S. REFLEXUM* Nutt. Branches but few; siliqua deflexed; petals yellow, white or red.— \odot . Presidio. Saucelito. Alameda. Summer.

9. *Cakile* Tourn. SEA-ROCKET.

Fruit a 2-articulate lomentum.

1. *C. AMERICANA* Nutt. Petals rose-colored. \odot . Berkeley salt marshes. Summer.

10. *Barbarea* R. Br. WINTER-CRESS.

Siliqua linear, 4-angular with concave, 1-nerved valves. Seeds ∞ , compressed, 1-seriate in each valve. Flowers yellow.

1. *B. VULGARIS* R. Br.— σ . San Mateo. Summer.

11. *Brassica* L. MUSTARD.

Siliqua linear, with concave, more than 1-nerved valves. Seeds ∞ , globose, 1-seriate in each valve. Flowers yellow.

1. *B. CAMPESTRIS* L. Upper leaves clasping, auriculate.— \odot . Common. Summer.

Native of Europe, but escaped from cultivation and naturalized.

2. *B. NIGRA* Boiss. All leaves petioled.— \odot . Very common. Summer.

Black mustard. Probably like the former escaped from cultivation, but at present a troublesome weed, plentiful enough to impart a disagreeable taste to milk and honey.

12. *Erysimum* L.

Siliqua linear, 4-angular, with carinate, 1-nerved valves. Seeds ∞ , globose, 1-seriate in each valve. Flowers yellow.

1. *E. ASPERUM* DC. $\odot\odot$. Common. Summer.

13. *Caulanthus* Watson.

Sepals equally saccate. Petals undulate. Siliqua elongated, terete, sessile; valves 1-nerved. Seeds ∞ , oblong, 1-seriate in each valve. $\odot\odot$.

1. *C. PROCERUS* Watson. Petals greenish white. Common. Summer.

14. *Cheiranthus* L. WALL-FLOWER.

Lateral sepals saccate. Petals unguiculate. Siliqua laterally compressed; replum broad; valves 1-nerved. Seeds compressed, 1-seriate in each valve.

1. *C. ASPER* Cham. & Schlecht. Petals in different localities of different coloration, viz., near San Francisco cream-color, at San Mateo orange-color, in Saucelito light yellow.—⊙⊙. Common. Spring.

15. *Streptanthus* Nutt.

Sepals colored and equal. Petals undulate. Siliqua laterally compressed; valves flattened, 1-nerved. Seeds ∞ , 1-seriate, marginate, compressed.

1. *S. GLANDULOSUS* Hook. Hispid; lateral pair of sepals not turned inward behind the upper petals; sepals and petals white or red.—⊙. Sunnyside. Spring.

2. *S. PERAMENUS* Greene. Hispid; lateral pair of sepals turned inward behind the upper petals; sepals dark purple, petals pale.—⊙. Wright's Station. Oakland Hills. Summer.

3. *S. NIGER* Greene. Glabrous, glaucous; sepals obtuse, dark purple; petals pale with purple claws.—⊙. Tiburon. Spring.

16. *Arabis* L. ROCK-CRESS.

Petals not undulate. Siliqua laterally com-

pressed, linear; valves flattened, with prominent nerve. Seeds ∞ , 1-seriate, compressed, marginate.

1. *A. BLEPHAROPHYLLA* Hook & Arn. Leaves conspicuously ciliate; cauline leaves oblong, sessile; petals purple.— \mathcal{L} . Presidio. Saucelito. Spring.

2. *A. PERFOLIATA* Lam. (*Turritis glabra* L.) Glauous; cauline leaves ovate, clasping by the sagittate base; petals pale.— \odot \odot . Common. Summer.

17. *Cardamine* L. BITTER-CRESS.

Siliqua laterally compressed, linear; valves flat, without nerve. Seeds ∞ , 1-seriate; funicle filiform.

1. *C. PAUCISECTA* Benth. Lower leaves simple or 3-lobed, cauline leaves 5-foliolate.— \mathcal{L} . Common. Spring.

2. *C. OLIGOSPERMA* Nutt. All leaves pinnate. \odot . Saucelito. San Mateo. Spring.

18. *Alyssum* L.

Silicula laterally compressed, orbicular; valves concave; replum broad. Seeds 1 or 2 in each valve.

1. *A. CALYGINUM* L. Petals white or yellow, but little exceeding the sepals.— \odot . San Francisco. Summer.

Native of the Mediterranean region, escaped from gardens,

2. *A. MARITIMUM* L. Petals white, twice longer than the sepals.—2f. Oakland. Summer.

Native of the Mediterranean region, escaped from gardens.

Family 2. FUMARIACEÆ.

Sepals 1 pair. Corolla irregular. Stamens 6 more or less connate, diadelphous.

1. *Dicentra* Bork. BLEEDING-HEART.

Sepals small, two lateral petals flat, anterior and posterior petals calcarate. Ovary 1-celled, with two parietal placentæ; ∞ -ovulate. Fruit a 2-valved capsule.—2f.

1. *D. CHRYSANTHA* Hook. & Arn. Flowers yellow, paniced on a leafy stem.—Crystal Springs. Summer.

2. *D. FORMOSA* DC. Flowers rose-color, racemose on a scape.—Colma. Taylorville. Summer.

Family 3. PAPAVERACEÆ.

Sepals 2 or 3, caducous. Corolla regular; petals two pairs, sometimes three. Stamens not connate. Ovary free.

1. *Platystemon* Benth. CREAM CUPS.

Sepals 3. Petals 6. Stamens ∞ ; filaments flattened their whole length. Carpidia several, ∞ -seeded, at first syncarpous; stigmas free, sessile, linear. Fruit at last apocarpous, dis-

solving into several follicles. Plant villous. Flowers cream color.—⊙.

1. *P. CALIFORNICUS* Benth. Common. Spring.

2. *Platystigma* Benth. CREAM CUPS.

Sepals 3. Petals 6. Stamens ∞ ; filaments flattened only at the base. Ovary 3-angular, 1-celled; placentæ 3; stigmas 3. Fruit a capsule, 3-angular, 3-valved, ∞ -seeded.—⊙. External petals yellow, internal cream color.

1. *P. LINEARE* Benth. Filaments flat.—Common. Spring.

2. *P. CALIFORNICUM* Benth. & Hook. Filaments terete.—San Mateo. Spring.

3. *Dendromecon* Benth.

Sepals 2. Petals 4. Stamens ∞ . Ovary terete, linear with 2 placentæ; ovules ∞ ; stigma sessile, 2-lobed. Fruit a capsule 1-celled, 2-valved; seeds on the margin of the valves. Flowers yellow. Leaves rigid, entire.— $\frac{1}{2}$.

1. *D. RIGIDUS* Benth.—Tamalpais. Wright's Station. Summer.

4. *Eschscholtzia* Cham. CALIFORNIA POPPY.

Calyx and corolla inserted into a turbinate thalamus. Calyx gamosepalus, circumscissile at base. Petals 4. Stamens ∞ , adhering to the base of petals; filaments very short; anthers extrorse. Ovary free, terete, 1-celled, with 2 placentæ; stigmas 4. Fruit a capsule,

10-nerved, 2-valved; seeds on the margins of the valves.

1. *E. CALIFORNICA* Cham. Flowers orange— \mathcal{f} . Common. Summer.

5. *Meconopsis* Viguiet.

Sepals 2, caducous. Petals 4. Stamens ∞ ; anthers laterally dehiscent. Ovary obovate, 1-celled, with more than 2 placentæ; style short, persistent; stigmas radiating. Fruit a 1-celled capsule, dehiscent by several slits near the apex.

1. *M. HETEROPHYLLA* Benth. Petals scarlet. \odot . Sunnyside. San Mateo. Niles. Summer.

6. *Argemone* TOURN. PRICKLY POPPY.

Sepals 2 or 3. Petals 4 or 6. Stamens ∞ ; anthers extrorse. Ovary ovate, 2-celled, with more than 2 placentæ; stigmas nearly sessile, radiating. Fruit a capsule, dehiscent by several valves near the apex.— \odot .

1. *A. HISPIDA* Gray. Petals white.—Niles. Summer.

ORDER 2. POLYCARPICÆ. Floral parts with a tendency to arrange themselves spirally, and gradually merge into each other.

Family 1. LAURINEÆ.

Perigonium 6-parted. Lobes 2-seriate, imbricate in æstivation. Disc adnate to the base

of the perigonium; Stamens inserted on the margin of a disc, 6 or its multiple; anthers dehiscent by valves. Ovary 1-celled, compound of 3 carpidia, but only 1 ovule developed; style 1. Fruit baccate or drupaceous. Leaves alternate; stipules 0.

1. *Oreodaphne* Nees (*Umbellularia* Nutt).

Margin of perigonium deciduous. Stamens 9-12, in 3 rows; anthers of the two external rows introrse, of the third extrorse, their filaments 2-glandular at the base. Fruit a drupe resting on an enlarged thalamus.—½.

1. *O. CALIFORNICA* Nees. Common. Summer.

Bay tree. California laurel.

Family 2. **BERBERIDEÆ.**

Sepals, petals and stamens 1-3-seriate, opposite each other. Anthers extrorse, dehiscent by valves. Ovary 1-celled, with ventral placenta ∞-ovulate. Fruit capsular, follicular or baccate, 1 to several-seeded. Leaves alternate; stipules 0.

1. *Vancouveria* Decaisne.

Sepals 6. Petals 6. Stamens 6. Fruit a follicle, several-seeded. Flowers white.—2f.

1. *V. HEXANDRA* Morr. & Decaisne.—Marin County. Spring.

2. *Berberis* L. BARBERRY.

Sepals 6, protected by bracts. Petals 6. Stamens 6. Fruit baccate, few-seeded. Flowers yellow.— $\frac{1}{2}$.

1. *B. PINNATA* Lag. Leaflets pinnately veined, shining above, acuminate; first pair of leaflets near the base of the petiole.—Colma. Mission Hills. Spring.

Berries glaucous, edible. Stem and root contain a yellow dye.

2. *B. AQUIFOLIUM* Pursh. Leaflets pinnately veined, shining above, acuminate; first pair of leaflets remote from the base of the petiole.—Crystal Springs. Spring.

3. *B. REPENS* Lindl. Leaflets pinnately veined, not shining above, not acuminate.—Marin County. Spring.

4. *B. NERVOSA* Pursh. Leaflets palmately nerved.—Marin County. Spring.

Family 3. RANUNCULACEÆ.

Petals alternate with sepals and imbricate in æstivation. Stamens ∞ ; anthers extrorse (lateral), longitudinally dehiscent. Ovaries several, rarely reduced to 1, apocarpous. Fruit akene, follicle or berry. Stipules 0.

1. *Pæonia* TOURN. PEONY.

Sepals 5, unequal, persistent. Petals equal 5 or 10. Stamens ∞ ; ovaries few, seated on a

fleshy disc, ∞ -ovulate; stigmas sessile. Fruit of few follicles, coriaceous, ∞ -seeded.

1. *P. BROWNII* Dougl. Petals dark red.— \mathcal{M} . Marin County. Summer.

2. *Actæa* L. BANEBERRY.

Sepals 4, equal, caducous. Petals 4. Ovary but 1, 1-celled with ventral placenta, ovules 2-seriate; stigma sessile. Fruit baccate, ∞ -seeded.— \mathcal{M} . Flowers white.

1. *A. SPICATA* L. Berry red.—Redwood Peak. Tamalpais. Nicasio. Spring.

Narcotic poison.

3. *Delphinium* Tourn. LARKSPUR.

Sepals 5, petaloid; upper one calcarate. Corolla irregular; upper petal 0; 2 lateral petals projecting by their appendices into the calyx-spur. Ovaries not more than 5, ∞ -ovulate. Fruit several follicles.

1. *D. NUDICAULE* Torr. & Gray. Leaves almost all radical, 3-5-lobed, segments broad, obtuse, mucronulate; flowers in lax racemes, scarlet.— \odot . Contra Costa Range. Marin County. Summer.

2. *D. CALIFORNICUM* Torr. & Gray. Stem smooth, foliose; leaves 3-5-cleft, divisions lobed; flowers in strict, dense racemes; pedicels and flowers velvety, pale, sometimes tinged with violet.— \mathcal{M} . Saucelito. Mission Dolores. Summer.

3. *D. DECORUM* Fisch. & Mey. Pubescent; lower leaves 5-lobed, sparingly toothed, upper leaves with narrow, linear divisions; flowers on long pedicels in an open raceme; follicles glabrous; flowers blue.—♀. Marin County. Summer.

4. *D. VARIEGATUM* Torr. & Gray. Pubescent; upper and lower leaves dissected, segments oblong or linear; flowers on long pedicels in an open raceme; follicles pubescent; flowers blue or white.—♀. Marin County. Crystal Springs. Summer.

5. *D. SIMPLEX* Dougl. Canescent; upper and lower leaves dissected, segments linear; flowers on short pedicels in dense racemes; follicles pubescent; flowers white, yellow, red, blue or dark purple.—♂. Marin County. Summer.

This genus needs investigation in regard to its toxic properties. One of the species is known to the sheep-herders as dangerous to their flocks. Perhaps all the perennial species may contain an acrid, narcotic substance.

4. *Aquilegia* Tourn. COLUMBINE.

Sepals 5, petaloid, equal. Petals 5, bilabiate, prolonged backward into a spur. Fruit 5 follicles.—♀.

1. *A. TRUNCATA* Fisch. & Mey. Flowers orange.—Common. Spring.

5. *Ranunculus* L. BUTTERCUP.

Sepals 5. Petals 5 or multiple, with a scale at the base. Ovaries ∞ , 1-ovulate. Fruit ∞ -akenes.

1. *R. HEBECARPUS* Hook. & Arn. Akenes hispid, roughened; floral parts of small size and reduced in their typical number.— \odot . Marin County. Niles. Spring.

Flaccid annual of the habit of *Bowlesia*.

2. *R. MURICATUS* L. Lower leaves rounded, 3-lobed, irregularly crenate; akenes tuberculate, spinose; floral parts typical and of ordinary proportions; petals yellow.— $\odot\odot$. Presidio. Summer.

3. *R. MAXIMUS* Greene. Leaves broad, alternately divided; radical leaves on very long petioles; leaflets laciniately lobed; akenes glabrous, flat, orbicular, thin-edged; styles long; petals more than 5, yellow.— \mathcal{L} . San Rafael. Corte Madera. Summer.

4. *R. CALIFORNICUS* Benth. Radical leaves pinnately ternate; leaflets laciniately cut; akenes glabrous, flat, firmly edged; styles short; petals more than 5; yellow.— \mathcal{L} . Common. Summer.

5. *R. BLOOMERI* Watson. Leaves 3-5-lobed, crenate; akenes lenticular; styles long, straight; petals yellow.— \mathcal{L} . Sunnyside. Summer.

6. *R. FLAMMULA* L. Leaves entire; akenes smooth; stamens more than 10; petals bright yellow.— \mathcal{L} . Marin County. Summer.

7. *R. PUSILLUS* Poir. Leaves entire; akenes granulate; stamens never more than 10; petals yellow.— \odot . Inundated places near Napa. Summer.

8. *R. LOBBII* Hiern. Aquatic; leaves floating, 3-lobed; submerged leaves 0; styles long, filiform, with small terminal stigma; petals white with yellow claws.— \mathcal{L} . Olema. Summer.

9. *R. AQUATILIS* L. Aquatic with submerged capillary, multifid leaves; styles subulate, not longer than the ovary, introrsely stigmatose; petals white with yellow claws.— \mathcal{L} . Presidio. Marin County. Summer.

All perennial species possess an acrid, poisonous principle, which disappears by drying, so that the plant, avoided by animals in its growing state, becomes harmless when made into hay.

6. *Myosurus* L. MOUSETAIL.

Sepals 5, calcarate. Petals 5, on filiform claws. Fruit ∞ -akenes, crowded on an elongated, spike-like receptacle.— \odot . Dwarf plants with radical leaves.

1. *M. MINIMUS* L.—Niles. Spring.

7. *Thalictrum* TOURN. MEADOW-RUE.

Involucre 0. Sepals 4 or 5, very caducous. Petals 0. Ovaries ∞ , 1-ovulate. Fruit ∞ -capitate akenes. Leaves ternately compound. \mathcal{L} .

1. *T. FENDLERI* Engelm. Flowers violet. Saucelito. San Mateo. Summer.

8. *Anemone* Hall.

Involucre 3-foliolate. Sepals petaloid, 5 or a multiple. Petals 0. Ovaries ∞ , 1-ovulate. Fruit ∞ , capitate akenes.— \mathcal{L} .

1. *A. GRAYI* Kellogg & Behr. Flowers white, tinged with blue or purple.—Tamalpais. Camp Taylor. Spring.

9. *Clematis* L. VIRGIN'S-BOWER.

Sepals 4, petaloid, not imbricate in aestivation. Petals 0. Ovaries 1-ovulate. Fruit ∞ akenes, caudate by the persistent style.

1. *C. LASIANTHA* Nutt. Tomentose; flowers solitary on long peduncles, pale.— \mathcal{L} . Niles. Summer.

2. *C. LIGUSTICIFOLIA* Nutt. Almost glabrous; Flowers paniculate, pale.— \mathcal{L} . San Mateo. Niles. Marin County. Summer.

ORDER 3. RANALES. Floral parts having a pronounced tendency to merge gradually into each other.

Family 1. NYMPHÆACEÆ.

Sepals and petals inserted into a fleshy, cup-shaped thalamus. Stamens ∞ , ∞ -seriate, inserted with the petals; filaments flat, petaloid, anthers introrse, adnate. Carpidia several, immersed in the thalamus, and presenting the appearance of a ∞ -celled ovary; ovules ∞ , inserted on the partitions; styles united and forming a radiate, peltate stigma, projecting beyond the thalamus and persistent.

1. Nuphar Smith. WATER-LILY.

Sepals more than 4, petaloid. Petals ∞ , short, densely crowded round the thalamus. Stigma stipitate. Fruit a baccate pseudocarp. \mathcal{Z} . Aquatic. Flowers yellow.

1. *N. POLYSEPALUM* Engelm. Sepals more than 7.—Near the Marine Hospital. Taylorville, Paper-mill Creek. Summer.

SERIES 3. PERIGYNÆ. Thalamus cup-shaped, bearing sepals, petals and stamens, and receiving the carpidia in shape of a receptacle, which in many instances surrounds the carpidia enough to form an inferior ovary.

Section 1. CALYCIFLORÆ. Petals present and distinct, 4 or 5. Stamens 4 or 5, or their multiples.

ORDER 1. MYRTALES. Calyx lobes valvate in æstivation. Petals as many, or a mul-

ple, rarely a divisor. *Carpidia* united into a single ovary.

Family 1. HALORRHAGIDÆ.

Calyx tube adnate to the ovary; limb 4-parted. Petals, if present, alternate with calyx lobes. Stamens not more than twice as many as petals; if of equal number opposite to them. Placentæ central, 1-ovulate. Styles as many as cells.

1. *Hippuris* L. MARE'S TAIL.

Flowers ♂, or sometimes polygamous, monandrous; filament subulate. Ovary 1-celled, 1-ovulate, style filiform. Fruit a caryopsis.—♀. Aquatic. Leaves verticillate; flowers minute, solitary in the axils.

1. *H. VULGARIS* L.—Marine Hospital (at present extinct). Summer.

2. *Myriophyllum* L. WATER-MILFOIL.

Monœcious. ♂ : petals caducous; stamens 8. ♀ : calyx limb and petals minute; stigma villos; ovary 4-celled, 4-ovulate.

1. *M. SPICATUM* L.—Marine Hospital. Camp Taylor. Summer.

Family 2. ONAGRACEÆ.

Calyx tube adnate to the ovary. Petals alternate with them, not valvate in æstivation. Stamens inserted with petals, as many or a

multiple; anthers introrse. Placentæ central, ∞ -ovulate. Style 1. Stipules 0.

1. *Jussiaea* L.

Calyx tube not prolonged beyond the ovary, but forming an epigynous disc; lobes 4-6, persistent. Petals 4-6; stamens twice as many. Cells of ovary as many as calyx-lobes; stigma capitate. Fruit a septicial capsule.

1. *J. REPENS* L. Flowers yellow; leaves alternate.— \mathcal{f} . Aquatic. Niles. Alvarado. Summer.

2. *Zauschneria* Presl.

Calyx-tube prolonged considerably above the ovary, funnel-shaped, colored, with 4-lobed, deciduous limb. Petals not exceeding the calyx-lobes, 2-cleft, erect. Stamens 8, included. Ovary 4-celled, 4-valved. Seeds with a hairy crown. Fruit a capsule, linear, imperfectly 4-celled, α -seeded. Flowers scarlet; upper leaves alternate.— \mathcal{f} .

1. *Z. CALIFORNICA* Presl. — Oakland Waterworks. Niles. Summer.

3. *Epilobium* L. WILLOW-HERB.

Calyx-tube prolonged but little beyond the ovary; limb 4-parted, soon deciduous. Petals 4, obovate, or obcordate, inserted into an annular disc on the summit of the calyx-tube. Stamens 8. Ovary 4-celled. Fruit a 4-angular,

4-celled, loculicidal capsule. Seeds with a crown of long hair.

1. *E. FRANCISCANUM* Barbey. Stem angular; leaves serrulate; stamens shorter than the style; petals purple.—♀. Fort Point. Summer.

2. *E. MINUTUM* Lindl. Stem terete; leaves almost entire; four of the stamens as long as the style; petals rose-color.—⊙. Tamalpais. Summer.

4. *Gayophytum* Juss.

Calyx tube not prolonged beyond the ovary, limb 4-parted, reflexed, deciduous. Petals 4. Stamens 8; anthers versatile, those opposite the petals smaller and usually sterile. Ovary 2-celled; style short; stigma capitate or clavate. Fruit a capsule, 2-celled, unequally 4-valved.—⊙. Slender herbs.

1. *G. DIFFUSUM* Torr. & Gray. Petals pale rose-color.—Saucelito. Spring.

5. *Œnothera* L. EVENING PRIMROSE.

Calyx tube more or less prolonged above the ovary; lobes reflexed. Petals 4. Stamens 8; anthers versatile. Ovary 4-celled; ovules ∞; style filiform. Fruit a 4-celled capsule, loculicidal; seeds ∞, smooth. Leaves alternate.

1. *Œ. BIENNIS* L. Stigma lobes linear; flowers in a leafy spike, yellow.—⊙ ⊙. San Mateo. Niles. Summer.

2. *Æ. CALIFORNICA* Watson. Stigma lobes linear; flowers axillary, white, at last pinkish. 2f. Niles. Summer.

3. *Æ. OVATA* Nutt. Stigma capitate; calyx tube filiform, above the ovary; petals yellow. 2f. Common. Spring.

Leaves and flowers formerly in use as a salad.

4. *Æ. CHEIRANTHIFOLIA* Hornem. Stigma capitate; calyx tube obconic; capsule 4-angular with acute angles; canescent; leaves thick, entire; petals yellow.—2f. Seashore. Summer.

5. *Æ. MICRANTHA* Hornem. Stigma capitate; calyx tube obconic; capsule 4-angular with acute angles, very much contorted; leaves denticulate; petals yellow.—⊙. Niles. Alvarado. Summer.

6. *Æ. DENTATA* Cav. Stigma capitate; calyx tube obconic; capsule elongated, linear, shortly beaked; petals yellow.—⊙. Common. Summer.

7. *Æ. STRIGULOSA* Torr. & Gray. Stigma capitate; calyx tube obconic; capsule elongated, linear, obtuse, petals yellow.—⊙. Common. Summer.

6. *Godetia* Spach.

Calyx-tube prolonged beyond the ovary, funnel-shaped, the lobes reflexed. Petals 4. Stamens 8, unequal; anthers basifixed. Capsule ovate to linear; seeds smooth, ∞.—⊙.

1. *G. PURPUREA* Watson. Flowers in a leafy, terminal cluster; capsule oblong; seeds in a double row in each cell; petals purple. Common. Summer.

2. *G. LEPIDA* Lindl. Flowers in short simple spike; capsule oblong, ovate; seeds in a simple row in each cell; petals rose-colored, with a darker spot near the apex.—Niles. Summer.

3. *G. QUADRIVULNERA* Spach. Capsule linear, sessile, villous, 2-costate; petals purple. Common. Summer.

4. *G. TENELLA* Watson. Capsule linear, sessile, puberulent, not costate; petals purple. Niles. Summer.

5. *G. AMÆNA* Lilja. Capsule linear, attenuate at each end, pedicillate; plant minutely puberulent; petals rose-color, tinged with purple.—Common. Summer.

6. *G. EPILOBIOIDES* Watson. Capsule linear, attenuate into a short pedicel, acuminate at the apex; plant tomentosely pubescent; petals rose-color.—Niles. Summer.

7. *Clarkia* Pursh.

Calyx-tube prolonged beyond the ovary, obconical, deciduous, the lobes reflexed. Petals 4, unguiculate. Stamens 8, alternately depauperate; anthers basifixed. Capsule linear; seeds ∞ , smooth.— \odot .

1. *C. ELEGANS* Dougl. Petals rose-color. Alameda. Summer.

8. *Eucharidium* Fisch. & Mey.

Calyx-tube linear, elongated beyond the ovary, its lobes reflexed, deciduous. Petals 4, unguiculate. Stamens 4, alternate with the petals; anthers basifixed. Seeds ∞ , smooth.

⊙.

1. *E. CONCINNUM* Fisch. & Mey. Flowers red.—Marin County. Summer.

9. *Boisduvalia* Spach.

Calyx-tube prolonged beyond the ovary; funnel-shaped, deciduous; the lobes erect. Petals 4, sessile, 2-lobed. Stamens 8, those opposite the petals shorter; anthers basifixed. Capsules 4-celled; seeds ∞ , smooth.—⊙. Flowers purple.

1. *B. DENSIFLORA* Watson. Canescent; capsules oblong; partitions of the dehiscent capsule separating from the valves.—Common. Summer.

2. *B. GLABELLA* Walpers. Capsules oblong, almost straight; partitions of the dehiscent capsule adhering to the valves.—Common. Summer.

3. *B. CLEISTOGAMA* Curran. Capsules oblong, 4-angular, curving outward from the stem; partitions of the dehiscent capsule adherent to the valves.—Suñol. Summer.

Family 3. LYTHRARIÆ.

Calyx tube not adnate to the ovary. Stamens inserted with the petals, and of definite number; anthers introrse. Ovary free, with central placentæ, compound of carpidia; ovules ∞ ; style 1.

1. Lythrum L. LOOSE-STRIPE.

Calyx tubular, 8-12-costate; limb 8-12-dentate; dentitions alternating in size. Petals 4-6, opposite the smaller dentitions of the calyx. Stamens 8-12. Ovary 2-celled, covered by the persistent calyx. Flowers purple.

1. *L. ALATUM* Pursh.— \mathcal{L} . Tamalpais. Summer.

ORDER 2. ROSIFLORÆ. Calyx lobes imbricate in æstivation, as many as the perigynous petals. Ovaries several, in different degrees of consolidation, in rare cases reduced to a single one, the eccentric position of which indicates the abortion of its fellows.

Family 1. AMYGDALACEÆ.

Calyx free, 5-cleft, deciduous. Petals 5, convolute in æstivation. Ovaries almost always reduced to one, 1-celled, 2-ovulate. Fruit a drupe, by abortion generally 1-seeded. Leaves alternate, stipulate; stipules caducous.

1. *Prunus* L. PLUM. CHERRY.

Drupe fleshy; endocarp not rugose.— $\frac{1}{2}$.

1. *EMARGINATA* Walpers. Flowers in a corymb; petals white.—Tamalpais. Spring.

Black, cherry-like fruit. Not poisonous, although not edible.

2. *P. DEMISSA* Walpers. Flowers in terminal racemes; petals white.—Mills Seminary. Marin County. Spring.

Dark red or purplish cherry. Edible. "Choke-cherry."

3. *P. ILICIFOLIA* Walpers. Flowers in leafless racemes from the axils of evergreen leaves; petals white. — Common. Spring. "Holly-leaf Cherry."

Red or purplish cherry, not of bad taste, but suspicious. The leaves at times prove poisonous to sheep and cattle, probably only when by their withering hydrocyanic acid is developed, the smell of it in the withering leaves becomes very perceptible.

2. *Nuttallia* Torr. & Gray.

Flowers polygamous. Ovaries 5. Fruit several 1-seeded drupes. Flowers white.— $\frac{1}{2}$.

1. *N. CERASIFORMIS* Torr. & Gray. Flowers white.—Common. Spring.

Drupes blue, of agreeable but slightly bitter taste, but suspicious.

Family 2. SPIRÆEÆ.

Ovaries verticillate, rarely reduced to 1. Fruit follicular.

1. Spiræa L.

Seeds small exalbuminous, with membranaceous testa.

1. *S. DISCOLOR* Pursh. Flowers reddish.— $\frac{1}{2}$. Common. Summer.

2. Neillia Don. NINE-BARK. BRIDAL-WREATH.

Seeds albuminous, with a shining crustaceous testa. Flowers white.— $\frac{1}{2}$.

1. *N. OPULIFOLIA* Benth. & Hook.—Common. Summer.

Family 3. DRYADACEÆ.

Ovaries collected around a convex receptacle, rarely reduced to 1, 1-ovulate. Fruit indehiscent, frequently drupaceous.

1. Rubus L. BLACKBERRY. RASPBERRY.

Calyx persistent, concave or flattened; limb 5-parted. Stamens ∞ . Ovaries on a convex receptacle. Fruit drupes ∞ , on a conical receptacle.

1. *R. NUTKANUS* Moc. Leaves palmately lobed; petals white.— $\frac{1}{2}$. Common. Summer. "Salmon-berry."

2. *R. SPECTABILIS* Pursh. Armed with stout prickles; leaves 3-foliolate, underneath slightly

pubescent; flowers almost solitary, red. — $\frac{1}{2}$. Colma. Saucelito. Summer.

Fruit looks like a raspberry, red or yellow, rather insipid.

3. *R. LEUCODERMIS* Dougl. Armed with stout prickles; leaves 3-5-foliolate, leaflets underneath white-tomentose; flowers in few-flowered corymbs, white.— $\frac{1}{2}$. Los Gatos Creek. Summer.

Fruit a raspberry, red, of agreeable flavor.

4. *R. URSINUS* Cham. & Schlecht. Armed with slender bristles or bristle-like prickles; leaves generally 3-foliolate, not tomentose underneath; flowers white.— $\frac{1}{2}$. Common. Summer. "Wild-blackberry."

2. *Fragaria* L. STRAWBERRY.

Calyx concave or flattened, persistent, 5-lobed, augmented by 5 bractlets. Petals 5. Stamens a multiple of 5. Ovaries ∞ , on a convex receptacle; styles lateral. Fruit ∞ akenes on an enlarged, fleshy receptacle.— $\frac{1}{2}$.

1. *F. CHILENSIS* Ehr. Leaves perfectly smooth and shining on their upper surface. Flowers white.—Presidio. Cliff House. Point Bonita. Summer.

2. *F. CALIFORNICA* Cham. & Schlecht. Leaves sparingly villous and not shining on their upper surface.—Common. Summer.

3. *Potentilla* L. FIVE-FINGER.

Calyx concave or flattened, 5-lobed, augmented by 5 bractlets. Petals 5. Stamens a multiple of 5. Ovaries ∞ , on a slightly conical receptacle. Styles lateral. Fruit ∞ akenes. Receptacle not enlarged.

1. *P. GLANDULOSA* Lindl. Flowers cymose; petals pale yellow, sometimes white.— \mathcal{Z} . Contra Costa range. Summer.

2. *P. ANSERINA* L. Flowers axillary, solitary, yellow.— \mathcal{Z} . Common on moist places. Summer.

4. *Horkelia* Cham. & Schlecht.

Calyx campanulate, limb 5-parted, augmented by 5 bractlets. Petals 5. Stamens a multiple of 5. Ovaries inserted on a conical receptacle; styles subterminal. Fruit akenes.— \mathcal{Z} .

1. *H. CALIFORNICA* Cham. & Schlecht. Glandular pubescent; calyx-tube deeply campanulate; petals white.—Common. Summer.

2. *H. KELLOGGII* Greene. Silky pubescent; calyx-tube spreading; cupuliform; petals white. Alameda. Summer.

Family 4. SANGUISORBACEÆ.

Calyx-tube contracted at its apex, including the ovaries. Petals often 0, Stamens and ovaries often reduced in number. Ovaries 1-ovulate.

1. *Adenostoma* Hook. & Arn.

CHEMISAL. GREASEWOOD.

Calyx funnel-shaped; tube 10-costate; limb 5-parted. Petals 5; stamens a multiple. Ovary 1, with truncate, pubescent apex. Fruit an akene, included in the persistent calyx-tube. Flowers white.— $\frac{1}{2}$.

1. *A. FASCICULATUM* Hook. & Arn.—Common. Summer.

2. *Alchemilla* Tourn. LADY'S MANTLE.

Calyx-tube urceolate, persistent; limb 4-parted, with 4 minute, deciduous bractlets. Petals 0. Stamens reduced, 1-4. Ovaries 1-4; styles basilar. Fruit 1-4 akenes.

1. *A. ARVENSIS* Scop.— \odot . Common. Spring.

3. *Acæna* Vabl.

Calyx-tube contracted at the throat, oblong, angular, the angles armed with glochidiate prickles; limb persistent. Petals 0. Stamens reduced 1-5. Ovaries 1 or 2; styles terminal; stigmas penicillate. Fruit an akene, enclosed in the persistent calyx.— $\frac{1}{4}$.

1. *A. TRIFIDA* Ruiz & Pavon.—Common. Summer.

4. *Cercocarpus* HBK. MOUNTAIN MAHOGANY.

Calyx-tube cylindrical, persistent; limb turbinate, 5-lobed, deciduous. Petals 0. Stamens multiple of 5. Ovary 1; style termi-

nal, long exerted. Fruit an akene, linear, terete, caudate by the elongated, plumose, twisted style.— $\frac{1}{2}$

1. *C. PARVIFOLIUS* Nutt.—San Leandro. Lake Chabot. Spring.

Family 5. ROSACEÆ.

Carpidia ∞ , 1-ovulate, indehiscent, included in the fleshy tube of the calyx. Stamens ∞ .

1. *Rosa* Tourn. ROSE.

Calyx-tube urceolate, constricted at the throat; limb 5-parted. Petals 5. Styles lateral, exerted. Fruit ∞ -akenes, included in the fleshy calyx-tube.— $\frac{1}{2}$.

1. *R. CALIFORNICA* Cham. & Schlecht. Foliage and inflorescence tomentose; fruit globose; calyx lobes persistent.—Common. Summer.

2. *R. GYMNOCARPA* Nutt. Glabrous; petioles and stipules glandular; fruit ovate; calyx lobes deciduous.—Tamalpais. Summer.

Family 6. CALYCANTHEÆ.

Calyx fleshy; tube obconical; limb divided into ∞ -seriate lobes. Petals 0. Stamens ∞ ; anthers extrorse. Ovaries ∞ , 1-ovulate inserted on the whole inner surface of the calyx-tube; styles ∞ , terminal, exerted. Fruit ∞ -akenes, enclosed in the fleshy calyx-tube. Leaves opposite. Stipules 0.

1. **Calycanthus** Lindl. SPICE-BUSH.

Inner lobes of the ∞ -seriate calyx-limb smaller than the external. Outer series of stamens fertile, inner sterile. Flowers dark purple.— $\frac{1}{2}$.

1. **C. OCCIDENTALIS** Hook. & Arn.—Sonoma. Napa. Summer. Leaves and branches fragrant when bruised.

Family 7. **POMACEÆ.**

Calyx-tube adnate to the ovary; limb 5-parted. Carpidia 2-5, 2-ovulate; styles 2-5. Fruit baccate.

1. **Photinia** Lindl.

(*Heteromeles* J. Røemer.) TOYON.

Carpidia 2, imperfectly united and only half immersed in the calyx-tube; styles 2. Calyx-limb growing fleshy in fruit and covering the upper half of the carpidia, which are 1-seeded by abortion. Flowers in corymbose panicles.
 $\frac{1}{2}$.

1. **P. ARBUTIFOLIA** Røemer. Flowers white. Common. Summer.

Berries coral red, of astringent taste, but used by the original settlers in the preparation of a refreshing drink.

2. **Amelanchier** Medik.

JUNE-BERRY. SERVICE-BERRY.

Ovules of carpidium separated by an incom-

plete septum, 1 ovule becoming abortive in the ripe fruit.— $\frac{1}{2}$. Flowers in racemes.

1. *A. ALNIFOLIA* Nutt. Flowers white. Common. Summer. Berries purplish, edible.

ORDER 3. LEGUMINOSÆ. Ovary 1, 1-celled. Fruit a legume. Leaves alternate, stipulate.

Family 1. PAPILIONACEÆ.

Corolla papilionaceous. Stamens 10.

1. Pickeringia Nutt.

Petals of the carina not connate. Stamens not diadelphous. Legume linear, compressed. Flowers purple; leaves palmately 3-foliolate, evergreen; stipules evanescent.— $\frac{1}{2}$.

1. *P. MONTANA* Nutt.—Tamalpais. Crystal Springs. Wright's Station. Summer.

2. Thermopsis R. Brown.

Petals of the carina partly connate. Stamens not diadelphous. Legume linear, compressed. Flowers yellow; leaves palmately 3-foliolate; stipules persistent, foliaceous.— $\frac{1}{2}$.

1. *T. CALIFORNICA* Watson.—San Rafael. Saucelito. Summer.

3. Lupinus L. LUPINE.

Stamens monadelphous; anthers alternately oblong and reniform. Leaves palmate; stipules adnate.

1. *L. ARBOREUS* Sims. Slightly silky pubescent, pubescence appressed; flowers on slender pedicels; bracts linear, equaling the calyx, deciduous; superior and inferior calyx lip nearly equal; ovules more than 6 in the legume; petioles short; flowers yellow.—☉ ☉. Common. Summer.

2. *L. CHAMISSONIS* Esch. Appressed silky pubescence; pubescence dense; flowers on slender pedicels; bracts lanceolate, shorter than the calyx, deciduous; superior and inferior calyx lip nearly equal; ovules more than 6; petioles short; flowers violet or pale.—½. Common. Summer.

3. *L. DOUGLASSII* Agardh. Almost tomentose; flowers on slender pedicels; bracts linear exceeding the calyx, deciduous; superior and inferior calyx lip nearly equal; ovules more than 6; petioles short; flowers purple.—¼. Common. Summer.

4. *L. POLYPHYLLUS* Lindl. Leaflets glabrous above; flowers on long pedicels; bracts lanceolate, not longer than the calyx; superior and inferior calyx lip nearly equal, entire; ovules more than 6; petioles considerably longer than the leaflets; flowers blue, purple or white.—¼. Common. Summer.

5. *L. RIVULARIS* Dougl. Leaflets glabrous above; flowers on long pedicels; bracts setaceous, exceeding the calyx; superior and inferior

calyx-lip nearly equal; ovules more than 6; petioles equalling the leaflets; flowers purple. 24. Tamalpais. Summer.

6. *L. LITTORALIS* Dougl. Leaflets glabrous above; flowers on short pedicels. Racemes short; bracts setaceous, exceeding the calyx; ovules more than 6; petioles longer than the leaflets; flowers violet marked with yellow.—24. Marin County near the seacoast. Summer.

7. *L. ALBICAULIS* Dougl. Bracts subulate, shorter than the calyx; deciduous; calyx-lips nearly of equal length, the upper narrowed and dentate; vexillum acute, its margins near the apex coherent; ovules about 6; petioles not longer than the leaflets; flowers pale blue. 24. Marin County. Summer.

8. *L. AFFINIS* Agard. Pubescent; racemes formed by regular whorls; bracts short; superior calyx-lip 2-cleft; legume linear; ovules about 6; petioles twice as long as the leaflets; flowers blue with a white spot on the vexillum, which afterwards turns red.—⊙. Common. Spring.

9. *L. NANUS* Dougl. Pubescent; racemes formed by regular whorls; bracts exceeding the calyx; superior calyx-lip 2-cleft; ovules 6-8; petioles much longer than the leaflets; flowers blue marked with white.—⊙. Berkeley. Spring.

10. *L. MICRANTHUS* Dougl. Villous; racemes formed by regular whorls; bracts shorter than the calyx; superior calyx-lip 2-lobed; lobes short 3-angular; lower lip almost entire; flowers blue, sometimes marked with white.—☉. Common. Spring.

11. *L. TRIFIDUS* Watson. Villous; racemes generally reduced to a single whorl; superior calyx-lip 2-lobed, inferior calyx-lip 3-cleft; flowers blue marked with white.—☉. San Francisco sand hills. Spring.

12. *L. TRUNCATUS* Nutt. Nearly glabrous; flowers of the raceme scattered, not in whorls; superior calyx-lip 2-cleft; leaflets truncate at the apex; legume 8-ovuled; petals dark purple.—☉. Contra Costa hills. Spring.

13. *L. MICROCARPUS* Sims. Villous; calyx densely villous, its lips dentate, the upper very short; legume ovate, 2-ovuled; flowers purple, rose-color or white.—☉. Common. Spring.

14. *L. DENSIFLORUS* Benth. Villous; calyx almost glabrous; legume ovate, 2-ovuled; flowers yellow or pale.—☉. Common. Spring.

4. *Trifolium* Tourn. CLOVER.

Diadelphous. Petals persistent and adnate to the staminal tube. Legume utriculate, irregularly dehiscent. Leaves palmately compound; stipules adnate.

1. *T. REPENS* L. All peduncles axillary, longer than the leaf. Petals white.—24. Common. Summer.

2. *T. MACRÆI* Hook. & Arn. Heads not involucrate; flowers sessile; petals dark purple. ☉. Marin County. Spring.

3. *T. CILIATUM* Nutt. Heads not involucrate; flowers shortly pedicillate; calyx teeth lanceolate, rigid, ciliate at the bottom; petals white or purplish.—☉. Common. Spring.

4. *T. GRACILENTUM* Torr. & Gray. Heads not involucrate; flowers shortly pedicillate; leaflets obovate; petal rose-color.—Common. Spring.

5. *T. BIFIDUM* Gray. Heads not involucrate; flowers shortly pedicillate; leaflets bifid at the apex; corolla rose-color.—☉. Common. Spring.

6. *T. INVOLUCRATUM* Willd. Heads involucrate; involucre herbaceous, deeply lobed, the lobes deeply dentate; corolla not becoming inflated; legume more than 4-ovulate; petals rose-color.—☉. Common. Spring.

7. *T. TRIDENTATUM* Lindl. Heads involucrate; involucre herbaceous, deeply lobed, the lobes deeply dentate; corolla not becoming inflated; legume 1-2-ovulate; calyx teeth rigid, abruptly narrowed into a spinulose apex; petals considerably exceeding the calyx, purple with paler or white tips.—☉. Common. Spring

8. *T. PAUCIFLORUM* Nutt. Heads involu-
crate; involucre herbaceous, deeply lobed, the
lobes deeply dentate; corolla not becoming in-
flated; legume 1-2-ovulate; petals scarcely ex-
ceeding the calyx, purple or rose-color.—☉.
Common. Spring.

9. *T. MICROCEPHALUM* Pursh. Heads invo-
lucrate; involucre membranaceous, its lobes
entire; corolla not becoming inflated; petals
pale.—☉. Common. Spring.

10. *T. MICRODON* Hook. & Arn. Heads in-
volucrate; involucre membranaceous, its lobes
3-dentate; corolla not becoming inflated; petals
white or pale.—☉. Common. Spring.

11. *T. BARBIGERUM* Torr. Heads involu-
crate; involucre broad, ∞ -dentate; corolla be-
coming inflated; petals purple.—☉. Com-
mon. Spring.

12. *T. FUCATUM* Lindl. Heads involucrate;
involucre deeply lobed; corolla becoming in-
flated, pale.—☉. Common. Spring.

13. *T. DEPAUPERATUM* Desv. Involucre al-
most 0, reduced to a mere disc; corolla becom-
ing inflated, white or purple.—☉. Common.
Spring.

5. *Melilotus* Tourn. MELILOT.

Diadelphous. Petals free from the column
of stamens and deciduous. Legume globose.
Flowers racemose. Leaves pinnately 3-foli-
olate; stipules adnate.

1. *M. PARVIFLORA* Desf. Flowers yellow.
⊙. Common the year round.

Native of the Mediterranean region, but at present perfectly naturalized.

6. *Medicago* TOURN. MEDICK.

Diadelphous. Petals free from the column of stamens and deciduous. Legume globose. Flowers racemose. Leaves pinnately 3-foliolate; stipules adnate.

1. *M. SATIVA* L. Legume without prickles, spirally rolled up, the spiral leaving an open space in the center; flowers purple. — ♀. Common all the year round. "Alfalfa."

Native of Europe, but escaped from cultivation in different countries.

2. *M. DENTICULATA* Willd. Legume spirally rolled up and armed with a double row of prickles; flowers yellow. — ⊙. Common all the year round.

Native of Europe, a fodder plant, but owing to the prickles that infest its legume, not desirable in wool-growing districts.

3. *M. LUPULINA* L. Legume without prickles, spirally rolled up, the spiral leaving no open space in the center; flowers yellow. — ⊙. Common all the year round.

Native of Europe, now naturalized.

7. *Hosackia* Dougl.

Diadelphous. Petals unguiculate. Legume cylindrical, linear, almost straight, dehiscent, sessile. Ovules more than 2. Flowers umbellate, or by depauperation solitary. Leaves imparipinnate, 2 to several-foliolate; stipules often minute.

1. *H. STIPULARIS* Benth. Stipules large, foliaceous; flowers purplish.—2f. Contra Costa range. Summer.

2. *H. GRACILIS* Benth. Stipules scarious; vexillum yellow; wings and carina purple.—2f. Colma. Talmalpais. Summer.

3. *H. STRIGOSA* Nutt. Stipules reduced to blackish glands; rachis of leaf flattened; vexillum attenuate below, carina obtuse; flowers yellow.—⊙. Common. Spring.

4. *H. PARVIFLORA* Benth. Stipules reduced to blackish glands; rachis of leaf flattened; vexillum cordate; carina acute; flowers pale.—⊙. Common. Spring.

5. *H. PURSHIANA* Benth. Stipules reduced to glands; leaves almost sessile; flowers peduncled, pale; vexillum pink.—⊙. Common. Summer.

6. *H. SUBPINNATA* Torr. & Gray. Stipules gland-like, leaves petioled; flowers almost sessile; legume 5-seeded; petals yellow.—⊙. Common. Summer.

7. *H. BRACHYCARPA* Benth. Stipules gland-like; leaves petioled; flowers almost sessile; legume less than 5-seeded; petals yellow.—☉. Common. Summer.

8. *Syrmatium* Vogel. (*Hosackia*, Dougl.)

Diadelphous. Petals unguiculate. Legume indehiscent, incurved. Ovules less than 3. Umbels few-flowered. Stipules reduced to dark colored glands.

1. *S. GLABRUM* Torr. Glabrous; calyx teeth narrow, erect; flowers yellow.—2♂. Common. Summer.

2. *S. CYTISOIDES* Benth. Glabrous. Calyx teeth attenuate, recurved; flowers yellow.—2♂. Half Moon Bay. Crystal Springs. Summer.

3. *S. TOMENTOSUM* Hook. & Arn. Pubescent throughout; calyx teeth filiform; flowers yellow.—2♂. Marine Hospital. Summer.

9. *Psoralea* L.

Diadelphous. Glandular. Wings united to the carina. Ovary sessile, 1-ovulate. Legume included in the calyx, indehiscent. Leaves imparipinnate, 3-5-foliolate; stipules not adnate.

1. *P. ORBICULARIS* Lindl. Stems prostrate; corolla purple, sometimes white.—2♂. Common. Summer.

2. *P. STROBILINA* Hook. & Arn. Stem erect; stipules large, membranaceous, acuminate; peduncles shorter than the leaves; flowers in short spikes; bracts large, deciduous; corolla purple.— $\frac{1}{2}$. Contra Costa range. Summer.

3. *P. MACROSTACHYA* DC. Stem erect; stipules small, lanceolate; peduncles much exceeding the leaves; spikes cylindrical; bracts large, acuminate; corolla purple.— $\frac{1}{2}$. Marin County. Summer.

4. *P. PHYSODES* Dougl. Stems erect; stipules small, lanceolate; peduncles about equaling the leaves; flowers in racemes; bracts small; corolla white or purplish.— $\frac{1}{2}$. Coast Range. Summer.

10. *Amorpha* L. FALSE INDIGO.

Monadelphous. Vexillum unguiculate. Wings and carina 0. Ovary sessile, 2-ovulate. Legume very late, dehiscent. Glandular. Leaves imparipinnate.— $\frac{1}{2}$.

1. *A. CALIFORNICA* Nutt. Flowers purple. Marine County. Summer.

11. *Glycyrrhiza* L. LIQUORICE.

Diadelphous. Calyx not bracteolate. Vexillum, wings and carina straight. Anthers confluent. Legume ovate, compressed, few-seeded, echinate. Glandular. Leaves imparipinnate.— $\frac{1}{2}$.

1. *G. LEPIDOTA* Nutt. Flowers ochroleucous.—Niles. Summer,

12. *Astragalus* L.

RATTLE-WEED. LOCO-WEED.

Diadelphous. Carina blunt. Legume more or less divided by the intrusion of the dorsal suture. Seeds reniform. Leaves imparipinnate.

1. *A. DIDYMOCARPUS* Hook. & Arn. Legume transversely wrinkled, 2-seeded; corolla white and violet.—Contra Costa. Summer.

2. *A. TENER* Gray. Legume ∞ -seeded, not inflated; corolla white, sometimes violet tipped. \odot .—Golden Gate Park. Cemetery. Summer.

3. *A. LEUCOPHYLLUS* Torr. & Gray. Legume inflated, stipitate in the calyx; corolla pale. \mathcal{L} .—Suñol. Summer.

4. *A. CROTALARIÆ* Gray. Legume inflated, sessile in the calyx; stipules distinct; corolla white, more than twice the length of the calyx. \mathcal{L} .—Sandhills near San Francisco. Summer.

5. *A. MENZIESII* Gray. Legume inflated, sessile in the calyx; stipules connate, opposite the petiole; corolla white, more than twice the length of the calyx.— \mathcal{L} .

6. *A. DOUGLASII* Gray. Legume inflated, sessile in the calyx; corolla pale, hardly twice the length of the calyx.— \mathcal{L} . Coast Range. Summer.

7. *A. PYCNOSTACHYUS* Gray. Legume coriaceous, not inflated, lenticular, reticulated; ovules 5; seeds less than 4; corolla pale.—24. Bolinas Bay. Summer.

Some species of *Astragalus* have the reputation of being poisonous to cattle, sheep and horses. The specimens sent to the Academy of Sciences do not belong to any species found in the territory of the local flora. The whole matter is still wrapt in a mystery, like the one connected with the properties of *Gastrolobium*, a leguminous plant in Australia.

13. *Vicia* L. VETCH.

Diadelphous. Style filiform; apex pilose. Leaves paripinnate, terminating in branched tendrils.

1. *V. GIGANTEA* Hook. Flowers in a raceme; leaflets more than 9 pairs; corolla pale-purple.—24. Coast Range. Summer.

2. *V. AMERICANA* Muhl. Flowers in racemes; leaflets less than 9 pairs; corolla purple.—24. Common. Summer.

3. *V. EXIGUA* Nutt. Flowers pedunculate, solitary, rarely 2; leaflets less than 5 pairs; corolla purple.—○. Contra Costa. Spring.

4. *V. SATIVA* L. Flowers almost sessile; leaflets more than 4 pairs; corolla violet.—○. Fields, hedges. Spring.



(265)

Escaped from cultivation, originally European.

14. Lathyrus L. WILD PEA.

Style ventrally flattened toward the apex, concave, pilose along the inner side. Leaves paripinnate, ending in branched tendrils.— $\frac{1}{2}$.

1. *L. VESTITUS* Nutt. Leaves cirrhate; peduncles ∞ -flowered; petals rose-color, changing before withering into violet.— $\frac{1}{2}$. Common. Summer.

2. *L. PALUSTRIS* L. Leaves cirrhate; peduncles 2-6-flowered; petals purple.—Marin County. Summer.

2. *L. LITTORALIS* Endl. Leaves not cirrhate; leaflets 1-3 pairs and a small terminal one; vexillum purple; wings and carina pale. Marin County. Summer.

15. Cercis L.

RED-BUD. JUDAS-TREE.

Stamens not diadelphous, anterior longer than the posterior ones; anthers versatile. Petals unguiculate, those forming the carina separate, and larger than the rest. Legume flat, stipitate, ∞ -seeded; ventral suture winged.— $\frac{1}{2}$. Flowers purple in axillary fascicles. Leaves simple, appearing after the flowers.

1. *C. OCCIDENTALIS* Torr. Petals rose-color. Suñol. Spring.

ORDER 4. DAPHNALES. Petals 0. Perigonium and stamens inserted in a perigynous disc. Stamens 2 or its multiple. Carpidium 1, ovules never basilar. Stipules 0.

Family 1. THYMELACEÆ.

Perigonium gamosepalous. Ovary 1-celled, 1-ovulate; ovule lateral, attached near the apex; style lateral.

1. *Dirca* L. LEATHERWOOD.

Flowers ♂. Perigonium corolline, campanulate; limb obliquely truncate. Stamens 8, the alternate ones shorter. Hypogynous scales 0. Style filiform, subterminal. Fruit a drupe.

1. *D. OCCIDENTALIS* Gray. Perigonium greenish-yellow; drupe orange.—Contra Costa hills. Coast Range. Spring.

Bark acrid, blistering when applied to the skin.

Family 2. LORANTHACEÆ.

Evergreen, dichotomous, parasitic perennials. Branches articulate. Leaves opposite; stipules 0. Flowers epigynous. Stamens inserted on the perigonium, opposite its lobes and of the same number. Ovary 1-celled, 1-ovuled. Fruit baccate. Seed often containing several embryos.

[Affinities doubtful, perhaps connecting the Gymnosperm families of *Taxineæ* and *Gnetaceæ* to the Angiosperm families mentioned here and the *Proteaceæ*.]

1. *Phoradendron* Nutt. MISTLETOE.

Flowers diclinous; ♂ and ♀, on separate spikes. Anthers 2-celled. Stigma sessile. Fruit crowned by the persistent perigonium.

1. *P. FLAVESCENS* Nutt.—½. Ross Station. Spring.

Parasite on oaks, poplars, etc.

2. *Arceuthobium* Bieb. PINE MISTLETOE.

Diœcious. ♂ flowers sessile. Lobes of the perigonium ovate, spreading. Anthers sessile, inserted on the middle of the perigonium lobes, 1-celled, transversely dehiscent. ♀ flowers shortly pedicillate, compressed, with 2-dentate limb. Stigma sessile. Berry transversely dehiscent at the base.

1. *A. OCCIDENTALE* Engelm.—½. Contra Costa mountains. Autumn.

Parasite on conifers.

Series 4. MONOCHLAMYDEÆ.

Flowers incomplete. Number of carpidia corresponding to number of floral parts.

ORDER 1. SERPENTARIÆ. Leafy plants containing chlorophyll, as distinguished from the other order of the series, the *Rhizanthææ*,

which are parasitic plants without chlorophyll.

Family 1. ARISTOLOCHIÆ.

Perigonium gamosepalous. Stamens 6, or a multiple. Ovary inferior, 6-celled.

1. Asarum Tourn. WILD GINGER.

Stamens 12, free. Perigonium entirely epigynous; limb regular. Rhizome dichotomous.—2f.

1. A. CAUDATUM Lindl. Flower dark-purple.—Strawberry Valley. Tamalpais. Spring.

2. Aristolochia Tourn. DUTCHMAN'S PIPE.

Stamens 6, gynandrous. Limb of perigonium oblique, irregularly cleft.

1. A. CALIFORNICA Torr. Flowers greenish; twining creeper.—2f. Saucelito. Ocean lake. Ross Station. Spring.

SERIES 5. JULIFLORÆ. Inflorescence dense; flowers apetalous. Leaves never compound.

ORDER 1. Flowers diclinous; ♂ flowers amentaceous. Fruit 1-seeded, indehiscent. Stipules deciduous.

Family 1. CUPULIFERÆ.

Epigynous. Style 1; stigmas several. Fruit a nut (akene), the base of which is surrounded by a cupula or a persistent involucre.

1. *Quercus* L. OAK.

Monœcious. Aments slender. ♂ perigonium 4-8-parted; ♀, 6-dentate; style short; stigmas 3. Ovary 3-celled, 6-ovulate, surrounded by a scaly, budlike involucre, which in fruit enlarges into a cup.— $\frac{1}{2}$.

1. 2. *LOBATA* Née. ♂ : aments pendulous; abortive ovules at the base of the developed seed; acorns maturing the first year; leaves deciduous; branchlets glabrous; nut conical. Common. Spring.

“White oak.” “Roble.” Large, stately tree.

2. *Q. DOUGLASHII* Hook. & Arn. ♂ : aments pendulous; abortive ovules at the base of the developed seed; acorns maturing the first year; leaves deciduous; branchlets pubescent; nut oblong.—Niles. Sonoma.

“Blue oak.”

3. *Q. DUMOSA* Nutt. ♂ : aments pendulous; abortive ovules at the base of the developed seed; acorns maturing the first year; leaves persistent, coriaceous; nut oval.—Tamalpais.

4. *Q. CHRYSOLEPIS* Liebm. ♂ : aments pendulous; abortive ovules scattered over the surface of the developed seed; acorns biennial; leaves persistent, coriaceous; nut obtuse.—Marin County. Spring.

Tall tree, frequently called live oak.

5. *Q. AGRIFOLIA* Née. ♂ : aments pendulous; abortive ovules apical; acorns maturing the first year; leaves persistent, coriaceous, not reticulate; nut acute.—Common. Spring.

Tall tree, the genuine “live oak” of the Californians.

6. *Q. WISLIZENI* DC. ♂ : aments pendulous; abortive ovules apical; acorns biennial; leaves persistent, coriaceous, reticulate, entire to sinuately lobed; petioles short; nut acute. Niles. Spring.

Stately tree, frequently mistaken for “live oak.”

7. *Q. KELLOGGII* Newb. ♂ : aments pendulous; abortive ovules apical; acorns biennial; leaves deciduous, pinnatifid-lobed; petioles long; nut obtuse.—Marin County. Spring. “Black Oak.”

8. *Q. DENSIFLORA* Hook. & Arn. Aments erect, androgynous.—Marin County. Santa Cruz mountains. Summer. “Chestnut.” “Tan-ner’s bark.”

2. *Castanopsis* Spach. CHINQUAPIN.

Monœcious. Aments slender, panicled on the young shoots. ♂ perigonium 5-lobed; stamens 10. ♀ flowers 1-3, in a scaly involucre, sessile at the base of the ament; perigonium 6-lobed, 2-seriate; fruit 1-3 nuts en-

tirely closed in a prickly, irregularly rupturing involucre.— $\frac{1}{2}$.

1. *C. CHRYSOPHYLLA* A. DC. — Tamalpais. Santa Cruz mountains. Summer.

In Northern California a tall tree, in our local flora, a shrub.

3. *Corylus* Tourn. HAZEL.

Monœcious. Aments slender. Scales (bracts) imbricated. δ flowers besides the scales 2-bracteolate. Stamens 8; anthers 1-celled. ♀ flowers immersed into a bud; uppermost scales containing flowers, the rest empty. Ovary 2-bracteolate, crowned by a minute perigonium, 2-celled; stigmas 2. Fruit a nut, surrounded by a large incised involucre, formed by the growth of the two bractlets.— $\frac{1}{2}$.

1. *C. ROSTRATA* Ait.—Common. Spring.

Family 2. **BETULACEÆ.**

Hypogynous. δ ament: bracts shield-shaped, each enclosing 3 flowers with minute perigonium and four stamens. ♀ aments: bracts 3-lobed, each enclosing 3 naked flowers; ovary 2-celled; cells 1-ovulate; styles 0; stigmas 2. Fruit an akene (nutlet) affixed to the lignescent bract, and collected into a strobilaceous inflorescence.

1. *Alnus* Tourn. ALDER.

Fruit lignescent; nutlets attached to lignescent bracts on a lignescent axis.— $\frac{1}{2}$.

1. *A. RUBRA* Bongard. Nutlets winged. Contra Costa Mountains. Spring.

2. *A. RHOMBIFOLIA* Nutt. Nutlet not winged, but having a thickened margin.—Niles. Santa Cruz mountains. Spring.

Family 3. MYRICACEÆ.

Flowers diclinous, amentaceous, each bract containing a single, naked, sessile flower. ♂ aments filiform; ♀ ovate. Ovary connate with some hypogynous scales, 1-celled, 1-ovulate; style very short; stigmas 2. Fruit a nutlet; drupaceous by the incrassate, hypogynous bracts.

1. *Myrica* L. WAX-MYRTLE.

Only genus.—♂.

1. *M. CALIFORNICA* Cham. — Common. Spring.

ORDER 2. URTICALES. Flowers diclinous. Stamens opposite to the lobes of the perigonium. Ovary superior. Fruit 1-seeded. Leaves stipulate.

Family 1. PLATANEÆ.

Monœcious. Flowers naked, capitulate on globose receptacles. ♂ head: stamens ∞ , irregularly mingled with scales. ♀ head: ovaries ∞ , placed irregularly among scales; fruit a coriaceous nutlet. Leaves alternate, palmately lobed.

1. *Platanus* L. SYCAMORE.

Only genus.— $\frac{1}{2}$.

1. *P. RACEMOSA* Nutt.—Common. Spring.

Family 2. **URTICACEÆ.**

♂ perigonium 4-lobed; stamens 4; filaments inflexed in æstivation; anthers introrse, 2-celled; ovary rudimentary. ♀ perigonium 2 to 4-lobed; ovary 1-celled, 1-ovulate; fruit an akene, sometimes baccate by the persistent and succulent perigonium. Stipules generally persistent.

1. *Urtica* Tourn. NETTLE.

♂ perigonium regular and spreading. ♀ perigonium 2-parted, with sessile, penicillate stigma. Herbs covered with stinging hairs.

1. *U. HOLOSERICEA* Nutt. Inflorescence unisexual; petioles short, stout.— $2\frac{1}{2}$. Common. Summer.

2. *U. LYALLII* Watson. Inflorescence unisexual; petioles long, slender.— $2\frac{1}{2}$. Marin County. Summer.

3. *U. URENS* L. Inflorescence androgynous. \odot . Cultivated lands. Spring. Introduced from Europe.

2. *Hesperocnide* Torr.

♀ perigonium gamophyllous, urceolate, compressed, minutely dentate.— \odot .

1. *H. TENELLA* Torr.—San Francisco. Saucelito. Spring.
Small stinging herb.

Family 3. POLYGONACEÆ.

Perigonium imbricate in æstivation. Stamens definite, inserted on the base of the perigonium. Ovary superior, originally compound of 2 or 3 carpidia; by abortion 1-celled, 1-ovulate; but showing the 2 or 3 styles of the original carpidia. Ovule basilar. Fruit indehiscent. Stems articulate. Leaves alternate, sheathing.

[Affinities of this family not yet settled. The articulate stem approaches it to the *Piperaceæ*; the basilar ovule to the *Chenopodiaceæ*.]

1. *Rumex* L. DOCK. SHEEP-SORREL.

Perigonium 6-sepalous, 2-seriate; 3 inner segments larger and more petaloid. Stamens 6, in pairs opposite the outer sepals. Styles 3 with penicillate stigmas. Fruit a 3-cornered akene inclosed in the 3 inner sepals, forming a spurious capsule. Flowers racemose.

1. *R. SALICIFOLIUS* Weinm. Calyx valves entire or denticulate, bearing large callosities; leaves linear to lanceolate, not undulate, attenuate into a short petiole.— γ . Common. Spring.

2. *R. CRISPUS* L. Calyx-valves entire, bearing callosities; leaves lanceolate, undulate, truncate at base.— γ . Common. Spring.

“Yellow dock.” Native of Europe, now naturalized.

3. *R. CONGLOMERATUS* Murray. Calyx valves entire, bearing callosities; leaves lanceolate, the lower leaves cordate, slightly undulate.

♀. Common. Spring.

Also called “yellow dock,” native of Europe, and at present naturalized.

4. *R. OBTUSIFOLIUS* L. Calyx valves with slender awn-like teeth and bearing callosities.

♀. Contra Costa. Spring.

7. *R. ACETOSELLA* L. Dioecious. Valves without callosities; leaves hastate.—♀. Common. Spring. “Sheep-sorrel.”

2. *Polygonum* L. KNOT-GRASS.

Perigonium 5, sometimes 4-phyllous. Stamens 3-8, single or in pairs opposite to the sepals. Styles 3, sometimes 2; stigmas capitate. Fruit a 3-cornered or lenticular akene, enclosed in the persistent perigonium.

1. *P. PARONYCHIA* Cham. & Schlecht. Flowers in leafy spikes; leaves lanceolate, with revolute margins; flowers pale, veined with green. ♀. San Francisco. Summer.

2. *P. AVICULARE* L. Flowers axillary, pale.—♂. Common. Spring, Summer.

3. *P. CALIFORNICUM* Meissn. Flowers spicate; bracts foliaceous, each bract containing

a single sessile flower; leaves linear; flowers rose-color.—⊙. Niles. Spring.

4. *P. NODOSUM* Pers. Flowers spicate; bracts scarious, each bract protecting a fascicle; stamens 6; styles 2, included; leaves cuneate at base; flowers pale.—⊙. Marine Hospital. Spring.

5. *P. MUHLENBERGII* Watson. Flowers spicate; bracts scarious, each bract protecting a fascicle; stamens 5; style 2-cleft, exserted; leaves cordate at base; flowers rose-color.—♀. Marine Hospital. Spring.

6. *P. CONVULVULUS* L. Stem twining; leaves hastate; flowers pale.—⊙. Cultivated grounds. Summer.

Ballast weed, introduced from Europe.

3. *Eriogonum* Michx.

Involucre ∞-flowered, campanulate, slightly angulate, 6-dentate. Flowers ♂ pedicillate, exserted, seated on a receptacle with scarious bracts. Perigonium 6-parted, 2-seriate. Stamens 9, in pairs opposite to the external lobes; singly to the internal ones. Styles 3; stigmas capitate. Fruit a 3-angled, rarely a lenticular, akene. Peduncles 2-3-chotomous.

1. *E. ANGULOSUM* Benth. Involucre nerveless, hemispherical; flowers pale.—⊙. Niles. Spring.

2. *E. LATIFOLIUM* Smith. Involucres 5-6-nerved, collected into a glomerule; peduncles stout, solid; flowers rose-color.—♂. Common. Summer.

3. *E. NUDUM* Dougl. Involucres 5-6-nerved, collected into a glomerule; peduncles slender, fistulose; flowers pale.—♂. Coast Range. Summer.

4. *E. TRUNCATUM* Torr. & Gray. Involucre 5 or 6-nerved, solitary, forming a cyme; leaves rosulate, mostly radical; flowers rose-colored. ⊙. Niles. Spring.

5. *E. VIRGATUM* Benth. Involucre 5 or 6-nerved, solitary, forming a dichotomous panicle; leaves lanceolate; bracts shorter than the involucre; flowers pale.—⊙. Coast Range. Spring.

6. *E. VIMINEUM* Dougl. Involucre 5 or 6-nerved, solitary, forming a dichotomous panicle; leaves orbicular; flowers pale.—⊙. Common. Spring.

7. *E. GRACILE* Benth. Involucre 5 or 6-nerved, solitary, forming a dichotomous panicle; leaves lanceolate; bracts longer than the involucre; flowers pale.—⊙. Santa Cruz mountains. Spring.

3. *Chorizanthe* R. Br.

Involucre 3-flowered, tubular, 3-angular, 6-dentate, mucronate or aristate. Flowers ♂,

scarcely exerted. Perigonium 6-parted, 2-seriate. Stamens typically 9, in pairs opposite to the external lobes; singly to the internal ones; occasionally by abortion 6 or 3. Styles 3; stigmas capitate. Fruit a 3-angular akene. Peduncles dichotomous.

1. *C. MEMBRANACEA* Benth. Stem erect; involucre equally 6-cleft, 3-flowered, only one flower developed; leaves scattered.—⊙. Niles. Spring.

2. *C. DOUGLASHI* Benth. Stem erect; involucre unequally cleft, 1-flowered; leaves verticillate.—⊙. Santa Cruz mountains. Spring.

3. *C. PUNGENS* Benth. Stem procumbent; involucre unequally dentate, teeth alternately smaller; leaves mostly opposite.—⊙. Presidio. Spring.

5 *Lastarriæa* Remy.

Involucre 0. Perigonium coriaceous, tubular, 6-dentate, the dentations uncinately aristate; stamens 3, inserted on the throat. Fruit a 3-angled akene.—⊙. Diffusely branched.

1. *L. CHILENSIS* Remy.—Common. Spring.

6. *Pterostegia* Fisch. & Mey.

Involucre a single 2-lobed bract, subtending a single ♀ flower. Perigonium 6-parted, persistent. Stamens 6 (rarely fewer). Fruit a 3-angular akene, loosely enveloped by the en-

larged involucre. Prostrate, diffusely dichotomous; leaves opposite.—⊙.

1. *P. DRYMARIOIDES* Fisch. & Mey. Common. Spring.

ORDER 3. PIPERALES. Flowers sessile, in dense spikes or racemes, bracteate, without perigonium. Stem articulate.

Family 1. SAURUREÆ.

Ovary central, compound of several follicular carpidia; stigmas several.

1. *Anemopsis* Nutt. YERBA MANSA.

Flowers in a dense, conical spadix, with a several-leaved, persistent, colored involucre; each flower subtended by a colored bract. Stamens adnate to the base of the ovary. Ovary immersed into a rachis, 1-celled, with parietal placenta. Fruit a capsule, dehiscent from the apex. Leaves mostly radical.—♂.

1. *A. CALIFORNICA* Hook. Bracts and involucre white.—San Pablo marshes. Summer. Herb and root-stock aromatic.

Family 2. CERATOPHYLLÆ.

Monœcious. Involucre 12-cleft; lobes linear, truncate. ♂ flowers: anthers ∞, sessile. ♀ flowers: ovary 1-celled, 1-ovulate; style terminal; fruit a nutlet, with persistent style and involucre. ♂. Submersed aquatics; stem ar-

ticulate; leaves verticillate, sessile, dichotomous.

Affinities doubtful. Further investigations even may remove them altogether from the *Angiosperms* and place them near the *Gnetaceæ*, together with the *Podostemoneæ*.

1. *Ceratophyllum* L.

Only genus of the family.

1. *C. DEMERSUM* L. Common aquatic.

DIVISION 2. MONOCOTYLEDONES.

SERIES 1. COROLLIFLORÆ.

Perigonium 2×3 parted. Ovary 3 carpida, syncarpous in different degrees.

ORDER 1. GYNANDRÆ.

Family 1. ORCHIDEÆ.

Placentæ parietal.

1. *Epipactis* Haller.

Anther 1, persistent, not connate with the column; pollinia 2, attached to the common gland. Perigonium spreading; labellum geniculate, inferior part concave. Ovary contorted only at base; column short, terete.— $\frac{2}{4}$ Cauliscent, flowers in a loose, few-flowered raceme.

1. *E. GIGANTEA* Dougl. Sepals brownish-green, labellum white, dotted.—San Francisco [extinct]. Camp Taylor. Summer.

2. *Spiranthes* Richard LADIES' TRESSES.

Anther 1, persistent, not connate with the column; pollinia 2, attached to the common gland. Perigonium oblique; labellum enclosed, canaliculate, embracing with its base the short column.—♂. Flowers forming a spiral spike.

1. *S. ROMANZOFFIANA* Cham. Callosities of the lip smooth, obscure; flowers greenish-white.—Presidio. Marin County. Autumn.

2. *S. PORRIFOLIA* Lindl. Callosities of the lip prominent, pointing downward; flowers greenish-white.—Marin County. Autumn.

3. *Habenaria* Willd.

Anther 1, persistent, entirely connate with the column; pollinia 2, divergent at the lower end. Perigonium ringent, galeate; lip elongate, spreading, calcarate. Column short.—♂. Caulescent.

1. *H. ELEGANS* Bolander. Sepals 1-nerved; flowers green. —Oakland hills. San Mateo. San Francisco [extinct]. Summer.

2. *H. LEUCOSTACHYS* Watson. Sepals 3-nerved; flowers white. —Saucelito. San Francisco [extinct]. Summer.

4. *Corallorhiza* Haller! CORAL-ROOT.

Anther 1, terminal, caducous, not adnate to the column; pollinia 4, globose. Perigonium

ringent. Lip adnate to the column, serrate at base, 3-lobed; lateral lobes very small. Column semi-terete.—2♂. Aphyllous and without chlorophyll; flowers in spikes.

1. *C. MULTIFLORA* Nutt. Flowers calcarate; sepals greenish; lip white, dotted with red.—Crystal Springs. San Mateo. San Francisco [extinct]. Summer.

2. *C. BIGELOVII* Watson. Spur 0, sepals purple.—Tamalpais. Summer.

5. *Cypripedium* L. LADY'S SLIPPER.

Anthers 2. Lip inflated.—2♂. Flowers few or solitary.

1. *C. MONTANUM* Dougl. Sepals brownish; lip pale, veined with purple.—Santa Cruz Mountains. Summer.

ORDER 2. EPIGYNÆ. Stamens free, 3 or 6. Ovary inferior.

Family 1. IRIDACEÆ.

Stamens 3, opposite to the external segments of the perigonium; anthers extrorse.

1. *Sisyrinchium* L. BLUE-EYED GRASS.

Divisions of the perigonium equal. Stamens monadelphous. Style short; stigmas 3, filiform, involute, alternate with the stamens. Fruit a capsule, obovate.

1. *S. BELLUM* Watson. Filaments united to the top; flowers blue with yellow center.— \mathcal{C} . Common. Summer.

2. *S. CALIFORNICUS* Ait. Filaments united only at the base; flowers yellow.— \mathcal{C} . Colma. Saucelito. Summer.

2. *Iris* L.

Perigonium tubular at the base; segments equal, but differing in shape; the 3 external reflexed; the 3 internal erect. Stigma 3-parted, petaloid.— \mathcal{C} .

1. *I. MACROSIPHON* Torr. Tube of the perigonium elongated, cylindrical; stem leafy; capsule ovoid, acute at each end; flowers variable in color, but most frequently lilac-purple. Marin County. Spring.

2. *I. DOUGLASIANA* Herbert. Tube of the perigonium elongated, cylindrical; stem leafy; capsule oblong, 3-gonal; flowers pale lilac, external sepals with white center, lined with purple.—Common. Spring.

3. *I. LONGIPETALA* Herbert. Tube of the perigonium short and funnel-shaped; stems almost naked; flowers lilac, external sepals white, veined with violet; midrib yellow. Common. Spring.

ORDER 3. CORONARIÆ. Perigonium 6-parted in different degrees. Stamens 3 or 6. Ovary superior.

Family 1. SMILACEÆ.

Anthers not extrorse. Ovary 3-celled, with central placentæ. Fruit a berry. Testa of the seed not crustaceous.

1. Smilacina Desf. FALSE SOLOMON'S SEAL.

Divisions of the perigonium equal. Stamens inserted at the base of the segments. Cells of the ovary 2-ovulate; cells of the ripe fruit 1-seeded.— γ . Leaves sessile and amplexicaul.

1. *S. AMPLEXICAULIS* Nutt. Flowers greenish-white, paniced; stamens exceeding the perigonium; berry red.—Common. Spring.

2. *S. SESSILIFOLIA* Nutt. Flowers in a simple raceme, white; stamens shorter than the perigonium; berry dark.—Common. Spring.

2. Maianthemum Mœnch.

Perigonium 4-parted. Stamens 4.— γ . Leaves petiolate.

1. *M. BIFOLIUM* DC. Flowers white; berry red.—Saucelito. Spring.

3. Prosartes Don.

Perigonium 6-sepalous, campanulate; segments saccate at base. Stamens 6, attached at the base of the sepals, and deciduous with them. Anthers adnate, dehiscing laterally. Ovary 3-celled; cells 2-ovulate; style decidu-

ous. Cells of the berry 1-2-seeded. — 2♂. Leaves alternate, sessile, amplexicaul.

1. *P. MENZIESII* Don. Perigonium gibbous at base, its sepals almost erect; flowers green; berry red.—Lagunitas Creek. Spring.

2. *P. HOOKERI* Torr. Perigonium narrow at base, its sepals spreading above; flowers green; berry red.—Common. Spring.

3. *Trillium* L. WAKE-ROBIN.

Perigonium 6-sepalous; 3 external sepals herbaceous, persistent; 3 internal petaloid, marcescent. Stamens 6; anthers linear, lateral; connective broad. Ovary 3-celled, ∞-ovulate; styles 3. Berry ∞-seeded.—2♂. Leaves 3, verticillate; nervation reticulate.

1. *T. SESSILE* L. Flowers sessile; corolla variable; berry red.—Common. Spring.

Rhizome, and probably also berry, an acrid poison.

2. *T. OVATUM* Pursh. Flowers pedunculate, white, gradually turning red.—Tamalpais. Camp Taylor. Spring. Poisonous.

Family 2. *LILIACEÆ*.

Anthers not extrorse. Ovary 3-celled. Testa of the seed crustaceous.

1. *Clintonia* Raf.

Perigonium 6-sepalous, campanulate, deciduous. Stamens 6, inserted on the base of the

sepals; anthers versatile. Style slender, deciduous. Fruit a berry.— \mathcal{L} . Flowers on a scape.

1. *C. ANDREWSIANA* Torr. Flowers scarlet; berry blue.—Los Gatos. Lagunitas Creek. Summer.

2. *Chlorogalum* Kunth. SOAP-ROOT.

Perigonium 6-sepalous; sepals linear, persistent, marcescent. Stamens 6, adnate to the base of the sepals; anthers versatile. Cells of the ovary 2-ovulate; style filiform, deciduous. Fruit a membranaceous, 3-lobed, loculicidal capsule.— \mathcal{L} . Bulbous; inflorescence racemose, paniculate.

1. *C. POMERIDIANUM* Kunth. Flowers white. Common. Summer.

In former times used in washing linen. Old settlers consider it beneficial to the growth of the hair, like the bark of the Chilean Rosaceous tree “*Quillaja*.”

3. *Allium* L. ONION.

Perigonium persistent. Stamens 6, adnate to the base of the segments; filaments naked with dilated base; anthers versatile. Ovary 3-lobed; cells 2-ovulate; style filiform, persistent; stigma simple. Fruit a loculicidal capsule.— \mathcal{L} ; bulbous; scapigerous; inflorescence umbellate.

1. *A. UNIFOLIUM* Kellogg. Capsule not crested; leaves linear, flat; flowers rose-color. Tamalpais. Berkeley. Summer.

2. *A. ATTENUIFOLIUM* Kellogg. Capsule 6-crested; leaves convolute, filiform; perigonium at last thin and lax; flowers pale.—Colma-Marin County. Summer.

3. *A. SERRATUM* Watson. Capsule 6-crested, crests narrow, central; perigonium fleshy, rigid; flowers rose-color. — Colma. Lake Chabot. Summer.

4. *A. LACUNOSUM* Watson. Capsule bearing toward the summit 6 obtuse ridges. Perigonium fleshy, rigid; flowers rose-color.—Alma. Summer.

4. *Cyanotris* Raf.

(*Camassia* Lindl.) CAMASS.

Perigonium 6-sepalous, persistent; 5 superior sepals ascending; inferior one deflexed. Filaments filiform, ascending; anthers versatile. Ovary ovate, 3-celled; cells ∞ -ovulate; style filiform, declinate; stigma slightly 3-cleft. Fruit a loculicidal, 3-angulate capsule; cells ∞ -seeded.— γ . Bulbous; scapigerous; flowers in a simple raceme.

1. *C. ESCULENTA* Lindl. Flowers of different shades of blue.—Marin County. Summer.

The bulb has a taste like garlic, and is eaten by the Indians.

5. *Brodiaea* Smith.

Perigonium funnel-shaped, angulate, persistent. Stamens 3, alternating with as many staminodia. Hypogynous disc 3-lobed. Ovules ∞ . Style persistent; stigma 3-lobed. Capsule loculicidal.— \mathcal{Z} . Scapigerous; umbellate.

1. *B. GRANDIFLORA* Smith. Leaves about equaling the scape; cells of the ovary 6-8-seeded; flowers purple to rose-color.—Common. Summer.

2. *B. MINOR* Watson. Cells of the ovary 3-seeded; flowers purple.—Marin County. Summer.

3. *B. TERRESTRIS* Kellogg. Leaves considerably longer than the short scape; cells of the ovary 6-8-seeded; flowers purple.—Crystal Springs. Summer.

4. *B. CONGESTA* Smith. Flowers subcapitate; segments of perigonium twice longer than the tube; flowers purple.—Common. Summer.

6. *Triteleia* Hook.

Perigonium salver-shaped, persistent. Fertile stamens 6; filaments short, 3 inserted on the throat, and 3 half way down the tube. Ovary stipitate; ovules ∞ . Capsule loculicidal.— \mathcal{Z} . Scapigerous umbellate; capitate.

1. *T. CAPITATA* Benth. Stamens in 2 rows; flowers subcapitate; segments of perigonium

little longer than the tube; flower blue, varying from purple to white.—Common. Summer.

2. *T. LAXA* Watson. Stamens in 2 rows; flowers umbellate; tube of perigonium exceeding the lobes; flowers blue.—Common. Summer.

3. *T. PEDUNCULARIS* Watson. Stamens in 2 rows; flowers umbellate; lobes of perigonium exceeding the tube; flowers purple, sometimes white.—Tiburon. Summer.

4. *T. IXIOIDES* Watson. Stamens almost in a simple row, inner filaments exceeding the 3 outer ones; flowers yellow, marked with purple.—San Rafael. Summer.

5. *A. LACTEA* Watson. Stamens almost in a simple row; all six filaments alike; flowers pale with green nerve.—Common. Summer.

7. *Muilla* Watson.

Perigonium subrotate, persistent. Stamens 6, inserted near the base; anthers versatile. Ovary sessile, ∞ -ovulate; style persistent, clavate, at length splitting. Capsule globose, loculicidal.— ζ . Scapigerous; umbellate.

1. *M. MARITIMA* Watson. Sepals white with 2 green nerves.—Common. Summer.

8. *Lilium* L. LILY.

Perigonium 6-sepalous; sepals equal, spreading with a nectariferous groove near the base,

deciduous. Ovary ∞ -ovulate; style undivided; stigma 3-angular.— γ . Bulbous. Pedicels bractless.

1. *L. PARDALINUM* Kellogg, Flowers orange-color, with purple dots.—Marin County. Contra Costa. Summer.

9. *Fritillaria* L.

Perigonium deciduous; 6-sepalous, campanulate, with a smooth nectariferous pit near the base of each sepal. Ovary ∞ -ovulate; style 3-parted.— γ . Bulbous. Flowers bracteate.

1. *F. LILIACEA* Lindl. Leaves in a single whorl near the base of the stem; capsule stipitate, with obtuse angles; flowers greenish-white.—South San Francisco. Mission Hills. Lake Chabot. Summer.

2. *F. LANCEOLATA* Pursh. Leaves in 1 to 3 whorls on the upper part of the stem; flowers dark purple, mottled with yellow.—Common. Summer.

Bulb of this species probably poisonous, like that of its European congener *F. Meleagris*.

10. *Calochortus* Pursh. MARIPOSA.

Perigonium 6-sepalous, deciduous; 3 external sepals smaller and lanceolate; 3 internal, broad, with a conspicuous nectariferous gland near the base. Ovary ∞ -ovulate; style 0; stigmas 3, reflexed, persistent.— γ . Bulbous.

1. *C. ALBUS* Dougl. Flowers subglobose, nodding; glandular pit of sepals shallow; with 4 transverse scales; flowers white.—San Mateo. Marin County. Summer.

2. *PULCHELLUS* Dougl. Flowers subglobose, nodding; glandular pit of sepals deep and covered by stiff appressed hairs; flowers yellow. Napa. Summer.

3. *C. LILACINUS* Kellogg. Flowers pale-lilac; erect; pedicels slender, recurved in fruit. Oakland Reservoir. Lagunitas. Summer.

4. *C. LUTEUS* Dougl. Flowers and capsules erect on stout pedicels; glands of the sepals rounded and about as broad as the claw; color of inner sepals variable (yellow, white or lilac) marked with purple.—Common. Summer.

5. *C. VENUSTUS* Benth. Flowers and capsules erect on stout pedicels; gland of the inner sepals much narrower than the claw; color of inner sepals white or lilac, with a reddish spot near the top, and a purple spot bordered with yellow.—Contra Costa Range. Summer.

Family 3. MELANTHACEÆ.

Anthers extrorse. Ovary 3-celled; styles 3. Capsule septicidal. Testa of the seeds not crustaceous.

1. *Scoliopus* Torr.

Perigonium 6-sepalous, spreading, deciduous; 3 external sepals lanceolate; 3 internal linear. Stamens 3, inserted at the base of the external sepals; anthers 2-celled, attached above the base. Ovary 3-quetrous, 1-celled, with 3 parietal placentaë; ovules 2-seriate on each; styles 3, linear, persistent, recurved, canaliculate. Capsule irregularly dehiscent.— \mathcal{L} . Leaves 2. Peduncles long, 1-flowered, arranged in an umbel.

1. *S. BIGELOVII* Torr. Outer sepals dark purple, inner sepals pale, lined with purple. Saucelito. Tamalpais. Spring.

2. *Zygadenus* Michx. ZYGADENE.

Perigonium 6-sepalous; sepals unguiculate, 2-glandular at the base, persistent; stamens 6, inserted on the claw, anthers reniform. Ovary 3-celled, ∞ -ovulate, styles 3, divergent; capsule septicidal.— \mathcal{L} . Bulbous. Leaves lanceolate. Inflorescence racemose.

1. *Z. FREMONTII* Torr. Outer sepals unguiculate; flowers greenish-white.—Marin County. Crystal Springs. Spring.

Has the reputation of being poisonous.

2. *Z. VENENOSUS* Watson. All sepals unguiculate; flowers greenish-white.—Common. Spring.

Has the reputation of being poisonous, but is certainly not so to hogs, who devour its bulbs greedily and without any injury to themselves.

3. *Xerophyllum* Rich.

Perigonium 6-sepalous; sepals sessile, persistent; 3 internal ones smaller. Stamens inserted on the base of sepals; anthers laterally dehiscent. Ovary 3-celled; cells few-ovulate; styles 3, recurved, persistent. Capsule septicidal.—2♂. Rhizomatous. Leaves linear, elongate. Inflorescence racemose.

1. *X. TENAX* Nutt. Flowers white.—Tamalpais. Wright's Station. Spring.

This plant is remarkable by its flowering plentifully at certain years and then disappearing altogether for a long period. This circumstance is probably owing to the fact that the plant requires a fixed age before flowering and then dies after having ripened its seeds. We find the same peculiarity in many species of Bamboo, in some palms, viz.: *Corypha*, and in *Agave*, all of which plants are annuals in a botanical sense, but require a number of years to complete their biological cycle.

Family 4. JUNCACEÆ.

Perigonium glumaceous. Anthers basifixed.

1. *Juncus* L. BOG-RUSH.

Perigonium 6-sepalous; 3 external sepals carinate. Ovary 3-celled; cells ∞ -ovulate. Capsule 3-valved, with central placentæ, ∞ -seeded. Mostly aquatics.

1. *J. ROBUSTUS* Watson. Stem naked. Inflorescence lateral; flowers of the panicle branches clustered; leaves terete and pungent. 24. Santa Clara. Summer.

2. *J. LESEURII* Bolander. Stem naked, terete; inflorescence lateral; flowers of the compound panicle branches solitary; leaves 0; capsule oblong, acute, but not rostrate.—24. Presidio. Summer.

3. *J. BALTICUS* Dethard. Stem naked, terete; inflorescence lateral; flowers of the compound panicle branches solitary; leaves 0; capsule oblong, rostrate.—Common. Summer.

4. *J. COMPRESSUS* HBK. Stem naked, somewhat flattened; inflorescence lateral; flowers of the loosely few-flowered panicle solitary; leaves terete, sometimes 0; capsule oblong. 24. Santa Clara. Summer.

5. *J. BREWERI* Engelm. Stem naked, somewhat flattened; inflorescence lateral; flowers of the small, but dense panicle, solitary on their branches; leaves 0; capsule oblong.—24. Santa Cruz. Summer.

6. *J. EFFUSUS* L. Stem naked; inflorescence lateral; flowers of the panicle solitary on their branches, triandrous; leaves 0; capsule clavate obtuse.— \mathcal{L} . Presidio. Summer.

7. *J. PATENS* Meyer. Stem naked; inflorescence lateral; flowers on panicle branches solitary, hexandrous; leaves 0; capsule subglobose. \mathcal{L} . Common. Summer.

8. *J. BUFONIUS* L. Stem leafy, branched; inflorescence terminal; flowers remote, hexandrous.— \odot . Common. Summer.

9. *J. KELLOGGI* Engelm. Stem leafy, branched; inflorescence terminal; peduncles 1 or 2-flowered; flowers triandrous.— \odot . San Francisco. Spring.

10. *J. TENUIS* Willd. Stem naked, simple; inflorescence terminal, a diffuse cyme of solitary flowers.— \mathcal{L} . Lake Chabot. Summer.

11. *J. FALCATUS* Meyer. Stem leafy; leaves sheathing, flat.— \mathcal{L} . Lone Mountain. San Francisco. Summer.

12. *J. XIPHIODES* Meyer. Stems leafy, ancipital; leaves equitant, laterally compressed; flowers capitate; heads paniculate; anthers exceeding the filaments.— \mathcal{L} . Coast Range. Summer.

13. *J. PHÆOCEPHALUS* Engelm. Stem leafy, ancipital; leaves equitant, laterally compressed; flowers capitate; heads paniculate; anthers ex-

ceeding the filaments.— \mathcal{L} . Coast Range. Crystal Springs. Summer.

2. *Luzula* DC. WOOD-RUSH.

Perigonium 6-sepalous, all the segments flat. Ovary 1-celled, 3-ovulate. Capsule 3-valved, 3-seeded.— \mathcal{L} .

1. *L. COMOSA* Meyer. Common. Spring.

SERIES 2. MICRANTHÆ. Flowers inconspicuous. Inflorescence ∞ -flowered.

ORDER 1. GLUMACEÆ. Flowers in the axils of bracts, and arranged in spikelets. Spathes 0. Perigonium depauperate. Ovary 1-celled, 1-seeded. Fruit a caryopsis, with endosperm.

Family 1. GRAMINEÆ.

Stem articulate. Leaves distichous, alternate, sheathing, with ligule (stipule) at base. Flowers protected by an anterior and a posterior bractlet (palet). Spikelet generally protected by one or two bracts (glumes). Stamens usually 3. Pistils 2.

1. *Panicum* L. PANIC-GRASS.

Spikelets 2-flowered; upper one ♀ , lower reduced to a single palet. Glumes unequal, the lower smaller. Leaves flattened.

1. *P. SANGUINALE* L. Spikelets in pairs; one sessile, the other pedicillate; crowded on

one side of the simple flattened branches, which are digitately clustered at the top of the culm.—⊙. Common. Summer.

Probably introduced from Europe.

2. *P. AGROSTOIDES* Spreng. Spikelets disposed in panicles; panicle elongated, racemose. ♀. San Jose. Summer.

3. *P. CAPILLARE* L. Panicle diffuse and spreading; spikelets scattered, disposed in panicles and pointed.—⊙. San Francisco. Summer.

4. *P. DICHOTOMUM* L. Panicle diffuse and spreading. Spikelets scattered, disposed in panicles and obtuse.—⊙. Marin County. Summer.

5. *P. CRUSGALLI* L. Spikelets crowded on the secund, spikelike branches of the panicle. ⊙. Common. Summer.

Introduced from Europe.

2. *Phleum* L. TIMOTHY.

Spikelets 1-flowered, ♀. Glumes 2; aristate; the upper palea sometimes bearing at its base the rudiment of a second flower. Leaves flattened. Panicles spikelike.

1. *P. PRATENSE* L. Awn shorter than its glume.—♀. Common. Summer. Native of Europe.

2. *P. ALPINUM* L. Awn about as long as its glume.—♀. San Francisco. Summer.

3. Alopecurus L. FOXTAIL.

Spikelet 1-flowered, ♀. Glumes 2, navicular, connate at their base; lower palet carinate; upper wanting.—♂: Leaves flattened. Panicles spikelike.

1. *A. FRATENSIS* L. Culm erect.—Niles. Summer.

2. *A. GENICULATUS* L. Culm procumbent, ascending from the lower nodes.—Marin County. Summer.

4. Beckmannia Host. CATERPILLAR GRASS.

Spikelets 1 or 2-flowered. Flowers ♀, sessile. Glumes 2, navicular; lower palet ovate, 3-nerved, embracing the upper 2-cleft, 2-nerved, one.—♂. Leaves flattened. Spike sessile; spikelets alternate, sessile, 1-sided, 2-seriate.

1. *B. ERUCÆFORMIS* Host.—Marin County. Summer.

5. Phalaris L. CANARY GRASS.

Spikelets 3-flowered, upper ♀, 2 lower neutral, depauperate. Glumes 2, carinate. Palets 2, navicular; the lower larger and embracing the upper. Spikelets pedicillate. Leaves flattened.

1. *P. CANARIENSIS* L. Panicle dense and spikelike; glumes broad with a broad keel.—⊙. Common near houses. Spring.

“Canary grass.” Native of the Mediterranean region; used as bird’s food and probably escaped with the rubbish of birds’ cages.

2. *P. INTERMEDIA* Bosq. Panicle dense and spike-like; glumes pointed with a broad keel. ☉. Common. Spring.

3. *P. AMETHYSTINA* Trin. Panicle dense and spike-like; glumes with a narrow keel.— \mathcal{L} . Contra Costa hills. Summer.

4. *P. ARUNDINACÆ* L. Panicle branched; glumes not all carinate, pointed, 3-nerved.— \mathcal{L} . Niles. Summer.

6. *Hierochloa* Gmelin. HOLY GRASS.

Spikelets 3-flowered; flowers sessile. Glumes equal, carinate. Upper flower ♂; lower palet carinate; upper 1-nerved. 2 lower flowers ♂; lower palet carinate, upper 2-carinate.— \mathcal{L} . Spikelets pedicillate. Leaves flattened.

1. *H. MACROPHYLLA* Thurb.—Tamalpais. Summer.

Fragrant grass, perhaps containing cumarine like our *Galium triflorum* and the European *Asperula odorata* (Waldmeister).

7. *Anthoxanthum* L. SWEET VERNAL-GRASS.

Spikelets 3-flowered. Glumes 2, carinate; lower glume shorter, 1-nerved; upper 3-nerved. Upper flower ♀, with 2 navicular palets without awns; lower palet wrapt round the upper.

Stamens 2. Lower 2 flowers neutral, with a single, aristate, canaliculate palet.—2♂. Leaves flattened. Panicle contracted.

1. *A. ODORATUM* L.—Marin County. Summer.

Fragrant grass, probably introduced from Europe.

8. *Polypogon* Desf. BEARD GRASS.

Spikelets 1-flowered. Glumes 2, carinate, aristate, much longer than the flower. Palets 2, upper 2-carinate, lower truncate at the apex. Leaves flattened.

1. *P. MONSPELIENSIS* Desf. Glumes notched, their awns 2 or 3 times their own length.—⊙. Coast Range. Spring.

2. *P. LITTORALIS* Smith. Glumes tapering into an awn about their own length.—2♂. San Francisco. Summer.

9. *Agrostis* L. BENT-GRASS.

Spikelets 1-flowered. Glumes 2, carinate, awnless, larger than the flowers. Palets 2; lower sometimes aristate; upper 2-carinate, sometimes wanting. Tufted. Leaves sometimes involute. Panicles with verticillate branches.

1. *A. ALBA* L (*vulgaris* With). Glumes nearly equal, upper palet nearly half the length of the lower.—2♂. Common, although not indigenous. Summer.

Native of Europe.

2. *A. VERTICILLATA* Vill. Glumes nearly equal; upper palet nearly as long as the lower. 2f. San Francisco. Summer.

3. *A. EXARATA* Trin. Glumes nearly equal; upper palet minute, about the length of the ovary; lower palet 5-nerved and marked on the back by a longitudinal furrow.—2f. San Francisco. Summer.

4. *A. SCABRA* Willd. Glumes unequal, the lower longer, acute and scabrous; upper palet very minute, sometimes 0.—○. San Francisco. Spring.

10. *Gastridium* Pal.

Spikelets 1-flowered. Glumes 2, ventricose at the base, much larger than the flower and closed. Palets 2; the inferior sometimes aristate, embracing the superior 2-carinate one.—○. Leaves flattened. Panicles contracted, spikelike.

1. *G. AUSTRALE* Beauv.—San Francisco. Spring.

11. *Stipa* L. FEATHER-GRASS.

Spikelets 1-flowered, flowers stipitate. Glumes 2, membranaceous, larger than the flower. Palets 2, involute, superior shorter, 2-nerved; inferior aristate, with a simple twisted awn. Ovary stipitate; caryopsis terete, and closely wrapt in the palets.—2f. Spikelets pedicellate, paniculate.

1. *S. SETIGERA* Presl. Panicle open, with spreading rays; lower palet tuberculate, only the nerves hairy.—Common. Summer.

2. *S. EMINENS* Cas. Panicle open, with spreading rays; lower palet hairy all over.—Coast Ranges. Summer.

3. *S. VIRIDULA* Trin. Panicle narrow, with short erect rays.—Coast Range. Summer.

All the species of *Stipa* are by their long awns more or less injurious to the wool of sheep, which they make impure and intractable. Some species even endanger the life of the animals, as their awns are liable to work their way from the wool through the skin into the lungs.

12. *Deyeuxia* Clar. (*Calamagrostis* Adans).

Spikelets 1-flowered, flowers sessile, with a bearded base and the rudiment of a superior flower, reduced to a plumose pedicel. Glumes 2, canaliculate, awnless. Palets 2; inferior aristate, superior 2-carinate.— σ . Leaves flattened. Panicles branched.

1. *D. ALEUTICA* Trin.—San Francisco. Summer.

13. *Spartina* Schreb. MARSH GRASS.

Spikelets 1-sided, sessile, 1-flowered; flower naked, sessile. Glumes 2, carinate, awnless; the upper embracing the lower, which is much smaller. Palets 2, awnless; the lower compressed; the upper larger navicular. Ovary

sessile.— σ . with creeping rhizome, rigid, and with involute leaves.

1. *S. STRICTA* Roth.—Salt marshes. San Francisco. Summer.

14. *Cynodon* Rich. BERMUDA GRASS.

Spikelets 1-sided, sessile, 1-flowered, often with rudiment of a superior flower. Glumes 2, carinate, awnless, the upper embracing the lower. Palets 2; the lower carinate, acute. Ovary sessile.— σ . Rhizome creeping. Leaves flattened. Spikes in our species digitate.

1. *C. DACTYLON* Pers.—San Jose. Summer.

15. *Danthonia* DC.

Spikelets 2- ∞ -flowered; rachis hairy. Flowers distichous, the uppermost depauperate. Glumes 2, awnless, somewhat longer than the flower. Palets 2; upper 2-carinate; lower concave, ∞ -nerved, 2-cleft, aristate. Ovary stipitate; caryopsis compressed, free.— σ . Turfy. Leaves flattened. Spikelets pedicellate, paniculate.

1. *D. CALIFORNICA* Bolander.—Contra Costa hills. Summer.

16. *Avena* L. OAT.

Spikelets 3-flowered, the uppermost depauperate. Glumes 2, awnless. Palets 2; lower 2-cleft, aristate; upper 2-carinate, awnless.

Ovary sessile, hirsute at the apex; caryopsis terete, adherent to the upper palet.—⊙.

1. *A. FATUA* L.—Common. Summer.

17. *Trisetum* Kunth.

Spikelets 2–4-flowered, the uppermost depauperate. Glumes 2, carinate, awnless, shorter than the flower. Palets 2; lower 2-dentate, aristate. Ovary sessile; caryopsis compressed, free. Leaves flattened.

1. *T. CANESCENS* Buckl. Spikelets 2 or 3-flowered, narrow; lower glume narrow, considerably shorter than the broad ovate upper one.—⊙. Common. Spring.

2. *T. BARBATUM* Steud. Spikelets 3–5-flowered, large, much flattened; both glumes narrow, the lower but little shorter than the upper one.—⊙. Common. Spring.

18. *Aira* L. HAIR-GRASS.

Spikelets 2-flowered, often with the rudiment of a superior one. Flowers sessile. Glumes 2, carinate, awnless, larger than the flower. Palets 2; the lower 2-cleft, dorsally aristate. Ovary sessile; caryopsis free.

1. *A. CÆSPITOSA* L. Glumes not longer than the florets; spikelets much compressed.—♀. San Francisco. Summer.

2. *A. HOLCIFORMIS* Steud. Glumes not longer than the florets; spikelets nearly terete.—♀. Common. Summer.

3. *ELONGATA* Hook. Glumes longer than the florets; panicle long and narrow, its rays unequal, distant, mostly appressed.—24. San Francisco. Summer.

4. *A. DANTHONOIDES* Trin. Glumes longer than the florets; panicle loose and open, the lowermost rays in threes, the upper in pairs or solitary.—24. Oakland. Summer.

19. *Arrhenatherum* Beauv.

Spikelets 3-flowered : lower ♂ ; middle ♀ ; upper neutral. Glumes 2, concave, awnless; the upper larger than the lower. Palets of ♂ flower 2; lower dorsally aristate near the base; upper 2-carinate, awnless. Palets of ♀ flower 2; lower dorsally aristate near the apex, upper adnate to the caryopsis.—24. Leaves flattened. Panicles with verticillate branches. Spikelets pedicillate.

1. *A. AVENACEUM* Beauv.—Common. Summer.

European species, probably introduced as a fodder grass.

20. *Holcus* L.

Spikelets 2-flowered ; flowers pedicillate ; lower ♀ ; upper ♂ . Glumes 2, navicular. Palets of the ♀ flower 2; lower navicular, awnless; upper 2-carinate. Ovary sessile, pyriform; caryopsis free. Palets of the ♂ flower 2; upper 2-carinate ; lower aristate near the

apex. Leaves flattened. Panicles branched. Spikelets pedicellate.

1. *H. LANATUS* L.— \mathcal{L} . Common. Summer.

21. Phragmites Trin. REED.

Spikelets 3-6-flowered; flowers distichous, somewhat remote; the lower one δ , the rest \varnothing . Glumes 2, carinate, acute; the upper larger. Palets 2; lower elongate, subulate, upper 2-carinate. Ovary sessile; caryopsis free.— \mathcal{L} . Aquatic. Leaves broad, flattened. Panicles diffused.

1. *P. COMMUNIS* Trin.—Common on river banks. Summer.

22. Dactylis L. ORCHARD GRASS.

Spikelets 1-sided, 2-7-flowered. Glumes 2, carinate, mucronate aristate. Palets 2; upper 2-carinate, lower 5-nerved, mucronate aristate; carina ciliate. Ovary sessile. Caryopsis free. \mathcal{L} . Leaves carinate. Panicles glomerate, 1-sided.

1. *D. GLOMERATA* L.

Introduced from Europe as a fodder-grass.

23. Kæleria Pers.

Spikelets 2-7-flowered. Flowers distichous. Glumes 2, carinate, awnless, unequal. Palets 2; lower enlarged, upper 2-carinate, 2-cleft. Caryopsis terete, free. Leaves flattened. Pani-

cles contracted, spikelike. Spikelets pedicellate.

1. *K. CRISTATA* Pers.— \mathcal{M} . San Francisco. Summer.

24. *Melica* L. MELIC-GRASS.

Spikelets 3-5-flowered, 2 inferior flowers \varnothing , the rest depauperate. Glumes 2, concave, awnless, unequal. Palets 2, sessile, smooth. Caryopsis terete, free. Leaves flattened. Spikelets pedicellate.

1. *M. IMPERFECTA* Trin. Glumes nearly equaling the florets; lower palet scarious margined, 7-nerved; spikelets one \varnothing floret and one or two sterile.— \mathcal{M} . Cemetery. Summer.

2. *M. BULBOSA* Geyer. Glumes nearly equaling the florets; lower palet scarious-margined, 7-nerved; spikelets contain 2 or 3 \varnothing florets.— \mathcal{M} . Coast Range. Summer.

3. *M. BROMOIDES* Gray. Glumes distinctly shorter than the lower floret; lower palet acute. \mathcal{M} . Marin County. Summer.

4. *M. HARFORDI* Bolander. Glumes distinctly shorter than the lower floret; lower palet truncate.— \mathcal{M} . Nicasio. Summer.

25. *Brizopyrum* Link. (*Distichlis* Raf.).

Dioecious. Spikelets compressed, ∞ -flowered. Glumes 2, narrow, carinate. Palets 2: lower coriaceous, ∞ -nerved, not carinate; up-

per carinate with involute margins. Ovary stipitate; caryopsis obovate, free.— γ . Creeping. Leaves rigid, distichous, spreading. Panicle spicate-racemose.

1. *B. MARITIMUM* Raf.—San Francisco. Salt marshes. Summer.

26. *Lophochlæna* Nees.

Spikelets ∞ -flowered. Flowers δ . Rachis articulate, deciduous. Glumes 2, shorter than the flowers, upper 3-nerved, lower smaller, 1-nerved. Palets 2: lower chartaceous, ∞ -nerved, membranaceous at the apex, 2-lobed, aristate from the cleft; upper complicate, chartaceous, margin and apex membranaceous, 2-nerved, 2-carinate, margin dentate, apex emarginate. Caryopsis compressed, 2-horned. Leaves short. Panicle secund (one-sided), simply racemose.

1. *L. CALIFORNICA* Nees.— \odot . Contra Costa. Spring.

27. *Glyceria* R. Br.

Spikelets ∞ -flowered. Glumes 2, obtuse, the lower shorter. Palets 2: upper 2-carinate, lower concave, ovate, rotund, 7-nerved. Caryopsis free. *Aquatics*. Leaves flattened. Branches of the panicle semi-verticillate.

1. *G. PAUCIFLORA* Presl.— \odot . San Francisco. Spring.

28. *Poa* L.

Spikelets 2- ∞ -flowered. Glumes 2, obtuse. Palets 2, both deciduous, lower carinate, upper 2-carinate. Ovary sessile.

1. *P. DISTANS* Grieseb. Lower palet rounded; all leaves short and narrow, mostly convolute.— \mathcal{L} . San Francisco. Summer.

2. *P. CALIFORNICA* Munro Ms. Dioecious; lower palet rounded; radical leaves half as long as the culm, mostly flat; culm-leaves short, frequently reduced to a mucro.— \mathcal{L} . San Francisco. Summer.

3. *P. TENUIFOLIA* Nutt. Lower palet rounded; radical leaves exceedingly narrow, mucronate; culm-leaves but little wider.— \odot . San Francisco. Spring.

4. *P. SCABRELLA* Gray. Dioecious; lower palet rounded. leaves carinate.— \mathcal{L} . Oakland. Summer.

5. *P. ANNUA* L. Lower palet acute; branches of the panicle single or in pairs; all leaves flat. \odot . Common. Spring.

6. *P. PRATENSIS* L. Florets \mathcal{F} ; lower palet acute; branches of the panicle mostly in fives; the entire plant smooth, only the margins of the leaves slightly scabrous.— \mathcal{L} . Common. Summer.

7. *P. TRIVIALIS* L. Florets \mathcal{F} : lower palet acute; branches of the panicle mostly in fives; the entire plant rough.— \mathcal{L} . Common. Summer.

8. *P. DOUGLASII* Nees. Dioecious; lower palet acute; radical leaves setaceously convolute.— \mathcal{L} . San Francisco. Sand dunes. Summer.

29. *Eragrostis* Beauv.

Spikelets 2- ∞ -flowered. Glumes 2, obtuse. Palets 2; upper palet persistent.

1. *E. POÆOIDES* Beauv.— \odot . San Francisco. Spring.

Probably introduced from Europe.

30. *Briza* L. QUAKING-GRASS.

Spikelets ∞ -flowered; flowers imbricate, distichous. Glumes 2, concave, rounded, ventricose. Palets 2: lower rotund, concave, with cordate base; upper much smaller, 2-carinate. Caryopsis compressed. Leaves flattened. Spikelets pedicellate, paniculate.

1. *B. MEDIA* L.— \mathcal{L} . Common. Summer.

31. *Festuca* L. FESCUE GRASS.

Spikelets 2- ∞ -flowered; flowers distichous. Glumes 2, carinate, unequal. Palets 2; lower not carinate, mucronate, sometimes aristate; upper 2-carinate. Ovary sessile, smooth.

1. *F. MYURUS* L. Monandrous; panicle contracted, spikelike, narrow, rays appressed; lower glume minute; upper glume half as long as the lowest floret.— \odot . San Francisco. Spring.

2. *F. TENELA* Wild. Diandrous; panicle contracted, spikelike, lower rays in unequal pairs; glumes acute, the lower at least half as long as the upper.—⊙. San Francisco. Spring.

3. *F. MICROSTACHYS* Nutt. Monandrous; panicle contracted spike like; branches not appressed; glumes acute, the lower 1-nerved and smaller than the 3-nerved upper; upper glume almost as long as the lowest floret.—⊙. Common. Spring.

4. *F. OVINA* L. Panicles loose, not spike-like; rays mostly solitary; glumes about equal; lower palet indistinctly nerved.—♂. Coast Range. Summer.

5. *F. SCABRELLA* Torr. Panicles loose, not spikelike; lower rays distant in pairs; upper glume somewhat longer than the lower and half as long as the lowest floret; lower palet membranaceous, distinctly 5-nerved.—♂. Oakland. Summer.

6. *F. PAUCIFLORA* Thunb. Panicles loose, not spikelike; lower rays distant in pairs; glumes narrow, the upper twice as large as the lower; lower palet distinctly 5-nerved.—♂. San Francisco. Summer.

32. *Bromus* L. BROME-GRASS.

Spikelets 3-∞-flowered; flowers distichous; glumes 2, unequal. Palets 2: lower rotund, convex, sometimes aristate; upper 2-carinate,

the carina ciliate. Ovary sessile; hiscite at the apex. Leaves flattened; spikelets pedicellate, paniculate.

1. *B. MAXIMUS* Desf. Lower glume 1-nerved, upper 3-nerved; glumes hyaline, the upper almost equalling the floret; lower palet scabrous.—○. San Francisco. Mission. Spring.

2. *B. SECALINUS* L. Lower glume 3-5-nerved, upper 5-7-nerved; lower palet, inside rounded, convex, outside, carinate; rays of the panicle spreading, even when in fruit.—○. Mission. Spring.

3. *B. RACEMOSUS* L. Lower glume 3-5-nerved, upper 5-7-nerved; lower palet, inside rounded, convex, outside, carinate; rays of the panicle contracted when in fruit.—○ ○. Oakland. Spring.

4. *B. CILIATUS* L. Lower glume 1-nerved, upper 3-nerved; glumes acute; the upper more than half the length of the lower floret; lower palet silky.—ㄥ. San Francisco. Summer.

5. *B. GRANDIFLORUS* Hook (*Ceratochloa* Beauv.). Lower glume 5-nerved, the upper 9-nerved; lower palet compressed, carinate.—ㄥ. Nicasio. Summer.

33. *Lepturus* R. Br.

Spikelets 1-flowered, with rudiment of a superior flower. Glumes 2, rigid, subulate in the terminal spikelet opposite, in the lateral

ones collateral; lower glume sometimes wanting. Palets shorter than the glumes, the lower enclosing the upper.—⊙. Leaves narrow and flattened. Spikelets solitary, immersed into alternate excavations of the rachis, the inflorescence representing a slender spike.

1. *L. INCURVATUS* Trin. Tiburon. Marin County. Spring.

Native of the Mediterranean region.

34. *Lolium* L. Darnel.

Spikelets ∞ -flowered, flowers imbricate, distichous. Glumes 2, blunt, the posterior often wanting. Palets 2: the lower concave, sometimes aristate; upper 2-carinate with the caryopsis adhering. Leaves flattened. Spikelets solitary, immersed into alternate excavations of the rachis, placed edgewise with it, and representing a single spike.

1. *L. PERENNE* L. Glume much shorter than the spikelet.— \mathcal{L} . Common. Summer.

Native of Europe, but frequently escaped from cultivation.

2. *L. TEMULENTUM* L. Glume as long as the spikelet.—⊙. Common. Spring.

Has the reputation of being a narcotic poison, but varies considerably in regard to intensity and character of the symptoms caused by it.

35. Triticum L. WHEAT.

Spikelets 3- ∞ -flowered. Flowers distichous, placed with their flat sides to the rachis. Glumes 2, lower sometimes wanting. Palets 2, lower rounded, upper 2-carinate, the carinate ciliate. Leaves flattened. Spikelets solitary, immersed into the excavations of the rachis, representing a single spike.

1. *T. REPENS* L.—2f. Common. Summer.

36. Hordeum L. BARLEY.

Spikelets 1-flowered, with the rudiment of a superior floret, ternate; the central spikelet σ , the lateral ones depauperate. Glumes 2, linear-lanceolate, aristate; all anterior. Palets 2; lower concave, aristate. Leaves flattened. Spikelets representing a single spike.

1. *H. NODOSUM* L. Glumes all setaceous from a broad base.— \odot \odot . Oakland. Summer.

2. *H. MURINUM* L. Glumes of the central spikelet lanceolate with a long awn and ciliate; outer glumes of the lateral spikelets setaceous; inner glumes like those of the central spikelet.

\odot . San Francisco. Spring.

Native of Europe.

3. *H. JUBATUM* L. Glumes capillary, running into very long awns.— \odot \odot . Marin County. Summer.

“Squirrel-tail.”

37. *Elymus* L. WILD-RYE.

Spikelets 1- ∞ -flowered, all $\text{\textcircled{f}}$; florets distichous, the uppermost rudimentary. Glumes 2, equal, anterior. Palets 2, low or concave. Ovary hirsute.— $\text{\textcircled{f}}$. Leaves flattened. Spikelets representing a single spike.

1. *E. ARENARIUS* L. Glumes acuminate, shorter than the spikelet; lower palet carinate towards the cuspidate tip; leaves pungent.—Cliff House. Summer.

Introduced under the name "Esparto Grass," to keep the sand dunes from moving.

2. *E. CONDENSATUS* Presl. Glumes subulate-setaceous, shorter than the spikelet; lower palet 5-nerved, mucronate, sometimes shortly aristate; leaves ample, flat.—San Francisco. Summer.

3. *E. SIBIRICUS* L. Glumes linear, 3-5-nerved, acute or shortly aristate; lower palet aristate, the awn longer than the palet.—San Francisco. Summer.

4. *E. SITANION* Schult. Glumes very long, sometimes 2-parted, ending in several very long awns.—San Rafael. Summer.

38. *Gymnostichum* Schreb. BOTTLE-BRUSH GRASS.

Spikelets 1-4-flowered; florets remote, the uppermost rudimentary. Glumes rudimentary, often wanting. Palets 2, lower aristate, em-

bracing the upper. — 2f. Leaves flattened. Spikelets representing a spike.

1. *G. CALIFORNICUM* Bolander.—Saucelito. Summer.

Family 2. CYPERACEÆ.

Stem a calamus. Leaves all radical, sheathing but not split at their base; lamina parallel-veined. Anthers basifixed.

1. *Carex* L. SEDGE.

Flowers diclinic. ♂ spikelets 1-flowered; glume 1, external, ♀ spikelets 1-flowered with 2 glumes; the external like that of the ♂; the internal transformed into a utricle including the ovary.—2f. Leaves flattened. Stem triquetrous. Spikelets collected in diclinic or androgynous spikes.

1. *C. MARCIDA* Boott. Inflorescence the compound of many sessile spikes; spikes irregularly androgynous, sometimes the whole inflorescence ♀; stigmas 2; utricle (perigynium) ovate with a short 2-dentate apex; spikes almost black.—Santa Clara marshes. Summer.

2. *C. DOUGLASHII* Boott. Dioecious; inflorescence the compound of many sessile spikes; utricle (perigynium) ovate, acuminate into a slender beak; spikes pale-brown.—Marin County. Summer.

3. *C. MURICATA* L. Inflorescence the compound of many sessile spikes; ♂ spikes or androgynous spikes at the top; inflorescence oblong, the compound of 4-10 spherical spikes; stigmas 2; bracts ovate, aristate, longer than the spikes; spikes chestnut-color.—Marin County. Summer.

4. *C. GLOMERATA* Thunb. Inflorescence the compound of many sessile spikes; ♂ or androgynous spikes at the top; inflorescence elongated, the compound of ∞-small rounded spikes; stigmas 2; bracts setaceous, longer than the spikes; spikes brown.—Common. Summer.

5. *C. PANICULATA* L. Inflorescence the compound of many sessile spikes; ♂ or androgynous spikes on the top; inflorescence almost linear, the compound of oblong spikes (sometimes branched); stigmas 2; bracts setaceous; spikes brown or pale.—Presidio. Summer.

6. *C. FESTIVA* Dewey. Inflorescence the compound of many sessile spikes, forming a capitulum; ♂ spikes at the bottom or centre; spikes brown.—Oakland. Summer.

7. *C. DEWEYANA* Schwein. Spikes of the inflorescence distinct, androgynous; ♂ florets at the base; stigmas 2; spikes pale.—Napa Valley. Summer.

8. *C. BIFIDA* Boott. Spikes of inflorescence distinct, the uppermost androgynous with ♂ florets on the base, the rest of the spikes ♀,

stigmas 3; spikes dark-colored.—Marin county. Summer.

9. *C. NUDATA* Bolander. Spikes unisexual, the upper ♂, the rest ♀ (sometimes androgynous) stigmas 2; ♂ spike single (sometimes androgynous); utricule (perigynium) elliptic, compressed, slenderly nerved.—Tamalpais. Summer.

10. *C. AQUATILIS* Wahl. Spikes unisexual; the upper 1-4 spikes ♂, the lower (2-5) ♀; stigmas 2; utricule (perigynium) elliptic, stipitate, nerveless.—Wright's Station. Summer.

11. *C. SITCHENSIS* Prescott. Spikes unisexual; the upper 1-4 spikes ♂, the rest (3-5) ♀; stigmas 2, utricule (perigynium) orbicular, turgid, stipitate, coriaceous, nerveless.—Presidio. Summer.

12. *C. JAMESII* Torr. Spikes unisexual; upper 1-4 spikes ♂, the rest (3-4) ♀; stigmas 2, utricule (perigynium) oval, strongly nerved, abruptly ending in a 2-dentate beak; bracts about the length of the stem.—Sonoma. Summer.

13. *C. LACINIATA* Boott. Spikes unisexual; upper 1-2; spikes ♂, the rest (3-4) ♀; stigmas 2, utricule (perigynium) oval, lenticular compressed, bracts far exceeding the stem.—Coast Range. Summer.

14. *C. GLOBOSA* Boott. Spikes unisexual, upper spike ♂, rest (3-5) ♀; stigmas 3; utricule

(perigynium) globose, tapering to the base, abruptly rostrate; beak short.—Marin county. Summer.

15. *C. PSEUDOCYPERUS* L. Spikes unisexual, upper 1 ♂; rest (3-5) ♀, drooping; stigmas 3; utricle coriaceous, ovate, triquetrous, attenuate into a long slender beak.—San Francisco. Summer.

2. *Fimbristylis* Vahl.

Spikelets ♀, ∞-flowered; palets closely imbricate all round, the lowermost empty. Perigonium 0. Style incrassate at base, persistent. Leaves narrow. Spikelets, if solitary, bracteate; if capitate or umbellate, involucrate.

1. *F. MILIACEA* Vahl.—⊙. San Francisco. Folsom street. Spring.

This tropical species has not been found again since the filling in of that part of the city.

3. *Isolepis* L.

Spikelets ♀, ∞-flowered. Palets imbricate all round, the lowermost empty. Perigonium 0. Base of the style deciduous, cespitose, slender.

1. *I. RIPARIA* R. Br. Nutlets triquetrous, the sides convex.—⊙. Common. Spring.

2. *I. CARINATA* Hook. & Arn. Nutlets triquetrous, the sides straight.—⊙. San Francisco. Spring.

4. *Scirpus* L. BULRUSH.

Spikelets ♂, ∞-flowered. Palelets imbricate all round, the lowermost empty. Perigonium 6-setaceous. Style articulate with its base.

1. *S. LACUSTRIS* L. Inflorescence apparently lateral, with a single erect involucre leaf; stem almost terete; bristles of perigonium 6, slender, with scattered barbs—2f. Common. Summer.

2. *S. TATORA* Kunth. Inflorescence apparently lateral, with a single erect involucre leaf; stem almost terete; bristles of perigonium less than 6, stout and retrorsely plumose.—2f. Benicia. Summer.

3. *S. OLNEYI* Gray. Inflorescence apparently lateral, with a single erect involucre leaf; stem triquetrous.—2f. Presidio. Summer.

4. *S. MARITIMUS* L. Involucre spreading; spikelets large in a sessile cluster or sparingly umbellate.—2f. Common. Summer.

5. *S. SYLVATICUS* L. Involucre spreading; spikelets small in a supra-decompound inflorescence.—2f. Common. Summer.

6. *S. ACICULARIS* L. (*Eleocharis* R. Br.). Spike distichous or 3-ranked, few-flowered, terminating a leafless stem.—⊙. Common. Spring.

7. *S. PALUSTRIS* Reichenb. (*Eleocharis* R. Br.). Spike terete, ∞-flowered, terminating a leafless stem.—2f. Common. Summer.

5. **Eriophorum** L. COTTON-GRASS.

Spikelets ♂, ∞-flowered. Palets imbricate all round, the lowermost empty. Perigonium ∞-setaceous, in fruit exerted, and silky.—2f.

1. **E. GRACILE** Koch.—San Francisco. Summer.

Aquatic; formerly found in a swamp near the mouth of Mission creek. Not yet rediscovered in our neighborhood, but frequent in the higher Sierras.

6. **Cyperus** L.

Spikelets ♂. Palets imbricate, distichous, the lowermost empty. Perigonium 0. Inflorescence involucrate.

1. **C. DIANDRUS** Torr. Style 2-cleft; nutlet lenticular, the edge turned to the rachis of the spikelet.—⊙. San Francisco. Summer.

2. **C. ARISTATUS** Rottl. Style 3-cleft; nutlet triquetrous. Floret monandrous.—2f. Common. Spring.

ORDER 5. SPADICIFLORÆ. Inflorescence a spadix surrounded by a spathe. Bracts depauperate. Ovary superior, one to several-celled; cells 1-ovulate. Leaves alternate with sheathing base.

Family 1. **TYPHACEÆ.**

Flowers declinous, in a sometimes interrupted club-shaped spadix, each division of which is

protected by foliaceous spathes. Upper part of spadix ♂ ; perigonium 0; anthers basifixed, irregularly mixed with scales. Lower portion of spadix ♀ ; ovaries surrounded at their bases by ∞-bristles, sometimes by 3 scales; stigma 1, lateral. Fruit a utricle; endosperm copious. Leaves linear, straight-nerved. Aquatic.—24.

1. **Sparganium** TOURN. BUR-REED.

Flowers in crowded glomerulate heads, constituting a simple or a branched spadix. Filaments slender. Ovaries sessile, surrounded by some linear scales.

1. **S. EURYCARPUM** Engelm.—West Oakland (extinct); rediscovered at Niles. Summer.

2. **Typha** TOURN. CAT-TAIL. FLAG.

Flowers in a club-shaped, continuous spadix. ♂ : filaments connate, very short. ♀ : ovaries stipitate, surrounded by ∞ bristles.

1. **T. LATIFOLIA** L.—Common. Summer.

SERIES 3. HELOBIÆ. Aquatics. Type generally ternate. Seed nearly without endosperm. Embryo with very pronounced radicle.

ORDER 1. POLYCARPICÆ. Ovaries 3-∞. Endosperm entirely wanting.

Family 1. ALISMACEÆ.

Calyx 3-sepalous. Corolla 3-petalous. Stamens 6-∞. Ovaries 3-∞, with terminal styles

and ventral placentation. Carpidia dry, indehiscent. Lamina of leaves nervose.

1. **Alisma** L. WATER-PLANTAIN.

Flowers ♂. Calyx herbaceous. Corolla petaloid. Ovaries ∞ , 1-ovulate. Akenes ∞ , verticillate.— γ . Aquatic. Scapigerous. Inflorescence verticillate, paniculate.

1. A. PLANTAGO L. Flowers pale.—Common. Summer.

Family 2. JUNCAGINEÆ.

Calyx not differing from corolla. Lamina of leaves abortive.

1. **Triglochin** L. ARROW-GRASS.

Perigonium deciduous, 3 inner sepals sometimes wanting. Ovary 6-celled; ovules in each cell solitary, basal. Styles 3-6; stigmas plumose. Capsule separating into carpidia; finally opening ventrally.— γ . Scapigerous. Inflorescence spicate.

1. T. MARITIMUM L. Common. Summer.

Family 3. POTAMEÆ.

Perigonium if present, 4-cleft. Stamens 1, 2, or 4. Ovaries 4- ∞ , 1-ovulate. Carpidia indehiscent. Leaves stipulate.

1. **Potamogeton** TOURN. PONDWEED.

Flowers ♂. Perigonium 4-cleft. Anthers 4, sessile, inserted on the base of the sepals.

Ovaries 4, styles 0.— σ . Jointed aquatics. Inflorescence pedunculate; spikes axillary.

1. *P. NATANS* L. Floating leaves coriaceous; submerged leaves filiform.—Marine Hospital. Lake Merced. Taylorville. Summer.

2. *P. LUCENS* L. Leaves uniform, lanceolate.—Mission Dolores. Lake Merced. Summer.

3. *P. PAUCIFLORUS* Pursh. Leaves linear with distinct stipules.—Ocean Lake. Summer.

4. *P. PECTINATUS* L. Leaves linear, stipules united with the sheathing base of the leaf.—Common. Summer.

ORDER 2. CENTROSPERMÆ. Perigonium 0. Ovules basilar.

Family 1. NAIADACEÆ.

Flowers diclinous. Ovary 1, 1-celled, 1-ovulate. Fruit a nutlet.

1. *Lilæa* HBK.

Flowers monœcious. σ , spicate, 1-bracteate, monandrous, sometimes mingled with ♀ flowers. ♀ flowers partly spicate and 1-bracteate; partly solitary, axillary and without bracts. Styles of the upper ♀ flowers short, of the lower elongate; stigmas capitate.— \odot . Aquatic. Leaves radical, terete, sheathing at base.

1. *L. SUBULATA* HBK.—San Francisco. School House Station. Searsville. Spring.

2. *Zostera* L. EEL-GRASS.

Flowers monœcious. Spathe elongated into a lamina. Spadix flattened, dorsally naked, ventrally covered by alternate stamens and ovaries arranged in two rows. Anthers 1-celled, sessile. Ovary rostrate. Style persistent.— $\frac{1}{2}$. Stems and roots submerged.

1. *Z. MARINA* L.—Brackish inlets around the bay. Summer.

The bulbous root is edible, and is eagerly sought for by the canvas-back duck.

Family 2. **LEMNACEÆ.**

Aquatic plants without axis. Flowers enclosed in a thin membranaceous spathe. Stamens 1 or 2. Ovary sessile, 1-celled; ovules 1-6; style short. Fruit a utricle.

1. *Lemna* L. DUCKWEED.

Stamens 2.

1. *L. TRISULCA* L. Ovule solitary; fronds oblong, stalked, remaining connected.—Presidio. Autumn.

2. *L. VALDIVIANA* Philippi. Ovule solitary; fronds not stalked, soon separating; bract reniform.—Marine Hospital. Autumn.

3. *L. MINOR* L. Ovule solitary; fronds not stalked, soon separating; bract cucullate.—Outlet of a creek between the Fort and the Cliff House. Autumn.

4. *L. GIBBA* L. More than 1 ovule.—Common. Autumn.

Sub-Class 2. GYMNOSPERMÆ.

ORDER 1. CONIFERÆ.

Family 1. TAXINEÆ.

1. *Torreya* Arn. NUTMEG-PINE.

Flowers from scaly buds. ♂ : consisting of bracts on the base of the axis, imbricated in 4 rows; stamens ∞; connectives peltate; anthers 4 to each stamen. ♀ : ovule immersed into an urceolate arillus.—ꝯ. Branches opposite. Leaves linear, decurrent, rigid, mucronate.

1. *T. CALIFORNICA* Torr.—Tamalpais. Lagunitas Creek. Spring.

Wood and seed have the odor of nutmeg, the seed occasionally being used as such.

Family 2. ABIETINEÆ.

1. *Pinus* Tourn. PINE.

♂ flowers on elongated axes; the different axes crowded into an inflorescence round the base of a new shoot. ♀ flowers with fertile scales protected by bracts, which are much

smaller than the scales. Scales of the cone persistent and elevated into a tumor (umbo).— $\frac{1}{2}$. Leaves 1-5, from a squamous sheath.

1. *P. SABINIANA* Dougl. Leaves in threes; cones on a well developed peduncle; apophyses of the scales stout, projecting.—Napa Valley. Spring.

“Blue-pine.” Seeds eaten by the Indians.

2. *P. INSIGNIS* Dougl. Leaves in threes; cones on short peduncles in clusters. Scales toward the base enlarged, thick and hemispherical.— $\frac{1}{2}$. Bolinas heights. Santa Cruz mountains. Spring.

“Monterey pine.” Frequently cultivated.

3. *P. TUBERCULATA* Gordon. Leaves in threes; cones verticillate, peduncled; scales angular, enlarged by a conical umbo.—Marin County. Spring.

4. *P. MURICATA* Don. Leaves in pairs. Bolinas. Spring.

2. *Pseudotsuga* Carriere. DOUGLAS SPRUCE.

Flowers from the axils of last year's leaves, crowded; δ on elongated axes; scales of ♀ flower considerably smaller than the bracts, persistent. Cones pendulous.— $\frac{1}{2}$. Leaves distichous, petioled.

1. *P. DOUGLASHII* Carr.—Tamalpais. Santa Cruz mountains. Spring.

Family 3. CUPRESSINEÆ.**1. Sequoia Endl.**

.Cones ovate; scales ∞ , cuneiform, spreading at maturity, decussately arranged.— $\frac{1}{2}$. Leaves alternate, decurrent, carinate.

1. *S. SEMPERVIRENS* Endl. Leaves spreading, distichous. Branchlets spreading.—Marin County. Coast Range. Contra Costa Range. Spring. “Redwood.”

2. *S. GIGANTEA* Decaisne. Leaves not distichous. Branchlets pendulous.—In a gulch of the Santa Cruz mountains near Meyers' vineyard. Otherwise only in the high Sierras.

“Big tree.”

2. Cupressus TOURN. CYPRESS.

Cone globose; scales 6–10, thick, peltate, valvate, ∞ -ovulate. Seeds angulate, narrowly winged.— $\frac{1}{2}$. Leaves decussate, adnate, squamate, imbricate, not distichous.

1. *C. MACROCARPA* Hartw. Cones of 5 or 6 pairs of scales.—Monterey. In our local flora only escaped from cultivation. Spring.

Monterey Cypress.

2. *C. GOVENIANA* Gordon. Cones of 3 or 4 pairs of scales.—Nicasio. Spring.

3. Juniperus L. JUNIPER.

Fruit a galbulus. Scales succulent, uniting in fruit. Seeds osseous.— $\frac{1}{2}$. Leaves opposite, not distichous.

1. *J. CALIFORNICA* Carr.—Walnut Creek. Spring.

CLASS 2. CRYPTOGRAMÆ.

Section 1. *DICHO TOMÆ* (*Lycopodiaceæ*).
Ramification dichotomous. Each leaf producing only one sporangium, borne on the upper surface of the leaf or near its axil.

Family 1. *SELAGINELLEÆ*.

Macrosporangia and Microsporangia. Leaves small and of two different kinds.

1. *Selaginella* Beauv.

Sporangia axillary, globose, transversely dehiscent.— γ . Leaves 4–8-ranked.

1. *S. RUPESTRIS*.—Spring. Tamalpais.

Family 2. *ISOETEÆ*.

Heterosporous. All leaves long and grass-like.

1. *Isoetes* L. QUILLWORT.

Characters of the family of which it is the only genus known.

1. *sp.* Submerged aquatic.—Corte Madera. Found by Mrs. Curran, but not yet identified.

2. *sp.* Growing on places inundated only during the rainy season.—Olema.

Found by Mrs. Curran, but not yet identified.

Section 2. **FILICINEÆ.** Ramification not dichotomous. Leaves developed, frondose. Sporangia not single and never on the upper surface of the leaf.

ORDER 1. **RHIZOCARPÆ.** Sporangia in sporocarps (conceptacula); heterosporus; microsporangia, ∞ -spored; macrosporangia generally 1-spored.

Family 1. **MARSILÆACEÆ.**

Sporocarps 2-4-celled, 2-4-valved, containing macro- and microsporangia. Sporangia parietal; macrosporangia, always 1-s p o r e d. Vernation circinate.

1. **Marsilæa L.**

Macro- and microsporangia in the same sporocarp.— \mathcal{Z} . Fronds petiolate, 4-foliolate.

1. *M. VESTITA* Hook. & Grev.—Fort Point. (Extinct.)

2. **Pillularia L.**

\mathcal{Z} . Fronds filiform.

1. *P. AMERICANA* Al. Braun.—Marin County.

Family 2. **SALVINIACEÆ.**

Sporocarps 2 to ∞ , on the same petiole (transformed half of a frond), macro- and microsporangia in distinct sporocarps. Vernation not circinate.

1. **Azolla** Lam. WATER-FERN.

Sporocarps in pairs. Macrosporangia 1-spored. Aquatics, moss-like, floating, green or purplish.

1. *A. CAROLINIANA* Willd.—Presidio. Saucelito.

ORDER 2. FILICES. Isosporous. Sporangia formed from the epidermis, bursting by an elastic ring. Fronds without stipules. Vernation circinate.

Family 1. **POLYPODIACEÆ.**

Sporangia hypophyllous, somewhat globular; ring vertical.

1. **Gymnogramme** Desf. GOLD-FERN.

Sori linear, placed on branching primary veins. Indusium 0.—2f.

1. *G. TRIANGULARIS* Kaulfuss.—Common.

2. **Cheilanthes** Swartz. LIP-FERN.

Sori placed separately at the end of veinlets. Spurious indusium rudimentary, formed by the reflexed margin of the frond.—2f.

1. *C. CALIFORNICA* Nutt. Fronds green on both sides, 4-pinnatifid.—Tamalpais.

2. *C. MYRIOPHYLLA* Desv. Fronds covered underneath with brown scales; 3-4-pinnate.—Wright's Station.

3. Allosorus Benth. (*Pellaea* Link.) **ROCK-BRAKE.**

Sporangia on veins which are oblique to the midrib. Spurious indusium continuous, formed by the reflexed margin of the frond.— γ . Glabrous, cespitose ; fertile fronds contracted ; petioles not green.

1. *A. ANDROMEDÆFOLIUS* Fée. Pinnules oval, obtuse.—San Rafael. Tamalpais.

2. *A. ORNITHOPUS* Hook. Pinnules rhomboid, mucronulate.—Piedmont. Lagunitas reservoir.

3. *A. DENSUS* Hook. Pinnules linear, mucronate.—Lagunitas reservoir.

4. Polypodium L. **POLYPODY.**

Sori round, placed on the veins. Indusium 0.— γ .

1. *P. VULGARE* L. Veins of the segments uniformly free and none anastomosing.—Common.

2. *P. SCOULERI* Hook & Gray. Veins of the segments anastomosing by veinlets.—Saucelito. Tamalpais.

5 Pteris L. **BRACKEN. BRAKE.**

Sporangia at the end of the veinlets, connected into a vein-like receptacle which surrounds the frond without interruption, indusium continuous, membranaceous, attached at its marginal side, free on its inner side.— γ .

1. *P. AQUILINA* L. Common. The fronds before expanding are edible, and were used in former times as a pot herb.

6. *Adiantum* L.

MAIDEN'S HAIR. FIVE-FINGER.

Sporangia at the end of distinct veins. Sori covered by the semi-lunar reflexed margins. Indusium continuous with the margin of the leaf, free on its inner side.— \mathcal{L} . Petioles not green.

1. *A. EMARGINATUM* Hook. Rachis of the frond continuous to the apical pinnule (leaflet). Common.

2. *A. PEDATUM* L. Rachis of the frond 2-furcate, each partition bearing several pinnate branches on its apical side.—Camp Taylor. Saucelito.

7. *Blechnum* L. (*Lomaria*, Willd.) DEER-FERN.

Sporangia forming a linear sorus on each side of the mid-rib and parallel to it. Indusium membranaceous, attached to the receptacle and free on the inner side.— \mathcal{L} .

1. *B. SPICANT* Desvaux.—Tennessee Valley. Wild-wood Glen.

8. *Woodwardia* L. CHAIN-FERN.

Sporangia placed on reticulate veinlets, forming oblong sori on each side of the mid-

rib. Indusium coriaceous, attached to the receptacle, and opening at the inner side.—2f.

1. W. RADICANS Smith.—Common.

9. Asplenium L. SPLENWORT.

Sori linear, placed on veinlets. Indusium membranaceous, attached laterally to the veinlet and opening toward the mid-rib.—2f.

1. A. FILIX FŒMINA Bernh.—Marine Hospital. Tamalpais. “Lady-fern.”

10. Aspidium Swartz. SHIELD-FERN.

Sori round, placed at the end of the veinlets. Indusium umbilicate, free on all sides and affixed by a column.—2f.

1. A. RIGIDUM Swartz. Indusium reniform; fronds 2-pinnate.—Common.

Specific against the tapeworm, like its European congener, *A. Filix mas*.

2. A. MUNITUM Kaulfuss. Indusium pel-tate; fronds pinnate, pinnæ serrate.—Common.

3. A. ACULEATUM Swartz. Indusium pel-tate; fronds 2-pinnate, sometimes simply pinnate, the pinnæ deeply cut.—Tamalpais.

2. Cystopteris Bernh. BLADDER-FERN.

Sori round, placed on the middle of the veinlet, solitary in the disc of the lacinia. Indu-

sium hyaline, laterally affixed, free towards the margin of the frond.— ζ .

1. *C. FRAGILIS* Bernh.—Tamalpais.

ORDER 3 STIPULATÆ. Isosporous.
Leaves stipulate.

Family 1. OPHIOGLOSSEÆ.

Sporangia capsular developed in the parenchyma of half of the frond. Vernation not circinate.

1. *Botrychium* Swartz.

Fertile segment of the frond a compound spike. Sporangia distinct, glabrous, distichous, nearly 2-valved.— ζ .

1. *B. TERNATUM* Swartz.—San Francisco. (Extinct.)

Section 3. EQUISETACEÆ. Stem articulate; internodes surrounded by sheathing whorls of scarious leaves. Sporangia on the under side of verticillate, peltate receptacles forming a terminal spike. Spores ∞ , each furnished with two elaters.

1. *Equisetum* L.


HORSETAIL. SCOURING-RUSH.

Only genus.

1. *E. TELMATEIA* Ehrh. Sterile stems green with verticillate branches; fertile stems with-

out chlorophyll, unbranched.—Marin County.
Contra Costa. Spring.

2. *E. ROBUSTUM* Al. Braun. All stems green,
unbranched. — Fort Point. Marin County.
Spring.



INDEX.

<p>ABIETINEÆ.....326</p> <p>Abronia.....25, 200</p> <p style="padding-left: 2em;">latifolia.....200</p> <p style="padding-left: 2em;">umbellata.....200</p> <p>Acæna, 5, 6, 7, 8, 17, 25, 38</p> <p style="padding-left: 2em;">40, 41, 250.</p> <p style="padding-left: 2em;">trifida.....250</p> <p>Acanthomintha...7, 51, 136</p> <p style="padding-left: 2em;">lanceolata.....136</p> <p>Acer..23, 36, 74, 211</p> <p style="padding-left: 2em;">macrophyllum..... 211</p> <p>Achillea.....67, 108</p> <p style="padding-left: 2em;">millefolium.....108</p> <p>Achyrachæna..... 67, 102</p> <p style="padding-left: 2em;">mollis.....102</p> <p>Actæa.....48, 233</p> <p style="padding-left: 2em;">spicata.....233</p> <p>Adenocaulon.. 88</p> <p style="padding-left: 2em;">bicolor..... 88</p> <p>Adenostoma.....40, 44, 250</p> <p style="padding-left: 2em;">fasciculatum.....250</p> <p>Adenostyles 60</p> <p>Adiantum.....333</p> <p style="padding-left: 2em;">emarginatum.....333</p> <p style="padding-left: 2em;">pedatum333</p> <p>Æsculus.....35, 74, 211</p> <p style="padding-left: 2em;">Californica.....211</p> <p>Agrostis12, 300</p> <p style="padding-left: 2em;">alba300</p> <p style="padding-left: 2em;">exarata301</p> <p style="padding-left: 2em;">scabra301</p> <p style="padding-left: 2em;">verticillata301</p>	<p>Aira.....13, 304</p> <p style="padding-left: 2em;">cæspitosa..... 304</p> <p style="padding-left: 2em;">danthonoides.....305</p> <p style="padding-left: 2em;">elongata305</p> <p style="padding-left: 2em;">holciformis.....304</p> <p>Alchemilla.....35, 250</p> <p style="padding-left: 2em;">arvensis.....250</p> <p>Alfilerilla.....205</p> <p>Alder.....271</p> <p>Alisma.....35, 323</p> <p style="padding-left: 2em;">plantago.....323</p> <p>ALISMACEÆ.....322</p> <p>Allium33, 286</p> <p style="padding-left: 2em;">attenuifolium287</p> <p style="padding-left: 2em;">lacunosum287</p> <p style="padding-left: 2em;">serratum.....287</p> <p style="padding-left: 2em;">unifolium287</p> <p>Allosorus.....332</p> <p style="padding-left: 2em;">andromedæfolius.....332</p> <p style="padding-left: 2em;">densus332</p> <p style="padding-left: 2em;">ornithopus332</p> <p>Alnus74, 271</p> <p style="padding-left: 2em;">rhombifolia272</p> <p style="padding-left: 2em;">rubra272</p> <p>Alopecurus.....11, 298</p> <p style="padding-left: 2em;">geniculatus298</p> <p style="padding-left: 2em;">pratensis.....298</p> <p>Alsine.. 30, 42</p> <p>ALSINEÆ.....192</p> <p>Alum-root.....187</p> <p>Alyssum54, 227</p> <p style="padding-left: 2em;">calycinum.....227</p>
--	--

Alyssum—maritimum... 228	Anthemis—cotula..... 107
AMARANTACEÆ 197	Anthoxanthum..... 11, 299
Amarantus... 75, 77, 79, 197	odoratum 300
retroflexus 197	ANTIRRHINEÆ..... 151
Ambrosia 75, 92	Antirrhinum 53, 152
artemisiæfolia 92	vagans 153
psilostachya 92	Aphyllon 51, 142, 143
AMBROSIEÆ 92	comosum..... 143
Amelanchier 45, 252	APHANOCYCLICÆ..... 221
alnifolia..... 253	Apiastrum..... 181
Amorpha.. 56, 57, 262	angustifolium..... 181
Californica 262	Apium 27
AMPELIDEÆ..... 214	Aplopappus 61, 63, 84
Amsinckia 19, 155	ericoides 84
intermedia..... 155	linearifolius..... 84
lycopsoides..... 156	APOCYNACEÆ 129
spectabilis 155	Apocynum..... 25, 129
tesselata 155	androsæmifolium 129
AMYGDALACEÆ 245	cannabinum 129
ANACARDIACEÆ..... 208	Aquilegia 49, 234
Anagallis..... 20, 172	truncata 234
arvensis..... 172	Arabis 55, 226
Anaphalis..... 65, 90	blepharophylla..... 227
margaritacea..... 90	perfoliata..... 227
Anemone..... 49, 237	Aralia 31, 184
Grayi 237	Californica 184
Anemopsis..... 35, 38, 279	ARALIACEÆ..... 184
Californica..... 279	Arbutus 41, 174
Angelica 28, 178	Menziesii..... 175
tomentosa 178	Arceuthobium 77, 267
ANGIOSPERMÆ..... 80	occidentale 267
ANISOCARPÆ..... 80	Arctostaphylos..... 41, 174
Anoplangthus..... 51, 142	Andersonii..... 174
fasciculatus 143	pungens. 174
uniflorus 142	tomentosa..... 174
ANTHEMIDEÆ..... 107	Arenaria..... 42, 193
Anthemis 67, 107	Californica 193

Arenaria—Douglasii	193	Aster—radulinus	86
macrophylla	193	ASTEROIDEÆ	81
palustris.	193	Astragalus	58, 263
Argemone	47, 48, 230	crotalariae	263
hispidia	230	didymocarpus	263
Aristolochia	71, 268	Douglasii	263
Californica	268	leucophyllus.	263
ARISTOLOCHIÆ	268	Menziesii	263
Armeria	31, 170	pynostachyus	264
vulgaris	170	t-ner	263
Arnica	65, 111	Atriplex	75, 77, 79, 198
discoidea	111	Californicum	199
Arrhenatherum	12, 305	coronatum	198
avenaceum	305	leucophyllum	199
Arrow-grass	323	patulum	198
Artemisia	61, 69, 109	Audibertia	7, 137
Californica	110	grandiflora	137
dracunculoides	110	stachyoides	137
Ludoviciana	110	Avena	13, 303
pynoccephala	110	Azolla	331
Asarum	43, 268		
caudatum	268	Baccharis	61, 62, 67, 87
ASCLEPIADACEÆ	128	Douglasii	87
Asclepias	128	pilularis	87
Mexicana	129	viminea	87
speciosa	128	Bæria	64, 103
vestita	128	carnosa	104
Ash	132	chrysostoma	104
Aspidium	234	Fremontii	104
aculeatum	234	gracilis.	104
munitum	234	macrantha	103
rigidum	234	uliginosa	104
Asplenium	334	<i>Bahia</i>	104
Filix fœmina	334	Balsamorhiza	66, 94
Aster	63, 86	deltoidea	94
Chamissonis	86	Hookeri	94
divaricatus	86	Baneberry	233

Barbarea.....	55, 224	Blue-Curls.....	132
<i>vulgaris</i> ..	224	Blue-eyed Grass	282
Barberry.....	232	Blue-pine.....	326
Barley	314	Boisduvalia.....	37, 244
Bay-tree.....	231	<i>cleistogama</i>	244
Beard-grass	300	<i>densiflora</i>	244
Beckmannia	298	<i>glabella</i>	244
<i>erucæformis</i>	298	BORRAGINEÆ.....	155
Bed-straw	126	Botrychium	335
Beggar-ticks	96	<i>ternatum</i>	335
Bellflower... ..	122	Bottle brush grass	315
Bent-grass.....	300	Bowlesia.....	27, 183
BERBERIDÆ.....	231	<i>lobata</i>	184
Berberis.....	31, 232	Boykinia.....	30, 186
<i>aquifolium</i>	232	<i>occidentalis</i>	186
<i>nervosa</i>	232	Box-Elder	211
<i>pinnata</i>	232	Brass-buttons.....	109
<i>repens</i>	232	Bracken.....	332
Berula.....	28, 180	Brake	332
<i>angustifolia</i>	180	Brassica.	56, 225
BETULACEÆ.....	271	<i>campestris</i>	225
Bidens.....	62, 68, 96	<i>nigra</i>	225
<i>chrysanthemoides</i>	96	<i>Brickellia</i>	112
Big-root.....	122	Bridal-wreath.....	247
Big-tree.. ..	328	Briza.....	14, 310
Bigelovia.....	60, 63, 84	<i>media</i>	310
<i>arborescens</i>	84	Brizopyrum.....	77, 307
Bind-weed.....	168	<i>maritimum</i>	308
Bitter-Cress	227	Brodiaea	8, 32, 288
Blackberry.....	247	<i>congesta</i>	288
Bladder-fern.....	334	<i>grandiflora</i>	288
Bladderwort	170	<i>minor</i>	288
Blechnum.....	333	<i>terrestris</i>	288
<i>spicant</i>	333	Brome-grass.....	311
Bleeding-heart.....	228	Bromus.....	14, 311
Blennosperma.....	69, 106	<i>ciliatus</i>	312
<i>Californicum</i>	107	<i>grandiflorus</i>	312

Bromus—maximus	312	CALYCIFLORAE	238
racemosus	312	<i>Cumassia</i>	33, 287
secalinus	312	Campanula	22, 122
Brooklime	148	<i>exigua</i>	122
Brookweed	172	<i>preanthoides</i>	122
Buck-bean	131	CAMPANULACEÆ	121
Buckeye	211	Campion	191
Bulbostylis	60, 112	Canary-grass	298
<i>Californica</i>	112	Canchalagua	130
Bulrush	320	CAPRIFOLIACEÆ	124
Bur-reed	322	Capsella	54, 223
Burrielia	64, 103	<i>Bursa-pastoris</i>	223
<i>microglossa</i>	103	Caraway	181
Buttercup	235	Cardamine	54, 227
Button-bush	126	<i>oligosperma</i>	227
		<i>paucisecta</i>	227
Cakile	54, 224	Carex	72, 316
<i>Americana</i>	224	<i>aquatilis</i>	318
Calais	116	<i>bifida</i>	317
<i>Kelloggii</i>	117	<i>Deweyana</i>	317
<i>Lindleyi</i>	117	<i>Douglasii</i>	316
<i>linearifolia</i>	117	<i>festiva</i>	317
<i>Calamagrostis</i>	12, 302	<i>globosa</i>	318
Calandrinia, 30, 35, 42, 44		<i>glomerata</i>	317
45, 195.		<i>Jamesii</i>	318
<i>Menziesii</i>	195	<i>laciniata</i>	318
California Lilac	213	<i>marcida</i>	316
California Poppy	229	<i>muricata</i>	317
Calochortus	33, 290	<i>nudata</i>	318
<i>albus</i>	291	<i>paniculata</i>	317
<i>lilacinus</i>	291	<i>pseudocyperus</i>	319
<i>luteus</i>	291	<i>Sitchensis</i>	318
<i>pulchellus</i>	291	Carrot	176
<i>venustus</i>	291	Carthamus	62
CALYCANTHACEÆ	252	Carum	28, 181
Calycanthus	46, 252	<i>Kelloggii</i>	181
<i>occidentalis</i>	252	CARYOPHYLLALES	190

- Cascara sagrada* 212
Castanopsis 76, 270
 chrysophylla 271
Castilleia 52, 147
 affinis 147
 foliolosa 147
 latifolia 147
 parviflora 147
 Catchfly 191
 Caterpillar-grass 298
 Cat-tail 322
Caucalis 29, 175
 microcarpa 176
 nodosa 175
Caulanthus 55, 225
 procerus 225
Ceanothus 23, 213
 crassifolius 214
 euneatus 214
 dentatus 213
 divaricatus 213
 papillosus 214
 thrysiflorus 213
 CELASTRACEÆ 214
 CELASTRALES 212
Centaurea 62, 68, 114
 benedicta 114
 melitensis 114
 CENTROSPERMÆ 190, 234
Cephalanthus 126
 occidentalis 126
Cerastium 43, 192
 arvense 192
 pilosum 192
Ceratochloa 312
 CERATOPHYLLEÆ 279
Ceratophyllum 75, 280
Ceratophyllum — *demersum* 280
Cercis 39, 265
 occidentalis 265
Chænactis 61, 106
 lanosa 106
 Chain-fern 333
 Chamomile 108
 Chapparal 213
Cheilanthes 331
 Californica 331
 myriophylla 331
Cheiranthus 55, 226
 Chemisal 250
 CHENOPODIACEÆ 197
 CHENOPODIALES 197
Chenopodium 5, 7, 8, 25, 197
 album 197
 ambrosioides 198
 Californicum 198
 murale 198
 Cherry 246
 Chickweed 192
 Chinquapin 27
Chlorogalum 34, 286
 pomeridianum 286
Chorizanthe . 15, 34, 39, 277
 Douglasii 278
 membranacea 278
 pungens 278
Chrysanthemum 108
 segetum 108
Chrysopsis 61, 63, 84
 Oregana 84
 sessiflora 84
Cicendia 16, 130
 CICHORIACEÆ 115

Cicuta	28, 180	Collomia	21, 167
Bolanderi	180	gilioides	168
Californica	180	gracilis	167
maculata	180	heterophylla	168
CISTINEÆ	217	Columbine	234
Clarkia	37, 243	COMPOSITEÆ	80
elegans	244	CONIFERÆ	326
Claytonia	30, 195	Conium	29, 181
Chamissonis	196	maculatum	182
linearis	196	CONVOLVULACEÆ	168
parviflora	196	Convolvulus	21, 168
perfoliata	196	arvensis	169
Sibirica	195	Californicus	169
spathulata	196	luteolus	169
Cleavers	126	pentapetaloides	169
Clematis	49, 237	Soldanella	168
lasiantha	237	Coral-root	281
ligusticifolia	237	Corallorhiza	71, 281
Clintonia	33, 285	Bigelovii	282
Andrewsiana	285	multiflora	282
Clover	256	Cordylanthus	7, 52, 144
Cnicus	62, 112	filifolius	144
Americanus	113	maritimus	144
Breweri	114	mollis	144
edulis	113	pilosus	144
fontinalis	113	Corethrogyne	68, 85
Hallii	113	obovata	85
occidentalis	113	Corn-spurrey	194
quercetorum	113	CORNACEÆ	185
Cocklebur	93	Cornel	185
Coinogyne	63, 103	CORNICULATÆ	186
carnosa	103	Cornus	17, 185
Collinsia	53, 152	Californica	185
bartsiaefolia	152	Nuttallii	185
bicolor	152	CORONARIÆ	283
parviflora	152	Corylus	76, 271
sparsiflora	152	rostrata	271

Cotton-grass.....	321	Cyperus—diandrus.....	321
Cottonwood.....	217	Cypress.....	328
Cotula.....	109	Cypripedium.....	71, 282
<i>Australis</i>	109	<i>montanum</i>	282
<i>coronopifolia</i>	109	Cystopteris.....	334
Cotyledon.....	43, 189	<i>fragilis</i>	335
<i>cæspitosa</i>	189		
Cow-parsnip.....	176	Dactylis.....	13, 306
CRASSULACEÆ.....	189	<i>glomerata</i>	306
Cream Cups.....	228, 229	Dandelion.....	120
Cressa.....	169	Danthonia.....	13, 303
<i>cretica</i>	169	<i>Californica</i>	303
<i>Croton</i>	79, 201	DAPHNALES.....	266
<i>Californicus</i>	201	Darnel.....	313
CRUCIFERÆ.....	221	<i>Datisca</i>	15, 218
CRUCIFLORÆ.....	221	DATISCEÆ.....	218
CRYPTOGAMÆ.....	329	Datura.....	21, 154
CUCURBITACEÆ.....	122	<i>stramonium</i>	154
CUPULIFERÆ.....	268	Daucus.....	29, 176
CUPRESSINEÆ.....	328	<i>pusillus</i>	176
Cupressus.....	73, 328	DECANDRIA.....	39
<i>Goveniana</i>	328	MONOGYNIA.....	39
<i>macrocarpa</i>	328	DIGYNIA.....	41
Currant.....	188	TRIGYNIA.....	41
Cuscuta.....	18, 26, 169	TETRAGYNIA.....	42
<i>salina</i>	170	PENTAGYNIA.....	42
<i>subinclusa</i>	170	POLYGYNIA.....	43
Cyanotris.....	33, 287	Deer-fern.....	333
<i>esculenta</i>	287	Delphinium.....	48, 233
CYNAROIDEÆ.....	112	<i>Californicum</i>	233
Cynodon.....	11, 303	<i>decorum</i>	234
<i>dactylon</i>	303	<i>nudicaule</i>	233
Cynoglossum.....	19, 158	<i>simplex</i>	234
<i>grande</i>	158	<i>variegatum</i>	234
CYPERACEÆ.....	316	Dendromecon.....	47, 229
Cyperus.....	5, 9, 15, 321	<i>rigidus</i>	229
<i>aristatus</i>	321	Deweya.....	29, 182

Deweya—Hartwegi	182	Dirca—occidentalis	266
Kelloggii	182	DISCOPHORÆ	175
Deyeuxia	302	<i>Distichlis</i>	307
Aleutica	302	Dock	274
DIADELPHIA	57	Dodder	169
DIANDRÆ	131	DODECANDRIA	43
DIANDRIA	5	MONOGYNIA	43
DI-TRI-TETRAGYNIA	7	PENTAGYNIA	44
MONOGYNIA	5	Dodecatheon	19, 171
Dicentra	57, 228	Meadia	171
<i>chrysantha</i>	228	Dogbane	129
<i>formosa</i>	228	Dogwood	185
Dichondra	168	Douglas Spruce	327
<i>repens</i>	168	Downingia	22, 56, 70
DICHOTOMÆ	329	Duckweed	325
DICOTYLEDONES	80	Dutchman's Pipe	268
DIDYNAMIA	49	<i>Echeveria</i>	43, 189
ANGIOSPERMIA	51	<i>Echinocystis</i>	122
GYMNOSPERMIA	49	Echinospermum	19
DIGITALEÆ	151	Eel-grass	325
DIECIA	76	Elder	124
DIANDRIA	76	<i>Eleocharis</i>	9, 15, 320
DODECANDRIA	79	Ellisia	20, 26, 161
HEXANDRIA	79	<i>chrysanthemifolia</i>	162
ICOSANDRIA	79	Elymus	10, 315
OCTANDRIA	79	<i>arenarius</i>	315
PENTANDRIA	78	<i>condensatus</i>	315
POLYANDRIA	79	<i>Sibiricus</i>	315
SYNGENESIA	79	<i>Sitanion</i>	315
TETRANDRIA	77	Emmenanthe	26, 159
TRIANDRIA	77	<i>penduliflora</i>	159
Diplacus	52, 150	ENNEANDRIA	38
<i>glutinosus</i>	150	EPIGYNÆ	80, 282
DIPSACEÆ	123	Epilobium	36, 240
Dipsacus	16, 123	<i>Franciscanum</i>	241
<i>fullonum</i>	123	<i>minutum</i>	241
Dirca	38, 266		

Epipactis.....70, 280	E odium-macrophyllum.205
gigantea.....280	moschatum.....205
EQUISETACEÆ.....335	Eryngium.....27, 183
Equisetum.....335	petiolatum.....183
Telmateia.....335	Eryngo.....183
robustum.....336	Erysimum.....55, 225
Eragrostis.....13, 310	asperum.....225
pœoides.....310	Erythræa.....20, 130
Eremocarpus....75, 200	Douglasii.....131
setiger.....200	floribunda.....130
ERICACEÆ.....173	Muhlenbergii.....130
ERICALES.....172	trichantha.....131
Erigeron.....63, 68, 86	venusta.....131
Canadensis.....87	Eschscholtzia....45, 47, 229
glaucus.....87	Californica.....230
Philadelphicus.....87	Esparto Grass.....315
stenophyllus.....86	Eucharidium.....17, 244
Eriodictyon.....26, 158	concinnum.....244
glutinosum.....159	EUCYCLICÆ.....200
Eriogonum.....38, 276	Eunanus.....53, 151
angulosum.....276	Douglasii.....151
gracile.....277	Euonymus.....16, 23, 214
latifolium.....277	occidentalis.....214
nudum.....277	EUPATORIACEÆ.....112
truncatum.....277	Euphorbia.....71, 201
vimineum.....277	lathyris.....202
virgatum.....277	leptocera.....202
Eriophorum.....9, 15, 321	ocellata.....201
gracile.....321	serpyllifolia.....201
Eriophyllum.....64, 104	EUPHORBIACEÆ.....200
cæspitosum.....105	Evax.....68, 70, 89
confertiflorum.....105	caulescens.....89
stæchadifolium.....105	Evening Primrose.....240
<i>Eritrichium</i>19, 156, 157	Everlasting.....90
Erodium.....24, 39, 56, 205	False Indigo.....262
botrys.....205	False Solomon's-Seal....284
cicutarium.....205	

Fatsia.....	184	Fraxinus—Oregana.....	132
Feather-grass.....	301	Fritillaria.....	32, 290
Ferula.....	176	FUMARIACEÆ.....	228
California.....	176		
Fescue Grass.....	310	Galium.....	16, 126
Festuca.....	5, 7, 14, 310	Andrewsii.....	127
microstachys.....	311	aparine.....	126
myurus.....	310	asperimum.....	127
ovina.....	311	boreale.....	126
pauciflora.....	311	Californicum.....	127
scabrella.....	311	Nuttallii.....	127
FICOIDÆ.....	190	trifidum.....	126
FICOIDALES.....	190	triflorum.....	127
Fig-marigold.....	190	GAMOPETALÆ.....	80
Figwort.....	153	Garrya.....	78, 185
Filago.....	65, 68, 89	elliptica.....	185
California.....	90	Fremontii.....	185
FILICES.....	331	Gastridium... ..	12, 301
FILICINÆ.....	330	australe.....	301
Fimbristylis.....	9, 15, 319	Gaultheria.....	40, 173
miliacea.....	319	Shallon.....	174
Five-finger.....	233, 249	Gayophytum.....	17, 37, 241
Flax... ..	206	diffusum.....	241
Fleabane.....	86	Gentian.....	131
Fool's Parsley.....	178	Gentiana.....	16, 20, 26, 131
Foxtail.....	298	Oregana.....	131
Fragaria.....	47, 248	GENTIANACEÆ.....	130
California.....	248	GENTIANALES.....	128
Chilensis.....	248	GERANIACEÆ.....	204
Frankenia... ..	24, 32, 35, 219	GERANIALES.....	204
grandiflora.....	219	Geranium.....	39, 56, 205
FRANKENIACEÆ.....	219	Carolinianum.....	205
Franseria.....	75, 93	Giant Hyssop.....	137
bipinnatifida.....	93	Gilia.....	21, 163
Chamissonis.....	93	achilleæfolia.....	166
Fraxinus.....	6, 72, 76, 132	androsacea.....	163
dipetala.....	132	capitata.....	166

Gilia—ciliata	164	Grape	215
cotulæfolia	165	Gratiola	149
densifolia	166	ebracteata	149
densiflora	163	GRATIOLEÆ	148
dichotoma	163	Greasewood	250
inconspicua	167	Grindelia	61, 64, 81
intertexta	165	cuneifolia	82
micrantha	164	glutinosa	82
multicaulis	167	hirsutula	82
leucocephala	165	robusta	82
pusilla	163	Groundsel	111
squarrosa	164	Gum-plant	81
tenella	164	Gutierrezia	64, 81
tricolor	167	Californica	81
virgata	166	GUTTIFERALES	215
viscidula	165	Gymnogramme	331
Githopsis	22, 121	triangularis	331
specularioides	121	GYMNOSPERMÆ	326
Glaux	24, 171	Gymnostichum	10, 315
maritima	171	Californicum	316
GLUMACEÆ	296	GYNANDRÆ	280
Glyceria	14, 308	GYNANDRIA	70
pauciflora	308	DIANDRIA	71
Glycyrrhiza	58, 262	HEXANDRIA	71
lepidota	263	MONANDRIA	71
Godetia	37, 242	Habenaria	70, 281
amcena	243	elegans	281
epilobioides	243	leucostachys	281
lepada	243	Hair-grass	304
purpurea	243	HALORRHAGIDÆ	239
quadrivulnera	243	Hawkweed	119
tenella	243	Hazel	271
Golden Aster	84	Hedge-nettle	139
Golden-rod	85	HELENIÆ	103
Gold-fern	331	Helenium	65, 107
Gooseberry	188	puberulum	107
GRAMINEÆ	296		

HELIANTHEÆ.....	95	Hetercodon—rariflorus...	122
Helianthella.....	68, 95	<i>Heteromeles</i>	45, 252
California.....	95	Heuchera.....	26, 30, 187
Helianthemum	48, 217	micrantha.....	187
scoparium.....	217	pilosissima.....	188
Helianthella	68, 95	HEXANDRIA.....	31
California.....	95	HEX-POLYGYNIA.....	31
Helianthus.....	68, 95	MONOGYNIA.....	35
annuus.....	95	TRIGYNIA.....	34
Californicus	96	Hieracium.....	60, 119
exilis ..	95	albiflorum	119
scaberrimus	95	Hierochloa.....	12, 299
Heliotrope	158	macrophylla.....	299
Heliotropium.....	18, 158	Hippuris.....	5, 239
curassavicum	158	vulgaris.....	239
HELOBIÆ.....	322	Holcus.....	13, 305
Hemizonia... 66, 68, 70, 98		lanatus	306
angustifolia.....	99	Holozonia	66, 69, 100
corymbosa	99	filipes	100
luzulæfolia	98	Holy Grass	299
macradenia ..	98	Honeysuckle	125
multiglandulosa.....	100	Hop-tree	208
Parryi	99	Hordeum	10, 314
pungens.....	99	jubatum.....	314
truncata	100	murinum	314
Hemlock	181	nodosum	314
Hendecandra.....	79, 201	Horehound.....	138
procumbens	201	Horkelia.....	45, 46
HEPTANDRIA.....	35	Horsetail.....	335
MONOGYNIA	35	Hosackia.....	58, 260, 261
POLYGYNIA.....	35	brachycarpa	261
TRIGYNIA	35	gracilis.....	260
Heracleum.....	29, 176	parviflora.....	260
lanatum.....	176	Purshiana.....	260
Hesperocnide	73, 273	stipularis.....	260
tenella	274	strigosa	260
Hetercodon	22, 122	subpinnata	260

Hound's-tongue	158	Judas Tree.....	265
Huckleberry	175	JUGLANDEÆ	209
Hydrocotyle.....27, 184		Juglans.....74, 209	
prolifera	184	Californica	209
ranunculoides	184	JULIFLORÆ	268
HYDROPHYLLEÆ	158	JUNCACEÆ	293
HYPERICINEÆ	215	JUNCAGINEÆ	323
Hypericum.....46, 59, 215		Juncus	34, 294
anagalloides	216	Balticus.....	294
concinnum	215	Breweri	294
Hypochæris	59, 115	bufonius	295
glabra	116	compressus.....	294
HYPOGYNÆ	128	effusus	295
ICOSANDRIA	41	falcatus	295
DI - TRI - TETRA - PENTA -		Kelloggii.....	295
GYNIA	45	Leseurii	294
MONOGYNIA	47	patens.....	295
POLYGYNIA.....	46	phæocephalus.....	295
Indian Hemp.....129, 130		robustus	294
INULEÆ	88	tenuis	295
IRIDACEÆ	282	xiphoides.....	295
Iris.....8, 283		Jane-Berry.....	252
Douglasiana	283	Juniper	318
longipetala	283	Juniperus.....78, 328	
macrosiphon.....	283	Californica	329
ISOCARPICÆ	170	Jussiaea	36, 40, 240
ISOËTEÆ	329	repens	240
Isoëtes	329	Knot-grass.	275
Isolepis	319	Koeleria.....13, 306	
carinata.....	319	cristata.....	307
riparia	319	Krynitzkia	19, 156
Iva.....66, 70, 76, 92		ambigua.	156
axillaris	92	Californica.....	157
Jacob's-ladder	163	Chorisiana.....	57
<i>Jaumea</i>	103	leiocarpa	156
		muriculata.....	156

Krynitzkia—oxycarya....	157	Lemna—gibba.....	326
Torreyana.....	156	minor.....	326
LABIATÆ.....	132	trisulca.....	325
Ladies'-Tresses.....	281	Valdiviana.....	325
Lady's-Mantle.....	250	LEMNACEÆ.....	325
Lady's-slipper.....	282	LENTIBULARIA.....	170
Lagophylla.....	67, 69, 100	Lepidium.....	6, 54, 222
congesta.....	101	latipes.....	223
ramosissima.....	101	nitidum.....	223
LAMIALES.....	132	<i>Lepigonum</i>	194
Larkspur.....	233	Leptosyne.....	66, 96
Lastarriæa.....	15, 278	Stillmani.....	96
Chilense.....	278	Lepturus.....	10, 312
Lasthenia.....	64, 105	incurvatus.....	313
Californica.....	106	Lessingia.....	61, 63, 83
glaberrima.....	106	Germanorum.....	83
glabrata.....	106	leptoclada.....	83
Lathyrus.....	58, 265	ramulosa.....	83
littoralis.....	265	Libocedrus.....	73
palustris.....	265	LIGULIFLORÆ.....	115
vestitus.....	265	Ligusticum.....	178
LAURINEÆ.....	230	apiifolium.....	178
Lavatera.....	57, 202	Lilæa.....	71, 324
assurgentiflora.....	230	subulata.....	325
Layia.....	67, 101	LILIACEÆ.....	285
calliglossa.....	102	Lilium.....	33, 289
chrysanthemoides.....	102	pardalinum.....	290
carnosa.....	101	Lily.....	289
elegans.....	101	LIMNANTHÆÆ.....	207
gaillardiioides.....	102	Limnanthes.....	40, 207
heterotricha.....	101	alba.....	208
hieracioides.....	102	Douglasii.....	208
platyglossa.....	102	Limosella.....	51, 148
Leatherwood.....	266	aquatica.....	148
LEGUMINOSÆ.....	253	Linaria.....	53, 153
Lemna.....	6, 325	Canadensis.....	153
		LINEÆ.....	206

Linum	30, 31, 206	Lupinus—nanus	255
Breweri	207	polyphyllus	254
Californicum	207	rivularis	254
congestum	207	trifidus	256
perenne	206	truncatus	256
spergulinum	207	Luzula	34, 296
Liquorice	362	comosa	296
Lip-fern	331	LYCOPODIACEÆ	329
Lippia	53, 140	Lycopus	7, 133
nodiflora	140	lucidus	133
LOASACEÆ	217	LYTHRARIÆ	245
Loco-weed	263	Lythrum	32, 44, 245
Lolium	10, 313	alatum	245
perenne	313	Madia	62, 66, 70, 97
temulentum	313	dissitiflora	97
Lomaria	333	elegans	97
Lonicera	22, 125	filipes	98
involucrata	125	Nuttallii	97
hispidula	125	radiata	97
Loose-strife	245	sativa	97
Lophanthus	50, 137	Madroña	174
urticifolius	137	Maianthemum	17, 284
Lophochlæna	14, 308	bifolium	284
Californica	308	Maiden's-Hair	333
LORANTHACEÆ	266	Malacothrix	59, 118
Lovage	178	Californica	118
Lupine	253	Clevelandi	118
Lupinus	56, 57, 253	obtusa	118
affinis	255	Mallow	203
albicaulis	255	Malva	57, 203
arboreus	254	borealis	203
Chamissonis	254	MALVACEÆ	202
densiflorus	256	MALVALES	202
Douglasii	254	Malvastrum	204
littoralis	255	Thurberi	204
micranthus	256	Manzanita	174
microcarpus	256		

- Maple.....211
 Mare's-tail.....239
 Mariposa.....290
 Marrubium..... 49, 138
 vulgaris.....139
 Marsh-flleabane..... 88
 Marsilæa.....330
 vestita.....330
 Matricaria.....62, 108
 discoidea.....108
 occidentalis.....108
 Mayweed.....107
 Meadow-rue.....237
 Meconopsis.....47, 230
 heterophylla.....230
 Medicago..... 58, 259
 denticulata.....259
 lupulina.....259
 sativa.....259
 Medick.....259
 Megarrhiza.....73, 122
 Californica.....123
 fabacea.....123
 marah.....123
 MELANTHACEÆ..... 29
 Melica.....13, 307
 bomoides.....307
 bulbosa.....307
 Harfordii.....307
 imperfecta.....307
 Melic-grass.....307
 Melilot.....258
 Melilotus.....58, 258
 parviflora.....259
 Mentha.....50, 133
 Canadensis.....133
 piperita.....133
 Mentha—*viridis*.....133
 Menyanthes.....20, 131
 trifoliata.....131
 Mentzelia..... 46, 59, 218
 lævicaulis.....218
 Lindleyi.....218
 Mesembryanthemum..46, 47
 190.
 æquilaterale.....190
 MICRANTHÆ.....296
 Microcala.....16 130
 quadrangularis.....130
 Micromeria..... 50, 135
 Douglasii.....135
 Micropus.....65, 88
 Californicus.....88
 Microseris.....59, 116, 117
 acuminata.....117
 aphantocarpha.....118
 attenuata.....117
 Bigelovii.....118
 Douglasii.....117
 elegans.....118
 Milkweed.....128
 Milkwort.....171
 Mimetanthe..... 52, 149
 pilosa.....149
 Mimulus.....52, 54, 149
 cardinalis.....150
 inconspicuus.....149
 luteus.....150
 moschatus.....150
 nasutus.....149
 Miner's-lettuce.....195
 Mint.....133
 Mistletoe.....267
 Mistletoe (pine).....267

MONADELPHIA.....	56	Muilla	32, 289
DECANDRIA	56	maritima.....	289
OCTANDRIA	56	Mustard.....	225
PENTANDRIA	56	Myosurus.....	31, 49, 236
POLYANDRIA	57	minimus	236
TRIANDRIA.....	56	MYRICACEÆ.....	272
MONANDRIA.....	5	Myrica.....	76, 272
DI-TRI-TETRAGYNIA	5	Californica.....	272
MONOGYNIA.....	5	Myriophyllum.....	74, 239
Monardella.....	50, 134	spicatum.....	239
Breweri	134	MYRTALES	238
Douglasii.....	134		
lanceolata	134	NAIADACEÆ.....	324
undulata.....	134	Naias (Najas).....	71
villosa.....	134	Nardosmia	69, 110
Monkey-flower..	149, 150	palmata	111
MONOCHLAMYDEÆ.....	267	Nasturtium.....	55, 223
MONOCOTYLEDONES.....	280	officinale.....	223
MONGECIA	71	Negundo..	18, 25, 77, 78, 211
DIANDRIA	72	Californicum.....	212
MONANDRIA.....	71	Neillia.....	45, 247
PENT—POLYANDRIA.....	74	opulifolia.....	247
TETRANDRIA.....	73	Nemophila.....	20, 26
TRIANDRIA.....	72	Nettle.....	273
MONOGAMIA.....	70	Nicotiana	21, 154
Monolopia	64, 105	Bigelovii.....	155
gracilens.....	105	Nightshade.....	153
major	105	Nine-bark	247
Monterey-pine	327	Nuphar	238
Montia	8, 196	polysepalum.....	238
fontana.....	197	Nutmeg Pine.....	326
Morning-glory.....	168	Nuttallia	44, 45, 246
Mountain Mahogany.....	250	cerasiformis	246
Mountain Mint.....	133	NYCTAGINEÆ.....	199
Mouse-ear Chickweed	192	NYPHÆACEÆ.....	238
Mousetail	236		
Mudwort.....	148	Oak.....	269

OCTANDRIA.....	36	Panicum.....	10, 296
DI-TRIGYNIA.....	38	PAPAVERACEÆ.....	228
MONOGYNIA.....	36	PAPILIONACEÆ.....	253
TETRAGYNIA.....	38	PARIETALES.....	217
Enanthe... ..	28, 178	PARONYCHIÆ.....	194
California.....	178	Pearly-Everlasting.....	90
Enothera.....	37, 241	Pearlwort.....	193
biennis.....	241	Pectocarya.....	19, 157
California.....	242	pericillata.....	158
Cheiranthifolia... ..	242	Pedicularis.....	51, 143
dentata.....	242	densiflora.....	143
micrantha.....	242	<i>Pellæa</i>	332
ovata.....	242	Pentacæna... ..	14, 24, 27, 194
strigulosa.....	242	ramosissima.....	195
OLEACEÆ.....	131	Pentachæta.....	61, 63, 82
ONAGRACEÆ.....	239	alsinoides.....	83
Onion.....	286	bellidiflora.....	82
OPHIOGLOSSEÆ.....	335	exilis.....	83
Orache.....	198	PENTANDRIA.....	18
Orchard grass.....	306	DIGYNIA.....	25
ORCHIDACEÆ.....	280	MONOGYNIA.....	18
Oreodaphne.....	38, 231	PENTAGYNIA.....	31
California.....	231	POLYGYNIA.....	31
OROBANCHACEÆ.....	142	TRIGYNIA.....	31
Osmorrhiza.....	29, 179	Pentstemon.....	21, 53, 151
brachypoda.....	179	azureus.....	151
nuda.....	179	centranthifolius.....	151
OXALIDEÆ.....	206	Peony.....	232
Oxalis.....	43, 56, 206	Peppergrass.....	222
corniculata.....	206	PRIGYNÆ.....	238
Oregana.....	206	PERSONALES... ..	142
Pæonia... ..	49, 232	<i>Petasites</i>	110
Brownii.....	233	Peucedanum.....	29, 177
Painted-Cup.....	147	caruifolium.....	177
Panax.....	27, 184	dasycarpum.....	177
horridum.....	184	leiocarpum.....	177
		triternatum.....	177

Peucedanum — utricu-	Pinus—insignis.....	327
latum.....	muricata.....	327
Phacelia.....	Sabiniana.....	327
ciliata.....	tuberculata ..	327
circinata.....	PIPERALES.....	279
divaricata.....	Plagiobothrys.....	19, 157
distanis.....	canescens.....	157
hispida.....	nothofulvus.....	157
malvæfolia.....	PLANTAGINEÆ.....	140
tanacetifolia.....	Plantago.....	16, 141
Phalaris.....	lanceolata.....	141
amethystina.....	major.....	141
arundinacea.....	maritima.....	141
Canariensis.....	Patagonica.....	141
intermedia.....	Plantain.....	141
PHANEROGAMÆ.....	PLATANÆ.....	272
PHILADELPHÆÆ.....	Platanus.....	76, 273
Phleum.....	racemosa.....	273
alpinum.....	Platystemon.....	48, 228
pratense.....	Californicus.....	229
Phoradendron...73, 77, 267	Platystigma ..	48, 229
flavescens.....	Californicum.....	229
Photinia.....	lineare.....	229
arbutifolia....	Plectritis.....	8, 124
.....252	congesta.....	124
Phragmites.....	macrocera.....	124
communis.....	Pluchea.....	65, 88
Pickeringia.....	camphorata.....	88
montana.....	Plum.....	246
.....253	PLUMBAGINEÆ.....	170
Pigweed.....	Poa.....	13, 309
Pilularia.....	annua.....	309
Americana.....	Californica.....	309
.....330	distanis.....	309
Pimpernel.....	Douglasii.....	310
.....172	pratensis.....	309
Pimpinella.....	scabrella.....	309
apiodora.....		
.....181		
Pin-clover.....		
.....205		
Pine.....		
.....326		
Pinus.....		
.....72, 326		

Poa—tenuifolia.....	309	Pondweed.....	323
trivialis.....	309	Poplar.....	217
Pogogyne.....	50, 135	Populus.....	79, 217
Douglasii.....	135	trichocarpa.....	217
parviflora.....	135	POTAMEÆ.....	323
serpylloides.....	135	Potamogeton.....	18, 323
POLEMONIACEÆ.....	162	lucens.....	324
POLEMONIALES.....	153	natans.....	324
Polemonium.....	21, 163	pauciflorus.....	324
carneum.....	163	pectinatus.....	324
POLYADELPHIA.....	59	Potentilla.....	46, 249
POLYANDRIA.....	47	anserina.....	249
DI - TRI - TETRA - PENTA -		glandulosa.....	249
POLYGYNIA.....	48	Portulaca, 35, 36, 42, 43, 45	
MONOGYNIA.....	47	195.	
POLYCARPICÆ.....	230, 322	oleracea.....	195
Polygala.	56, 57, 210	PORTULACACEÆ.....	195
cucullata.....	210	Prickly Poppy.....	230
POLYGALACEÆ.....	210	PRIMULACEÆ.....	171
POLYGONACEÆ.....	274	PRIMULALES.....	170
Polygonum.....	24, 38, 275	Prosartes.....	33, 284
aviculare.....	275	Hookeri.....	285
Californicum.....	275	Menziesii.....	285
convolvulus.....	276	Prunella.....	50, 138
Muhlenberghii.....	276	vulgaris.....	138
nodosum.....	276	Prunus.....	246
paronychia.....	275	demissa.....	246
POLYPETALÆ.....	175	emarginata.....	246
POLYPODIACEÆ.....	331	ilicifolia.....	246
Polypodium.....	332	Pseudotsuga.....	327
Scouleri.....	332	Douglasii.....	327
vulgaris.....	332	Psilocarphus.....	67, 70, 89
Polypody.....	332	Oreganus.....	89
Polypogon.....	11, 300	tenellus.....	89
littoralis.....	300	Psoralea.....	57, 58, 261
Monspeliensis.....	300	macrostachya.....	262
POMACEÆ.....	252	orbicularis.....	261

Psoralea—physodes	262	Ranunculus	49, 235
strobilina	262	aquatilis	236
Ptelea, 17, 24, 73, 74, 77, 78		Bloomeri	235
208.		Californicus	235
angustifolia	208	flammula	236
Pteris	332	hebecarpus	235
aquilina	333	Lobbii	236
Pterostegia	34, 278	maximus	235
drymarioides	278	muricatus	235
Pugiopappus	66, 96	pusillus	236
calliopsideus	96	Raphanus	54, 221
Purslane	195	sativus	222
Pycnanthemum	50, 133	Raspberry	247
Californicum	134	Rattleweed	263
Pyrola	39, 173	Redwood	328
aphylla	173	Reed	306
PYROLACEÆ	172	RHAMNACEÆ	212
Quaking-grass	310	Rhamnus	23, 78, 212
Quercus	76, 269	Californica	212
agrifolia	270	Crocea	212
chrysolepis	269	Purshiana	212
densiflora	270	RHINANTHÆÆ	143
Douglasii	269	RHIZANTHÆÆ	267
dumosa	269	RHIZOCARPÆÆ	330
Kelloggii	270	Rhododendron	40, 173
lobata	269	occidentale	173
Wislizenii	270	RHODORACEÆ	173
Quillwort	329	Rhus	30, 74, 78, 82, 209
Quinine-Tree	185	aromatica	209
Radish	221	diversiloba	209
Rafinesquia	59, 115	Ribes	24, 30, 188
Californica	115	aureum	188
Ragweed	92	divaricatum	188
RANALES	237	Menziesii	188
RANUNCULACEÆ	232	sanguineum	188
		Rigiopappus	64, 106
		leptocladus—see <i>Errata</i> .	

Rock-brake.....	332	Salix—lasiolepis.....	216
Rock-cress.....	226	longifolia.....	216
Rock-rose.....	217	Salvia.....	7, 136
Romanzoffia.....	20, 159	carduacea.....	136
Sitchensis.....	159	Columbariæ.....	137
Rosa.....	46, 251	SALVINIACEÆ.....	330
Californica.....	251	Sambucus.....	30, 124
gymnocarpa.....	251	glauca.....	124
Rose.....	251	Samolus.....	22, 41, 172
ROSACEÆ.....	251	Valerandi.....	172
ROSIFLORÆ.....	245	Samphire.....	199
RUBIACEÆ.....	126	Sand-spurrey.....	194
RUBIALES.....	133	Sandwort.....	193
Rubus.....	46, 247	Sanicle.....	182
leucodermis.....	248	Sanicula.....	27, 182
Nutkanus.....	247	Sanguisorba.....	18
spectabilis.....	247	SANGUISORBACEÆ.....	249
ursinus.....	248	SAPINDACEÆ.....	210
Rumex.....	34, 79, 274	SAPINDALES.....	210
acetosella.....	275	SAURUREÆ.....	279
conglomeratus.....	275	Saxifraga.....	41, 186
crispus.....	274	Virginica... ..	186
obtusifolius.....	275	SAXIFRAGACEÆ.....	186
salicifolius.....	274	Saxifrage.....	186
RUTACEÆ.....	208	Scirpus.....	6, 15, 69, 320
Sage.....	136	acicularis.....	320
Sage-brush.....	109	lacustris.....	320
Sagina.....	31, 42, 193	maritimus.....	320
occidentalis.....	193	Olneyi.....	320
SALICINEÆ.....	216	palustris.....	320
Salicornia.....	5, 199	Tatora.....	320
ambigua.....	199	sylvaticus.....	320
Salix.....	76, 77, 216	Scoliopus.....	8, 292
Coulteri.....	217	Bigelovii.....	292
lævigata.....	216	Scorzonella.....	116
lasiandra.....	216	paludosa.....	116
		procera.....	116

Scorzonella—sylvatica ..	116	Shepherd's-purse	223
Scouring-rush	335	Shield-fern.....	334
Scrophularia	53, 153	Shooting-star	171
<i>Californica</i>	153	Sida	57, 204
SCROPHULARIÆ	143	<i>hederacea</i>	204
Scutellaria.....	50, 138	Sidalcea.....	57, 203
<i>Californica</i>	138	<i>diploscypha</i>	203
<i>tuberosa</i>	138	<i>humilis</i>	203
Sea-blite.....	199	<i>malachroides</i>	203
Sea-lavender	171	<i>malvaeflora</i>	203
Sea-Rocket.....	224	Silene.....	41, 191
Sedge	316	<i>antirrhina</i>	191
Sedum.....	43, 189	<i>Californica</i>	191
<i>spathulifolium</i>	189	<i>Gallica</i>	191
<i>stenopetalum</i>	189	<i>laciniata</i>	191
Selaginella.....	329	<i>verecunda</i>	192
<i>rupestris</i>	329	SILENEÆ	191
SELAGINELLEÆ	329	Silybum	114
Self-heal	138	<i>Marianum</i>	114
Selinum.....	28, 178	Sisymbrium.....	56, 224
<i>Pacificum</i>	178	<i>acutangulum</i>	224
Senebiera.....	54, 222	<i>officinale</i>	224
<i>didyma</i>	222	<i>reflexum</i>	224
Senecio.....	61, 65, 111	Sisyrinchium.....	56, 282
<i>aronicoides</i>	111	<i>bellum</i>	283
<i>Douglasii</i>	112	<i>Californicum</i>	283
<i>eurycephalus</i>	111	Sium	28, 179
<i>hydrophilus</i>	111	<i>cicutæfolium</i>	179
<i>vulgaris</i>	112	Skullcap	138
SENECIONIDÆ	110	SMILACEÆ	284
Sequoia.....	74, 328	Smilacina	34, 284
<i>gigantea</i>	328	<i>amplexicaulis</i>	284
<i>sempervirens</i>	328	<i>sessilifolia</i>	284
SERPENTARIÆ	267	Snapdragon	152
Service-berry.....	252	Sneezeweed	107
Setaria	10	Snowberry	125
Sheep-sorrel.....	274, 275	Soap-root.....	286

SOLANACEÆ	153	Spiranthes—porrifolia . . .	281
Solanum	22, 153	Romanzoffiana	281
nigrum	154	Spleenwort	334
umbelliferum	154	Spotted Thistle	114
Solidago	63, 85	Spurge	201
Californica	85	Squirrel-tail	314
occidentalis	85	St. John's-wort	215
sempervirens	85	Stachys	50, 139
Soliva	69, 109	ajugoides	139
sessilis	109	albans	139
Sonchus	60, 120	bullata	140
asper	120	Chamissonis	140
oleraceus	120	pycnantha	139
Sow-thistle	120	Star-flower	171
SPADICIFLORÆ	321	Star-thistle	114
Sparganium	72, 322	Statice	31, 171
eurycarpum	322	limonium	171
Spartina	11, 202	Stellaria	42, 192
stricta	203	media	192
Specularia	22, 121	nitens	193
biflora	121	Stephanomeria	59, 115
Speedwell	148	virgata	115
Spergula	31, 43, 194	Stipa	12, 301
arvensis	194	eminens	302
Spergularia	42, 194	setigera	302
macrotheca	194	viridula	302
media	194	STIPULATÆ	335
Sphacele	50, 136	Stone-crop	189
calycina	136	Stork's-bill	205
<i>Sphaeralcea</i>	57, 204	Strawberry	248
Spice-bush	252	Streptanthus	55, 226
Spikenard	184	glandulosus	226
Spindle-tree	214	niger	226
Spiræa	45, 46, 247	peramœnus	226
discolor	247	Stylocline	65, 67, 70
SPIRÆEÆ	247	Suæda	25, 199
Spiranthes	71, 281	Californica	199

Sunflower.....	95	TETRADYNAMIA.....	54
Sweet Cicely.....	179	SILICULOSA.....	54
Sweet Vernal-grass.....	299	SILIQUOSA.....	54
Sycamore.....	273	TETRANDRIA.....	16
Symphoricarpus.....	22, 125	DIGYNIA.....	16
mollis.....	125	MONOGYNIA.....	16
racemosus.....	125	Thalictrum.....	49, 79, 237
SYNANDRÆ.....	80	Fendleri.....	237
SYNGENESIA.....	59	Thermopsis.....	39, 253
ÆQUALIS.....	59	Californica.....	253
FRUSTRANEA.....	68	Thistle.....	112
NECESSARIA.....	68	Thorn-apple.....	154
SUPERFLUA.....	62	Thrift.....	170
Synthyris.....	7, 148	THYMELACÆ.....	266
rotundifolia.....	148	Thysanocarpus.....	54, 222
Syrmatium.....	261	curvipes.....	222
cytisoides.....	261	laciniatus.....	222
glabrum.....	261	pusillus.....	222
tomentosum.....	261	Tiarella.....	41, 187
Tanacetum.....	61, 109	unifoliata.....	187
camphoratum.....	109	Tillæa.....	14, 18, 31, 190
Tansy.....	109	angustifolia.....	190
Taraxacum.....	60, 120	minima.....	190
officinale.....	120	Timothy.....	297
Tar-weed.....	97, 98	Toad-Flax.....	153
Tassel-tree.....	185	Tobacco.....	154
TAXINÆ.....	326	Torreyia.....	79, 326
Teasel.....	123	Californica.....	326
Tellima.....	41, 186	Toyon.....	252
affinis.....	187	Tree-Mallow.....	202
Bolanderi.....	187	TRIANDRIA..	8
grandiflora.....	187	DIGYNIA.....	9
heterophylla.....	187	MONOGYNIA.....	8
TEREBINTHACÆ.....	208	TRIGYNIA.....	14
TEREBINTHALES.....	208, 209	Tricerastes.....	15, 218
TETRACYCLICÆ.....	80	glomerata.....	218
		Trichostema.....	51, 132

Trichostema—lanceola-	Troximon	60, 119
tum	apargioides	119
TRICOCCEÆ	grandiflorum	119
Trientalis	heterophyllum	120
latifolia	humile	119
Trifolium	laciniatum	119
barbigerum	TUBULIFLORÆ	81
bifidum	<i>Turritis glabra</i>	227
ciliatum	Twin-berry	125
depauperatum	Typha	72, 322
fucatum	latifolia	322
gracilentum	TYPHACEÆ	321
involueratum	UMBELLALES	175
Macræi	UMBELLIFERÆ	175
microcephalum	<i>Umbellularia</i>	231
microdon	URTICALES	272
pauciflorum	Utica	73, 77, 273
repens	holosericea	273
tridentatum	Lyallii	273
Triglochin	urens	273
maritimum	Utricularia	7, 170
Trillium	vulgaris	170
ovatum	VACCINEÆ	175
sessile	Vaccinium	37, 175
Trisetum	ovatum	175
barbatum	VALERIANACEÆ	124
canescens	Vancouveria	31, 231
Triteleia	hexandra	231
capitata	Venus Looking-glass	121
ixioides	VERBASCEÆ	153
lactea	Verbena	53, 140
laxa	officinalis	140
peduncularis	VERBENACEÆ	140
Triticum	Veronica	6, 148
repens	Americana	148
Tropidocarpum		
gracile		

Veronica—peregrina	148	Wild-Ginger	268
VERONICÆ	147	Wild-Pea	265
Vetch	264	Wild-Rye	315
Vicia	58, 264	Willow	216
Americana	264	Willow-herb	240
exigua	264	Winter cress	224
gigantea	264	Wintergreen	173
sativa	264	Wood-sorrel	206
Viola	23, 220	Wood-rush	296
aurea	220	Woodwardia	333
canina	220	radicans	334
lobata	221	Wormwood	109
ocellata	220	<i>Wulfenia</i>	148
odorata	220	Wyethia	66, 94
pedunculata	220	angustifolia	95
sarmentosa	221	glabra	94
VIOLACEÆ	219	helenioides	94
Virgin's Bower	237	Xanthium	93
Vitis	23, 78, 215	spinosum	93
Californica	215	strumarium	93
Wake-robin	285	Xerophyllum	293
Waldmeister	127	tenax	293
Wall-flower	226	Yarrow	108
Walnut	209	Yedra	209
Water-cress	323	Yellow-dock	275
Water-fern	331	Yerba buena	135
Water-hemlock	180	Yerba santa	158
Water-horehound	133	Yerba mansa	279
Water-lily	238	Zauschneria	240
Water-milfoil	239	Californica	240
Water-parsnip	179	Zostera	71, 325
Water-pennywort	184	marina	325
Water-plantain	323	Zygadene	292
Wax-Myrtle	272	Zygadenus	34, 292
Wheat	314	Fremontii	292
Whipplea	30, 42, 189	venenosus	292
modesta	189		

APPENDIX.

NOTE.—In this work, in order to conform with most American treatises on botany, the word *loculus* is replaced by the less exact word *cell*.

- ⊙. Annual.
- ⊙⊙. Biennial.
- ∩. Perennial.
- h. Shrub or tree.
- ♂. Male.
- ♀. Female.
- ♂. Hermaphrodite, *i. e.*, both sexes in the same flower.
- ∞. Many, or an indefinite number.
- Acaulescent*. Stemless.
- Accrescent*. Increasing with age.
- Acerose*. Needle-shaped.
- Aculeate*. Having sharp points or prickles.
- Acuminate*. Tapering to a point.
- Æqualis*. All flowers hermaphrodite.
- Acute*. Sharp-pointed.
- Adnate*. United by the surface.
- Alcene*. A dry, hard, indehiscent, 1-celled, 1-seeded fruit.
- Alveolate*. Pitted like a thimble.
- Ament, Amentum*. A unisexual spike with scaly bracts.
- Amplexicaul*. Clasping the stem.
- Anceps, Ancipital*. Two-edged.
- Andræcium*. Stamens and corolla.
- Androgynous*. Having both male and female flowers.
- Angiosperm, Angiospermæ*. Bearing seeds in a closed pericarp.
- Anisocarpæ*. Number of Carpels less than lobes of Corolla.
- Annular*. Ring-like.
- Anther*. The part of the stamen bearing pollen.
- Aphyllous*. Without leaves.

- Apetalous.* Without petals.
- Apocarpous.* Carpels (*Carpidia*) distinct.
- Apophyses.* E enlargements, structurally united.
- Appressed.* Pressed close.
- Arachmoid.* Cobweb-like.
- Arillus.* An expanded appendage attached to the seed.
- Aristate.* Having an awn.
- Articulate.* Jointed.
- Ascending.* Rising, but not erect.
- Aspergilliform.* Brush-like.
- Auriculate.* Having an ear-like lobe at the base.
- Awn.* A bristle-like appendage.
- Axil.* The angle formed by a leaf or branch with the stem.
- Baccate.* Berry-like.
- Barbellate.* Having minute awns with reflexed points.
- Basifixed.* Attached by the lower end.
- Bifid.* Two-cleft.
- Bipinnate.* Twice pinnate.
- Bract.* A leaf, or modification of one, subtending a flower or cluster.
- Bracteolate.* Having secondary bracts upon the pedicel.
- Caducous.* Falling very early.
- Cæspitose, Cespitose.* Growing in tufts.
- Calamus.* A stuffed scape.
- Calcarate.* Spurred.
- Calycine.* Calyx-like.
- Calyculate.* Calyx-like.
- Campanulate.* Bell-shaped.
- Canaliculate.* Fluted.
- Canescent.* Hoary with a gray pubescence.
- Capillary.* Very slender and hair-like.
- Capitate.* Collected into a head.
- Capitulate.* Diminutive of capitate.
- Carina.* Keel.
- Carpidia.* A simple pistil, or one of the several parts of a compound pistil.

Carpophore. A prolongation of the axis between the carpels.

Caryopsis. Seed inseparably united to the wall of the ovary.

Caudate. Tailed.

Caulescent. Having a manifest stem.

Cauline. Belonging to the stem.

Centrospermæ. Seeds affixed to the centre of a 1-celled ovary.

Chlorophyll. The green coloring matter of plants.

-chotomous. Forked.

Ciliate. Fringed with hairs.

Circinate. Coiled from the tip.

Circumscissile. Opening by a transverse circular line.

Cirrhate. Tendril-bearing.

Clavate. Club-shaped.

Cleistogamic. Having fertile but undeveloped flowers.

Coccus. The indehiscent segment of a compound ovary.

Cælospermæ. Internal surface of Endosperm concave.

Commissure. The surface by which two carpels adhere.

Complicate. Folded together.

Conduplicate. Doubled together lengthwise.

Connate. United in one.

Connective. The part of the filament which connects the anther cells.

Contorted. Twisted.

Convolute. Rolled together from one edge.

Cordate. Heart-shaped.

Coriaceous. Leathery.

Corniform. Horn-shaped.

Corolline. Like a corolla.

Corona. Crown.

Corymb. A depressed raceme.

Costa. A rib.

Cotyledon. The seed leaf or lobe of the embryo.

Cremocarp. Fruit of Umbelliferæ.

Crenate. Having rounded teeth.

Crenulate. Finely crenate.

Cryptogamæ. Fructifying without stamens and pistils.

Cucullate. Like a cowl or hood.

Cuneate. Wedge-shaped enlarging upward.

Cuneiform. Same as cuneate.

Cupuli. A cup-shaped involucre inclosing a nut as in the acorn.

Cusp. A sharp rigid point.

Cyathiform. Cup-shaped.

Cyme. A broad and flattish inflorescence, flowering from the centre outward.

Decandria Ten stamens.

Declined. Bent or curved downward.

Decomound. Repeatedly compound or divided.

Decumbent. Reclining at base, the summit ascending.

Decurrent. Running down the stem.

Decussate. In pairs or threes alternating at right angles.

Deflexed. Bent or turned down abruptly.

Dehiscent. Opening regularly by valves, slits, etc.

Deltoid. Broadly triangular like the Greek letter Delta.

Dentate. Toothed.

Denticulate. Minutely toothed.

Diadelphia. Stamens in two sets or clusters.

Diandria Two stamens.

Dichotomous. Forking regularly by pairs.

Diclinic. Of separate sexes; unisexual.

Dicotyledones. Having an embryo with two cotyledons.

Didynamia. Having two long and two short stamens.

Diffuse. Widely and loosely spreading or branched.

Digitate. Fingered; spreading like the fingers.

Digynia. Two pistils.

Dimidiate. Halved, as though one-half were wanting.

Dimorphous. Occurring in two forms.

Diœcia. Male and female flowers on separate plants.

Distichous. Two ranked.

Divaricate. Widely diverging.

Dodecandria. Having twelve stamens.

- Dorsal*. Relating to the *dorsum* or back.
- Drupe*. A fleshy or pulpy fruit with the seed inclosed in a hard or stony casing.
- Ebracteate*. Without bracts.
- Echinate*. Beset with hooked prickles.
- Eliater*. Coiled elastic thread.
- Emarginate*. Notched at the extremity.
- Endosperm*. Layer of substance in the seed usually enveloping the germ.
- Enneandria*. Nine stamens.
- Epicarp*. Outer layer of the pericarp.
- Epigynæ*. Above the ovary.
- Equitant*. Astride, as in leaves of Iris.
- Exocarp*. The outer portion of a pericarp.
- Exserted*. Projecting beyond an envelope.
- Extrorse*. Directed outward.
- Fascicle*. A close bundle or cluster.
- Filament*. The part of the stamen which supports the anther.
- Filiform*. Thread-shaped.
- Fimbriate*. Fringed with narrow cuttings.
- Floret*. A small flower; one of a head.
- Foliate*. Having leaves, leafy.
- Foliolate*. Having leaflets or small leaves.
- Follicle*. A pod formed from a simple pistil dehiscing only along the ventral suture.
- Foveolate*. Marked by minute pits.
- Frond*. The leaf of ferns, liverworts or of Lemna.
- Frustranea*. Florets of the disk hermaphrodite, of the ray neutral.
- Frutescent*. Shrubby.
- Funicle*. The stalk of an ovule or seed.
- Furfuraceous*. Scurfy.
- Galbulus*. An indehiscent cone, becoming berry-like.
- Galeate*. Formed like a helmet.
- Gamopetalæ*. Petals more or less united.

Gamosepalous. With more or less united or coalescent sepals.

Geminate. Twinned; in pairs.

Geniculate. Bent like the knee.

Gibbous. Swelling out at one side.

Glabrous. Smooth; without hairs or roughness.

Glaucous. Bluish-hoary; covered with a fine whitish bloom.

Glochidiate. Barbed like a fish-hook,

Glomerule. A compact somewhat capitate cyme.

Glume. The chaff-like bracts subtending the spikelets.

-gonal. Relating to the angles.

Gymnospermia. Plants having naked seed.

Gynandria. Having the stamens adnate to or borne upon the pistil.

Gynobase. A short thick prolongation of the axis upon which the pistil rests.

Gynœcium. The pistil or aggregate pistils of a flower.

Gynophore. A stem bearing the ovary.

Hastate. Triangular or arrow-shaped.

Heptandria. Having seven stamens.

Hermaphrodite. Both sexes in the same flower.

Heterosporous. Bearing more than one kind of spores.

Hexandria. Having six stamens.

Hexagynia. Having six pistils.

Hirsute. Having rather coarse or stiff hairs.

Hispid. Beset with rigid or bristly hairs.

Hyaline. Transparent or partially so.

Hypogynous. At the base of, or below the pistil.

Icosandria. Having more than twelve stamens inserted on the calyx.

Imbricate. Overlapping

Imparipinnate. Pinnate, with an odd terminal leaflet.

Incised. Irregularly, sharply and deeply cut.

Incrassate. Thickened.

Incumbent. Lying upon.

- Induplicate.* With margins folded inward.
- Indusium.* In ferns, the scale-like cover of the fruit spot.
- Inferior.* Lower; below the corolla.
- Inflexed.* Bent or turned abruptly inward.
- Inflorescence.* The flowering portion of a plant.
- Introrse.* Turned inward toward the axis.
- Involucre.* A circle or circles of scales, bracts or leaves, distinct or united, surrounding a flower or flower-cluster.
- Involute.* Rolled inward.
- Labellum.* A lip, as in orchids.
- Labiate.* Lipped; having an irregular calyx or corolla unequally divided into two lips.
- Laciniate.* Cut into narrow slender teeth or lobes.
- Lactescent.* Yielding milky juice.
- Lamina.* The blade or dilated part of a leaf.
- Lanceolate.* Shaped like a lance-head.
- Legume.* A 1-celled capsule opening by two valves, as the pea.
- Lenticular.* Lens, or lentil-shaped.
- Lignescens.* Woody; becoming woody.
- Ligule.* A tongue or strap-shaped body, as the corolla of ray-flowers in *Compositæ*.
- Lobate.* Divided into lobes.
- Loculicidal.* Dehiscence of the cells of a capsule through the dorsal suture.
- Lomentum.* *Loment.* A legume jointed, and usually constricted between the seeds.
- Macro.* Large or long.
- Marcenscent.* Withering and persistent.
- Marginate.* Furnished with a border.
- Membranaceous.* Thin and translucent.
- Meri-carp.* Division of schizocarp.
- merous.* In parts.
- Micro.* Small.
- Monadelphia.* Stamens united by their filaments into one set.

- Monandria*. One stamen.
- Moniliform*. Like a string of beads.
- Monocotyledones*. Embryo with one cotyledon.
- Monœcia*. Stamens and pistils in separate flowers on the same plant.
- Monogamia*. A syngenetic flower not belonging to a head.
- Monogynia*. Having one pistil.
- Mucro*. A short and small abrupt tip.
- Mucronate*. Having a mucro.
- Mucronulate*. Diminutive of mucronate.
- Multifid*. Cleft into many lobes and segments.
- Muricate*. Rough, with short hard points.
- Navicular*. Boat-shaped.
- Nectaria, Nectary*. Organs which produce a sweet secretion within a flower.
- Necessaria*. Florets of the disk male; of the ray female; or hermaphrodite.
- Nervose*. Having veins or ribs.
- Neutral*. Neither male nor female.
- Obcompressed*. Flattened dorsally.
- Obconic*. Like an inverted cone.
- Obcordate*. Inverted cordate.
- Obovate*. Inverted ovate.
- Obtuse*. Blunt.
- Ochroleucous*. Yellowish white.
- Octandria*. Eight stamens.
- Orthospermæ*. Having the endosperm straight (flat).
- Osseous*. Bony.
- Ovoid*. Egg-shaped.
- Palea, Palet*. Chaff, or a chaffy bract; in grasses the two inner bracts of the flower.
- Palmate*. Compound, with the leaflets radiating.
- Panicle*. A compound raceme.
- Papilionaceous*. Butterfly-like. The peculiar corolla of the Leguminosæ.
- Papillate*. Beset with minute thick projections.

Pappus. The hairs, bristles or scales crowning the akenes in Compositæ.

Parietal. Relating to or upon the wall of a cavity.

Paripinnate. Evenly pinnate; without the odd terminal leaflet.

Pedicel. The footstalk of a flower.

Peduncle. A general or primary flower-stalk.

Peltate. Shield-shaped.

Pendulous. Drooping, or hanging nearly inverted.

Penicillate. Resembling a brush of fine hairs.

Pentandria. Having five stamens.

Pentagynia. Having five pistils.

Perianth. The floral envelopes—*i. e.*, the calyx and corolla, so far as present.

Pericarp. The seed-vessel or ripened ovary.

Perigonium. A simple floral envelope.

Perigynium. The sac-like envelope, or the bristles or scales which in the Cyperaceæ represent the perianth.

Personate. Closed, as in Labiate flowers with prominent palates.

Petal. One of the parts of a polypetalous or nearly divided corolla.

Petiole. The footstalk of a leaf.

Petaloid. Petal-like.

Phalanges. Stamens united in sets by their filaments.

Phanerogamæ. Plants which fructify by means of stamens and pistils.

-phyllous. Leafy, leaved.

Pilose. Hairy, with soft distinct hairs.

Pinnæ. Primary divisions of a compoundly pinnate leaf.

Pinnatifid. Pinnately cleft into opposite nearly equal segments.

Pinnule. A secondary pinna—*i. e.*, one of the pinnate divisions of a pinna.

Pistil. The female organ of a phanerogam consisting of the ovary with its styles and stigmas.

Placenta. That part of the ovary or fruit which bears the ovules and seeds.

Plicate. Folded into plaits like a fan.

Plumose. Plume-like; having fine hairs on each side, like a feather.

Polyadelphia. Having stamens united by the filaments into many sets.

Polyandria. Having many stamens.

Polygamia. Having both hermaphrodite and unisexual flowers.

Polygynia. Having many pistils.

Pollinia. Concrete masses of pollen.

Pomum, Pome. A fleshy fruit like the apple, inclosing several leathery or bony carpels.

Procumbent. Lying upon the ground.

Pruinose. Covered with a minute bloom or powder.

Pseudocarp. Fruit including other organs in addition to the ovary.

Puberulent. Very minutely pubescent.

Pubescent. Covered with hairs usually short and soft.

Pungent. Terminating in a rigid and stout point or prickle.

Pyrena. The stone of a drupe.

Pyriform. Pear-shaped.

-quetrous. Cornered.

Quinate. In fives.

Raceme. A form of inflorescence, with pedicellate flowers developing from below upward upon a simple prolonged axis.

Rachis. The axis of a spike, etc.

Radicle. The part of the embryo below the cotyledon.

Ray. The marginal flowers in compositæ.

Receptacle. The more or less expanded surface forming a support for the organs of a flower or a head of flowers.

Reflexed. Bent abruptly down or backward.

Reniform. Kidney-shaped.

Replum. The frame-like placenta left by the falling away of the valves in Cruciferæ.

Retrorse. Turned back or downward.

Revolute. With the margin or apex rolled backward.

Rhizome. An underground, somewhat horizontal, rooting stem.

Ringent. Gaping; applied to labiate flowers with open throats.

Rostrate. Beaked.

Rotate. Wheel-shaped.

Rugose. Wrinkled; ridged.

Runcinate. Deeply-toothed or lobed, with the segments directed backwards.

Saccate. Sac-shaped.

Samara. An indehiscent, membraneously-winged fruit, as in the ash and maple.

Sagittate. Shaped like an arrow-head

Scabrous. Rough to the touch, with minute rigid points.

Scape. A naked peduncle rising from the ground.

Scapigerous. Producing scapes.

Scarious. Thin, dry and membranaceous.

Schizocarp. An ovary which divides into its component parts, each part usually remaining closed over its seed.

Secund. One-sided.

Septicidal. Opening through the lines of junction of the carpels.

Septifragal. Breaking away from the partitions in dehiscence.

Serrate. Having teeth directed forward, like those of a saw.

Serrulate. Finely serrate.

Sessile. Without footstalk.

Setaceous. Bristle-like.

Siliculosa. Short cruciferous pod, as *Capsella*.

Siliquosa. Long cruciferous pod, as in mustard.

Sinuate. With a strongly wavy margin.

Sinus. An indentation, either angular or rounded, separating parts.

Sordid. Rusty or dirty-colored.

Sorus, Sori. A cluster of sporangia.

Spadix. A spike with a thickened, fleshy rachis, usually subtended by a spathe.

Spathe. A sheathing bract.

Spathulate. Narrowly attenuate downward from an abruptly rounded summit.

Spiculate. Like a spike.

Spike. Resembling a raceme, but the flowers sessile, or nearly so.

Spinescent. Ending in a spine or spiny point.

Spinulose. Having diminutive spines.

Sporangia. Cases which contain spores.

Spores. The minute bodies in cryptogams, which answer somewhat to the seeds of other plants.

Squamose. Furnished with scales.

Squarrose. Irregularly spreading.

Stipitate. Borne upon a stipe.

Stipule. An appendage to the base of a petiole.

Stamen. The male organ of flowering plants, consisting of an anther borne usually upon a filament and containing the pollen.

Staminodia. A sterile or undeveloped stamen.

Strobilaceous. Inflorescence formed of imbricated scales, as in the Coniferæ.

Subulate. Awl-shaped.

Suffrutescent. Slightly shrubby; woody at base.

Sulcate. Grooved or furrowed.

Suture. A line of union.

Synandræ. Having united anthers.

Syncarpous. Having united carpidia.

Syngenesia. United anthers.

Tendril. A thread-like production in climbers.

Terete. Cylindrical, or nearly so.

Ternate. In threes.

Testa. The outer seed-coat.

Tetradynamia. Having four long and two short stamens;

Cruciferae.

Tetragynia. Having four pistils.

Tetrandria. Having four anthers.

Thalamus. End of peduncle.

Thallus. A cellular expansion taking the place of stem and foliage.

Thyrse. A contracted or close ovate cyme.

Tomentose. Pubescent, with matted wool.

Triandria. Having three stamens.

Trigynia. Having three pistils.

Triquetrous. Triangular, with the sides concave or channeled.

Truncate. Ending abruptly as if cut off transversely.

Tumid. Swelled.

Turbinate. Top-shaped.

Turgid. Inflated.

Umbel. An umbrella-shaped inflorescence, the pedicels radiating from the summit of the common peduncle.

Umbilicate. Pitted in the centre.

Uncinate. Hooked at the extremity.

Undulate. Wavy.

Unguiculate. Narrowed below into a claw.

Urceolate. Cylindrical or ovoid, but contracted at or below the open orifice.

Utricle. A small, bladdery, usually 1-seeded, pericarp.

Vallecula. Groove between the ribs of umbelliferous fruits.

Valvate. Opening by valves.

Ventral. Belonging to the anterior or inner face of a carpel; the opposite of dorsal.

Ventricose. Swelling unequally or on one side.

Versatile. Swinging; turning freely on its support.

Verticillate. Arranged in whorls.

Vexillum. The large upper petal of a papilionaceous flower.

Villos. Bearing long and soft straight, or straightish, hairs.

Virgate. Slender, straight and erect.

Vittæ. The longitudinal oil-tubes in the pericarp of most umbelliferæ.

Whorl. An arrangement of leaves, flowers, etc., around the stem.



ERRATA.

- Page 106. After generic description of Rigiopappus read
"R. leptocladus.—Niles. Spring."
- Page 163. For "Gillia" read "Gilia."
- Page 163. For "androscaea" read "androsacea."
- Page 171. 6th line from bottom, for "decusate" read
"decussate."
- Page 172. 9th line from bottom, for "L" read "S."
- Page 215. 7th line from top, after "h." insert "—
Niles. Summer."
- Page 262. 10th line from bottom, omit comma between
"late" and "dehiscent."
- Page 275. 12th line from top, for "7" read "5."
- Page 283. 4th line from top, for "Californicus" read
"Californicum."
- Page 299. 11th line from top, after "not" insert "at."
- Page 311. 1st line, for "tenela" read "tenella."
- Page 330. For "Pillularia" read "Pilularia."

9#4618

U. C. BERKELEY LIBRARIES



058353551



