


Kelly (H.A.)

ABSTRACT OF A PAPER ON
URINALYSIS IN GYNÆCOLOGY.

BY
HOWARD A. KELLY, M.D.



REPRINT FROM
American Medico-Surgical Bulletin,
October, 1893.

TO CONTRIBUTORS.

Contributions of Original Thought and Experiences, on Medical and Surgical Topics, are desired by the AMERICAN MEDICO-SURGICAL BULLETIN on the following conditions:

1.—Authors of Scientific Papers or Clinical Reports accepted by us will receive—according to their own preference expressed *with each communication*, either:

a.—A number of REPRINTS of their article in neat pamphlet form (pocket size); or,

b.—Instead of the above, an Equivalent value therefor in CASH.

2.—All contributions are received only on the express understanding:

a.—That they have not been printed anywhere, nor communicated to any other journal.

b.—That, if they have been read anywhere to an audience, this fact be stated in full detail by a note on the manuscript.

Contributors will serve their own interest by heeding the following suggestions:

Write Concisely and Clearly.

What we desire to print, and what professional men like to read, is information, not verbiage. An article will stand better chances of acceptance, and of being widely read and copied and discussed, the fewer its words are, in proportion to the facts or ideas it embodies. Of course, a thought too thinly clad must suffer. Use, therefore cheerfully, as many words as appear needed to convey your meaning,—but no more.

Aim at Fact.

It is not to be expected that every Medical or Surgical paper should be a mere array of statistical data, hospital records, tabulated figures, or graphic summaries; some room must be allowed to theory, or even conjecture, in its proper place; but the true aim of theory should never be left out of sight,—which is, to lead to fact; to a rule or result of practice.—And it should be likewise borne in mind that the Reader will attach little weight to mere generalizing statements (such as, that a certain remedy, procedure, or line of treatment has uniformly proved successful, etc.);—to be convinced, he wants to see positive evidence: recorded in clinical detail of cases; conditions found, courses pursued, and results achieved.

Do Not Fear,

however, that a communication you may be inclined to make would be devoid of value because you have but little time to spend on writing it! If your thought be a good one to yourself and for your patients' benefit, it will be equally so to your colleagues and their practice, and will be worth communicating. It need not come in the garb of an elaborate Scientific treatise; a simple "Letter to the Editor" will often be just as acceptable.

Some Rules of Order we should like to have our esteemed Contributors comply with:


Do *not* write on both sides of the sheet.

Write as legibly as you conveniently can (*names* especially so).

Leave a liberal margin on the sheet, or space between the lines. (*Close* writing is not conducive to correct typography; and what you save in writing material has to be expended a thousand-fold by us in eyesight, labor, and expense for printer's corrections.)

Address:—P. O. Box, 2535,
New York City.

Editor American Medico-Surgical Bulletin.



**ABSTRACT OF A PAPER ON URINALYSIS IN
GYNECOLOGY.***

By **HOWARD A. KELLY, M.D.**

A CAREFUL analysis of catheterized specimens of urine should be made before and after all gynæcological operations.

HELLER'S and the heat test are the most practical for the qualitative estimation of albumin, while FEHLING'S and NYLANDER'S solutions have proven efficient in testing for sugar.

The results of each analysis should be transferred to a special urinary chart, and filed with the patient's history.

In cases subjected to cœliotomy, the urine should be examined at least three times : the day before, and three days after operation, and during convalescence, about the twentieth day.

From an analysis of two hundred urinary charts, the following conclusions were drawn :

- 1) Albuminuria in pelvic inflammatory cases is usually due to cystitis.
- 2) Secondary changes in the kidneys are due either to absorption of septic products,

* Read before the Obstetric and Gynæcological Section of the ACADEMY OF MEDICINE, New York, May 25, 1893.

from pressure on the ureters, or from extension by contiguity.

3) Albumin is most frequently noted three days after operation, and in the majority of cases disappears during convalescence.

4) Decreased excretion of urine following cœliotomy, averaging 516 c.c. the second day, and gradually reaching normal on the tenth.

5) Increased specific gravity, probably due to decrease in watery constituents, while the solids are normal or increased in amount.

6) To concentration of the urine is probably due the vesical irritation so often observed following operations.

7) The increase in the percentage of cases showing albumin, is also due to vesical irritation.

Albumin was found in 46 (23%) of the 200 cases before operation, and in 66 (33%) after operation.

Casts were observed in 10 (5%) cases before, and in 30 (15%) after operation.

At time of discharge from the hospital, in only 8% of cases were casts still present. Although there is a definite relation between pelvic and kidney diseases, it is seldom that an operation is refused on account of renal changes.

In all cases dying after operation in the Gynæcological Department of the JOHNS

HOPKINS HOSPITAL during the past two years, not a single death was found at the autopsy to be due to kidney disease.

In conclusion, the practical results of this analysis may be summarized as follows :

1) Pelvic disease does not occasion renal disease of serious nature in a very large percentage of cases.

2) All urinary examinations must be made from catheterized specimens.

3) Presence of albumin, even in a large amount, is not significant unless casts are present.

4) The presence of small granular and hyaline casts does not contraindicate operation unless vascular and associated symptoms indicate renal disease.

5) Diminished secretion following operation, as low as 290 c.c. a day, is not unusual and need occasion no alarm as to the possibility of a ureter being ligated or the existence of any serious interference with the renal function.

6) Persistent diminished secretion, associated with elevated temperature and increasing rapidity of pulse, is not due to nephritis, but to septicæmia.

BALTIMORE, Md.

Issued Monthly.

\$2.00 per year.

American Medico-Surgical Bulletin

A JOURNAL OF PRACTICE AND SCIENCE.

THE BULLETIN PUBLISHING COMPANY, 73 William St., New York.

EDITORIAL STAFF.

WILLIAM HENRY PORTER, M.D.,
Chief Editor.

FREDERICK PETERSON, M.D.,
Associate Editor.

SAMUEL LLOYD, M.D.,
Department of Surgery.

J. RIDDLE GOFFE, M.D.,
Department of Obstetrics, Gynæcology
and Pædiatrics.

WILLIAM FANKHAUSER, M.D.,
Department of Materia Medica and
Therapeutics.

H. BAILLON, M.D., Paris,
Foreign Department.

ADOLPH ZEH, M.D.,
Pathology and General Medicine.

ANGELO ZUCCARELLI, M.D., Naples
Foreign Department.

WILLIAM C. GUTH, M.D.,
Pathology and General Medicine.

ADOLPH BARON, M.D.,
Diseases of Children.

GEORGE G. VAN SCHAICK, M.D.,
Pathology and Clinical Medicine.

HENRY F. QUACKENBOS, M.D.,
General Medicine.

W. TRAVIS GIBB, M.D.,
Gynæcology.

T. HALSTED MYERS, M.D.,
Orthopædic Surgery.

GUSTAV A. KLETZSCH, M.D.,
Gynæcology.

GEORGE THOMAS JACKSON, M.D.,
Dermatology.

J. CLIFTON EDGAR, M.D.,
Obstetrical Surgery.

JONATHAN WRIGHT, M.D.,
Laryngology.

T. S. SOUTHWORTH, M.D.,
Pædiatrics.

WILLIAM OLIVER MOORE, M.D.,
Ophthalmology and Otology.

WILLIAM B. COLEY, M.D.,
General Surgery.

16,199 ACTUAL SUBSCRIBERS BY SWORN COUNT.

