

QD  
421  
M8

UC-NRLF



QB 621 936

Committee of Revision and Publication of the Pharmacopoeia of the United States of America: 1890-1900.

---

# ALKALOIDAL ESTIMATION:

A BIBLIOGRAPHICAL INDEX OF CHEMICAL RESEARCH  
PREPARED FROM ORIGINAL LITERATURE  
FOR THE COMMITTEE OF REVISION

BY

PAUL I. MURRILL

UNDER THE DIRECTION OF

ALBERT B. PRESCOTT.

---

PUBLISHED BY THE COMMITTEE OF REVISION

---

ANN ARBOR.  
1898.

[THIS PAMPHLET IS NOT FOR SALE.]

LIBRARY  
OF THE  
UNIVERSITY OF CALIFORNIA.

GIFT OF


**PROF. W.B. RISING**

*Class*

105

1884

1884



Committee of Revision and Publication  
of the Pharmacopoeia of the  
United States of America: 1890-1900.

---

# ALKALOIDAL ESTIMATION:

A BIBLIOGRAPHICAL INDEX OF CHEMICAL RESEARCH  
PREPARED FROM ORIGINAL LITERATURE  
FOR THE COMMITTEE OF REVISION

BY

PAUL I. MURRILL

UNDER THE DIRECTION OF

ALBERT B. PRESCOTT.

---

PUBLISHED BY THE COMMITTEE OF REVISION,

UNIVERSITY OF  
MICHIGAN

ANN ARBOR.  
1898.

[THIS PAMPHLET IS NOT FOR SALE.]

Q 11421  
M 8

ANN ARBOR:  
THE INLAND PRESS.  
1898.

ANN ARBOR  
MICHIGAN

BIBLIOGRAPHICAL INDEX OF ALKALOIDAL ESTIMATION.

1877-1897.

The object of this work is to furnish a descriptive index of the work that has been done on the subject, rather than to abstract or summarize it.

For convenience, as large a list of references is given for each article as is practicable, the first referring to the original publication, and the others to republications or abstracts. In preparing it, the following publications have been examined up to the end of 1897:

Journal of the Chemical Society, London.

Chemisches Centralblatt.

American Journal of Pharmacy.

Proceedings of the American Pharmaceutical Association.

Proceedings of the Michigan State Pharmaceutical Association.

Jahresbericht der Pharmacie (ending with '95).

Répertoire de Pharmacie.

Also,

Lyon's Manual of Pharmaceutical Assaying (1886).

Dragendorff's Chemische Werthbestimmung einiger stark-wirkender Drogen, (1874).

Guareschi, Einführung in das studium der Alkaloide, Kunz-Krause's translation, (1896).

Other periodicals to which references are given are the following :

American Druggist.

Analyst, (London.)

Annalen der Chemie (Liebig).

Archiv der Pharmazie.

Archives de Médecine et de Pharmacie militaires.

Apotheker-Zeitung.

- Berichte der deutschen chemischen Gesellschaft.  
Berichte der deutschen pharmaceutischen Gesellschaft.  
Bulletin de la Société chimique de Paris.  
Chemist and Druggist.  
Chemical News.  
Chemiker-Zeitung.  
Comptes rendues des Séances de l'Academie des Sciences.  
Dingler's Polytechnisches Journal.  
Dorpat Dissertations.  
Druggists' Bulletin.  
Druggists' Circular.  
Ephemeris (Squibb).  
Festschr. 50-jährigen Stiftungsfeier des Schweiz. Apot.  
Ver.  
Fortschritt (Der), or Le Progrès, (Geneva.)  
Gazzetta chimica italiana.  
Helfenberger Annalen.  
Journal de Pharmacie et de Chimie.  
Journal für praktische Chemie.  
Journal of the American Chemical Society.  
Journal of Analytical and Applied Chemistry.  
Journal of the Society of Chemical Industry.  
Zhurnal russkova chimicheskova.  
Mitteil. pharm. Inst. Lab. Erlangen.  
New Remedies.  
Nederl. Tijdschrift voor Pharmacie, etc.  
Oesterreichisches Zeitschrift für Pharmacie.  
L'Orosi (Florence).  
Pharmaceutical Era.  
Pharmaceutical Journal and Transactions.  
Pharmaceutical Record.  
Pharmaceutische Centralhalle.  
Pharmaceutische Post.  
Pharmaceutische Rundschau.  
Pharmaceutische Zeitung.  
Pharmaceutische Zeitschrift für Russland.



Recueil des Travaux chimiques des Pays-Bas.  
Repertorium der analytischen Chemie.  
Repertorium der Pharmacie.  
Revue internationale scientif. et popul. des falsific des den-  
reés, aliment.  
Rundschau für die Interessen der Pharmacie, Chemie,  
u. s. w.  
Schweizerische Wochenschrift für Pharmacie.  
Transactions of the Academy of Science at St. Louis.  
L'Union pharmaceutique (Paris).  
Vierteljschr. Fortschr. Chem. Nahrungsm.  
Zeitschrift des österreich. Apot. Ver.  
Zeitschrift für analytische Chemie.  
Zeitschrift für physiologische chemie.  
Zeitschrift für Hygiene.  
Year Book of Pharmacy. London.

Abbreviations used are those of Bolton's "Bibliography of Chemistry," as far as possible. Others are believed to be self-explanatory.

References have been verified as far as practicable, and fully wherever any doubt has arisen in regard to possible errors.



# ALKALOIDAL ESTIMATION.

## PART I.—CHRONOLOGICAL INDEX.

- 1861: WAGNER. Method of separation, using iodo-potassium iodide.  
Dingler's Polyt. J. **161**, 40; Ztschr. anal. Chem. **1**, 102; Lyons' Manual, p. 33.
- 1864: GROVES, T. B. Methods of separation and purification of alkaloids.  
Pharm. J. Trans. (2), **6**, 275; Lyons' Manual, p. 33.
- 1872: (1) ZINOFFSKY. Estimation of alkaloids with Mayer's volumetric reagent.  
Dissert. Dorpat. 1872; Chem. Werthbest, p. 8.
- 1872: (2) CARLES. Estimation of quinine.  
Bull. Soc. chim. **18**, 98; Guareschi, Einf. Stud. Alk. 1896, p. 521.
- 1874: (1) ZINOFFSKY. Volumetric estimation coniine with phosphomolybdic acid.  
Chem. Werthbest, p. 44; Guareshi. Einf. Stud. Alk., p. 283.
- 1874: (2) DRAGENDORFF. Estimation of strychnine, gravimetric.  
Chem. Werthbest. p. 61; Lyons' Manual, p. 112.
- 1876: (1) SOKOLOFF, N. Estimation of coniine, alkalimetric.  
Chem. Centrbl. 1876, 604; J. Chem. Soc. **32**, 368.
- 1876: (2) MARKOWNIKOFF, B. W. Estimation of caffeine in tea.  
Ber. d chem. Ges. **9**, 1312; J. Chem. Soc. **31**, 110.
- 1876: (3) LYNN, J. Morphine estimation; comparison of methods.  
A. J. Pharm. **48**, 358; Chem. Centrbl. 1878, 158; J. Chem, Soc. **34**, 612.

- 1877: (1) BLYTH, A. W. Estimation of caffeine in tea.  
Analyst. **2**, 39; J. Chem. Soc. **32**, 517.
- 1877: (2) CAZENEUVE AND CAILLOL. Extraction and estimation of piperine in pepper.  
Bull. Soc. Chim. (2) **27**, 290; J. Chem. Soc. **32**, 516.  
Chem. Centrbl. 1877, 343;
- 1877: (3) MASING, E. Estimation of veratrine and physostigmine, with Mayers' reagent.  
Arch. Pharm. (3) **9**, 310; J. Chem. Soc. **32**, 367.
- 1877: (4) TESCHEMACHER, E. F. Estimation of morphine in opium.  
Chem. News, **35**, 47; Proc. A. Ph. A. 1877, 299; J. Chem. Soc. **32**, 231; Chem. Centrbl. 1877, 299.
- 1877: (5) PRESCOTT, A. B. Gravimetric estimation of quinin, by potassium mercuric iodide and by phospho-molybdate.  
Am. J. Pharm. **49**, 482.
- 1878: (1) PRESCOTT AND THUM. Comparison of methods for separation and estimation of the cinchona alkaloids.  
Proc. A. Ph. A. 1878, 823,
- 1878: (2) HEINTZ, E. Modification of Stas' method for extraction of alkaloids, using white bole, or quartz sand.  
Ztschr anal. Chem. 1878, 166; New Rem. 1878, 200;  
Proc. A. Ph. A., 1879, 478.
- 1878: (3) SCHRAGE, F. Separation of the cinchona alkaloids, using potassium sulphocyanide.  
Arch. Pharm. (3) **13**, 25; Chem. Centrbl. 1878, 582.
- 1878: (4) DRAGENDORFF. Extraction of theobromine from cacao.  
Jsb. d. Pharm. 1878, 493.
- 1878: (5) DWARS. Estimation of quinine in salts.  
Nieuw Tijds. Pharm. 1878, 267; Arch. Pharm. (3) **14**,  
149 and 329; Chem. Centrbl. 1879, 427 and 428; J.  
Chem. Soc. **36**, 488 and 492; Jsb. d. Pharm. 1878, 471.
- 1879: (1) DWARS. Separation of quinine and strychnine.  
Arch. Pharm. (3) **15**, 463; Chem. Centrbl. 1880, 78.

- 1879: (2) YVON. Polariscopic estimation of morphine and opium.  
J. Pharm. Chim. (4) **29**, 332; Chem. Centrbl. 1879, 363.  
Jsb. d. Pharm. 1879, 185.
- 1879: (3) PRUNIER, L. Estimation of cinchona alkaloids.  
J. Pharm. Chim. (4) **29**, 135; J. Chem. Soc. **36**, 489.
- 1879: (4) WOLFRAM, G. Estimation of theobromine in cacao and chocolate.  
Dingl. Polyt. J. **230**, 240; J. Chem. Soc. **36**, 406; Chem. Centrbl. 1879, 40.
- 1879: (5) LÖSCH, A. Quantitative estimation of alkaloids in plants.  
Pharm. Ztg., 1879, 89; Apoth.-Ztg., 1898, 15; Pharm. Ztschr. Russ. **18**, 545; Jsb. d. Pharm. 1879, 165.
- 1879: (6) LÖSCH, A. Methods of extractions of alkaloids using saturated alum solution.  
Pharm. Ztg. 1879, No. 87; Am. J. Pharm. 1880, 16; Lyons' Manual, p. 18; Phar. Ztschr. Russ. **18**, 545; Arch. Pharm. **16**, 317; Rep. de Pharm. 1880, 261.
- 1879: (7) PETIT, A. Rapid estimation of morphine in opium.  
J. Pharm. Chim. (4) **29**, 159; Chem. Centrbl. 1879, 363; Pharm. J. Trans. (3) **9**, 937. Rep. de Pharm. 1879, 100.
- 1879: (8) FLEURY, G. Estimation of morphine in opium.  
J. Pharm. Chim. (4), **29**, 341; Chem. Centrbl. 1879, 363.
- 1879: (9) FLUECKIGER, F. A. Estimation of morphine, gravimetric.  
Pharm. Ztg. 1879, Nos. 57, 59, and 74; Pharm. J. Trans. (3) **10**, 254; Jsb. d. Pharm. 1879, 183.
- 1879: (10) ————. Estimation of quinine in cinchona bark. Extraction and titration.  
Chem. News **40**, 209; J. Chem. Soc. **38**, 190.
- 1879: (11) PRESCOTT, A. B. Modification of the Hager-Jacobson method of estimation of morphine in opium.  
Proc. Am. Ph. A. 1879, 807; Jsb. d. Pharm. 1879, 186; Pharm. J. Trans. (3) **10**, 66, 128, and 132; J. Chem. Soc. **38**, 191 and 193.

- 1880: (1) PRESCOTT, A. B. Estimation of alkaloids by potassium mercuric iodide.  
Am. Chem. J. 1880, 2, 294; Ber. d. Chem. Ges. 14, 1421; J. Soc. 42, 646; Chem. News 45, 114 and 123; Proc. Am. Ph. A. 1882; 399.
- 1880: (2) THRESH, J. C. Volumetric estimation by means of bismuth potassium iodide.  
Pharm. J. Trans. (3) 10, 809, and 825; Jsb. d. Pharm. 1880, 156; J. Chem. Soc. 38, 763.
- 1880: (3) PATRONOULLARD. Estimation of caffeine in tea.  
Schweiz. Wochens. 1880, No. 15; Chem. Centrbl. 1880, 427; Rep. de Pharm. 1880, 52.
- 1880: (4) HAGER. Recognition and analytical determination of cinchona alkaloids.  
Pharm. Centralh. 21, 41I; Jsb. d. Pharm. 1881-2, 673.
- 1880: (5) HILBIG, C. Estimation of alkaloids in cinchona bark.  
Pharm. Ztschr. Russ. 19, 289 and 481; Chem. Centrbl. 1880, 551 and 713.
- 1880: (6) DE VRIJ. Estimation of quinine as herapathite.  
Am. J. Pharm. 1880, 394; Jsb. d. Pharm. 1881-2, 670; Rep. de Pharm. 1880, 485.
- 1880: (7) HESSE, O. The officinal test of quinine.  
Ber. d. chem. Ges. 13, 1517; J. Chem. Soc. 40, 63.
- 1880: (8) KERNER, G. Testing commercial quinine sulphate for for foreign alkaloids.  
Arch. Pharm. (3) 16, 186; J. Chem. Soc. 40, 63.
- 1880: (9) MYLIUS, E. Opium testing.  
Arch. Pharm. (3) 15, 310; J. Chem. Soc. 38, 829.
- 1880: (10) SCHMITT, E. Assay of opium.  
J. des Sci. Med. de Lille, — —; Rep. de Pharm. 1880, 442.
- 1880: (11) ROTHER. Assay of opium.  
Am. J. Pharm. 9, 541; Arch. Pharm. 17, 75; Rep de Pharm. 1880, 417.

- 1881: (1) PORTES and LANGLOIS. Rapid estimation of morphine in opium.  
J. Pharm. Chim. (5), 4, (December '81); Rep. de Pharm. 1882, 100; Chem. Centrbl. 1882, 60.
- 1881: (2) HAGER. Weighing of alkaloids as picrates.  
Pharm. Centrbl. 1881, 399; Jsb. d. Pharm. 1881-2, 631; Am. J. Pharm. 1882, 58, Proc. A. Ph. A. 1882, 399.
- 1881: (3) SKALWEIT, J. Estimation of nicotine in tobacco, volumetric.  
Arch. Pharm. 1881, 36; Rep. anal. Chem. 1, 165 and 303; Chem. Centrbl. 1881, 476 and 772; J. Chem. Soc. 42, 108; Am. J. Pharm. 1882, 60; Jsb. d. Pharm. 1881-2, 130 and 634; Lyons' Manual, p. 132.
- 1881: (4) PROLLIUS. Estimation of alkaloids in cinchona bark, gravimetric.  
Arch. Pharm. 1881, 85; Am. J. Pharm. 1882, 59; Proc. A. Ph. A. 1882, 204; J. Chem. Soc. 42, 246; Chem. Centrbl. 1881, 773.
- 1881: (5) SCHACHT, C. Estimation of alkaloids in wine of quinine.  
Arch. Pharm. (3) 14, 81; J. Chem. Soc. 40, 204.
- 1881: (6) HESSE, O. Optical estimation of cinchonidine in commercial quinine sulphate.  
Ann. d. Chem. 205, 217; J. Chem. Soc. 40, 315.
- 1881: (7) BLYTH, A. W. Estimation of quinine, volumetric with Mayer's reagent.  
Analyst. 1881, 162; J. Chem. Soc. 40, 1176.
- 1881: (8) EYKMAN. Assay of cinchona.  
Arch. Pharm. 1881, 814.
- 1881: (9) POEHL. Assay of Jaborandi for pilocarpine, gravimetric, using sodium phospho-molybdate.  
Year Book, 1881, 28 and 141; Lyon's Manual, p. 122.
- 1881: (10) MYLIUS, E. Estimation of morphine in opium, colorimetric.  
Phar. J. Trans. (3) 12, 6; J. Chem. Soc. 40, 945; Pharm. Centrbl. 1881, 97 and 105; Chem. Centrbl. 1881, 203; Jsb. d. Pharm. 1881-2, 256; 1883-4, 339.

- 1882: (1) SQUIBB, E. R. Modification of Flückiger's process for opium essay.  
Ephemeris, p. 14; Pharm. J. Trans. (3) 12, 724; J. Chem. Soc. 42, 666.
- 1882: (2) DE VRIJ. Modification of Prollius' method for estimation of total alkaloids in barks.  
Nieuwe Tijds. Pharm. 1882, Jan.; J. Pharm. Chim. (5) 6, 36; Pharm. J. Trans. (3) 12, 765; J. Chem. Soc. 42, 665.
- 1882: (3) DE VRIJ, J. E. Determination of quinine as herapathite.  
Am. J. Pharm. 1882, 58; Rund. Int. Pharm. 1883, 557; Pharm. Ztschr. Russ. 1883, 613; J. Pharm. Chim. (5) 5, 500; Rep. de Pharm. 1882, 243.
- 1882: (4) KISSEL. Estimation of alkaloids in cinchona barks.  
Pharm. Centrbl. 22, 556; Chem. Centrbl. 1882, 60; J. Chem. Soc. 42, 899.
- 1882: (5) ALESSANDRI. Assay of cinchona, extracting with oxalic acid.  
Phar. J. Trans. (3) 12, 994; Proc. A. Ph. A. 1882, 206; J. Chem. Soc. 42, 1003.
- 1882: (6) BIEL, J. Estimation of alkaloids of cinchona by Prollius' method.  
Arch. Pharm. (3) 20, 350; J. Chem. Soc. 42, 1139; Pharm. Ztschr. Russ. 1882; Mar. 28; J. Pharm. Chim. (5) 6, 37; Rep. de Pharm. 1882, 371.
- 1882: (7) MEYER, H. Estimation of cinchona alkaloids.  
Arch. Pharm. (3) 20, 721 and 812; J. Chem. Soc. 44, 388; Chem. Centrbl. 1883, 200.
- 1882: (8) SQUIBB. Assays of cinchona, extracting with amyl alcohol.  
Ephem., pp. 76, 105, 146; Chem. and Drug. 1882, 327; Rep. de Pharm. 1882, 417.
- 1882: (9) SKALWEIT, J. Estimation of nicotine.  
Arch. Pharm. (3) 20, 113; J. Chem. Soc. 42, 1005.



- 1882: (10) KISSLING, R. Estimation of nicotine in tobacco.  
Volumetric.  
Ztschr. anal. Chem. **21**, 75 and 383; J. Chem. Soc. **42**, 1005; Jsb. d. Pharm. 1881-2, 132; Lyons' Manual, p. 132; Pharm. Centrbl. **23**, 81; Chem. Centrbl. 1882, 252.
- 1882: (11) KISSLING, R. Estimation of nicotine in tobacco smoke.  
Dingl. Polyt. J. **244**, 64 and 234; Chem. Centrbl. 1882, 571.
- 1882: (12) GERRARD, A. W. Estimation of atropine.  
Pharm. J. Trans. (3) **13**, 190; Arch. Pharm. 1883, 74; Apoth. Ztg. 1898, 17.
- 1882: (13) FAIRTHORNE. Assay of cinchona.  
Am. J. Pharm. 1882, 548; Rep. de Pharm. 1882, 545.
- 1883: (1) HAGER. Method of extraction.  
Handb. d. Pharm. Praxis, Supplement, p. 62; Lyon's Manual, p. 19; Apoth. Ztg. 1898, 15.
- 1883: (2) KISSLING, R. Estimation of nicotine in tobacco, volumetric.  
Ztschr. anal. Chem. **22**, 2 and 199; Arch. Pharm. (3) **21**, 525; Jsb. d. Pharm. 1883-4, 169; 1885, 168; Chem. Ztg. 1885, 1336; Chem. Centrbl. 1883, 425.
- 1883: (3) SHIMOYANA, Y. Estimation of cinchona alkaloids.  
Jsb. d. Pharm. 1883-4, 234.
- 1883: (4) CONROY, M. Assay of nux vomica, tincture.  
Pharm. J. Trans. (13) **14**, 461.
- 1883: (5) DUNSTAN and SHORT. Assay of preparations of nux vomica.  
Pharm. J. Trans. **13**, 665; **14**, 441 and 621; **15**, 156; Year Book, 1883, 233 and 475; 1884, 274; Am. Drug. 1883, 124; Am. J. Pharm. 1883, 268; Lyons' Manual, p. 116.
- 1883: (6) DUNSTAN and SHORT. Separation of strychnine and brucine, using ferrocyanide.  
Pharm. J. Trans. (3) **14**, 290; Year Book, 1883, 469; Am. J. Pharm. 1883, 579; Arch. Pharm. (3) **22**, 40; Pharm. Ztg. 1884, No. 22; Pharm. Rund. 1883, 248; Jsb. d. Pharm. 1883-4, 768; Proc. A., Ph. A., 1884, 312.

- 1883: (7) SCHEIBE, E. Separation and estimation of morphine in chemico-legal investigations.  
Pharm. Ztschr. Russl. 1883, 49; Ber. d. chem. Ges. **16**, 797; Chem. Centrbl. 1883, 332; Arch. Pharm. (3) **21**, 290; Chem. News, **47**, 217; J. Chem. Soc. **44**, 1036; and **46**, 373.
- 1884: (1) CONROY, M. Estimation of morphine. Modification of the method of Portes and Langlois.  
Pharm. J. Trans. **15**, 473.
- 1884: (2) BERNHARDT, W. Estimation of morphine in opium.  
Pharm. Post. —; Schweiz. Wochens. **22**, 95; Chem. Centrbl. 1884, 479.
- 1884: (3) VON PERGER. Assay of opium, using baryta water, gravimetric.  
J. Prakt. Chem. (2) **29**, 97; Ber. d. chem. Ges. **17**, 241, R.; Chem. Centrbl. 1884, 479 and 684; J. Chem. Soc. **46**, 1217; Bull. Soc. Chim. 1885, 1, 43; Pharm. J. Trans. **15**, 384; Lyons' Manual, p. 119.
- 1884: (4) SQUIBB. Assay of coca.  
Ephemeris, p. 603.
- 1884: (5) SQUIBB. Assay of tea, coffee, and guarana.  
Ephemeris, pp. 606, 614, 616, 637, 641, 644.
- 1884: (6) LYONS. Three general assay processes.  
Drug. Cir. 1884, 114; Lyons' Manual, p. 20.
- 1884: (7) DUNSTAN and RANSOM. Assay of belladonna.  
Pharm. J. Trans. (13) **14**, 623; Lyons' Manual, p. 56; Am. J. Pharm. 1884, 279; Proc. A. Ph. A. 1884, 134.
- 1884: (8) REDWORD. Assay of belladonna.  
Pharm. J. Trans. (3), **14**, 623; Pharm. Ztg. 1884, 230; Apoth. Ztg. 1898, 17.
- 1885: (1) DUNSTAN and RANSOM. Assay of belladonna.  
Pharm. J. Trans. (3), **16**, 154 and 237; Pharm. Ztg. 1885, 813; Apoth. Ztg. 1898, 17; J. Chem. Soc. **50**, 104; Lyons' Manual, p. 56; Am. J. Pharm. 1885, 582; Am. Drug. 1885, 217; Proc. A. Ph. A. 1886, 392.
- 1885: (2) COBLENTZ. Assay of belladonna.  
Pharm. J. Trans. (3) **16**, 89; Arch. Pharm. 1885, 897; Proc. A. Ph. A. 1886, 393; Apoth. Ztg. 1898, 17; Am. Drug. 1885, 126.

- 1885: (3) SQUIBB. Assay of belladonna leaves.  
Ephemeris, pp. 848, 854, 859; Lyons' Manual, p. 57.
- 1885: (4) SQUIBB. Assay of coca.  
Ephemeris, pp. 726, 784, 788, 795, 871.
- 1885: (5) SCHEFFER, E. Estimation of nicotine in tobacco,  
volumetric with Mayer's reagent.  
Am. J. Pharm. **56**, No. 10; Arch. Pharm. (3), **23**, 153 and  
352; Jsb. d. Pharm. 1885, 372; Pharm. J. Trans. (3)  
**15**, 425; J. Chem. Soc. **48**, 604.
- 1885: (6) DE VRIJ. Estimation of quinine as oxalate.  
Arch. Pharm. (3), **23**, 209; Pharm. Ztg. 1885, 319; Pharm.  
Rund. 1885, 132; Rund. Int. Pharm. 1885, 314; Jsb. d.  
Pharm. 1885, 352.
- 1885: (7) DE VRIJ. Quantitative estimation of cinchonidine in  
quinine sulphate.  
Am. Drug. 1885, 153; Pharm. Centralb. 1885, 547; Arch-  
Pharm. (3) **23**, 442; Pharm. Rund. 1885, 97; Rund-  
Int. Phar. 1885, 756; Jsb. d. Pharm. 1885, 354.
- 1885: (8) DE VRIJ. Assay of quinine sulphate.  
Jour. Pharm. (5) **9**, 454; Chem. Centrbl. 1885, 968; J.  
Chem. Soc. **48**, 302; **50**, 397.
- 1885: (9) DE VRIJ. Assay of cinchona.  
Am. J. Pharm. 1885, 623; Chem. and Drug. 1885, 431;  
Proc. Am. Ph. A. 1885, 219.
- 1885: (10) SHIMOYANA, Y. Estimation of cinchona alkaloids.  
Herapathite method.  
Arch. Pharm. **23**, 81, and 209; J. Chem. Soc. **48**, 845;  
Lyons' Manual, p. 71; Jsb. d. Pharm. 1885, 351; 1886,  
302; Chem. Centrbl. 1885, 393.
- 1885: (11) LLOYD, J. U. Estimation of berberine, gravimetric.  
Drug. Circ. 1885, 22; Lyons' Manual, p. 101.
- 1885: (12) LYONS. Modification of Dragendorff's method for  
estimation of emetine, titrating with Mayer's reagent.  
Am. J. Pharm. 1885, pp. 531, 540, and 579; Pharm. Ztg.  
1885, 994; Proc. Am. Ph. A. 1886, 422; Lyons' Manual,  
p. 108; Proc. Mich. S. Ph. A. 1885, 185.

- 1885: (13) LYONS, A. B. Assay of nux vomica.  
Proc. Mich. S. Ph. A. 1885, 172; Drug. Circ. 1885, 253;  
1886, 137.
- 1885: (14) SCHWEISSINGER. Separation of strychnine and brucine, or estimation of strychnine in presence of brucine (double titration).  
Arch. Pharm. 1885, 611; Am. Druggist, 1885, 230; Proc. Am. Ph. A. 1886, 407; Lyons' Manual, 116.
- 1885: (15) LALIEN. Separation of a morphine salt from a mixture.  
J. Pharm. d'Anv. 1885, 6; Arch. Pharm. (3) 23, 512; Chem. Centrbl. 1885, 389.
- 1885: (16) WAINWRIGHT, J. H. Estimation of morphine in opium, gravimetric. U. S. Custom House method.  
J. Am. Chem. Soc. 7, 48; Ber. 18, 303 R.; Chem. Centrbl. 1885, 412; Pharm. J. Trans. 15, 990; Lyons' Manual, p. 118.
- 1885: (17) FLUECKIGER, F. A. Assay of opium, gravimetric.  
Arch. Pharm. 1885, 254 and 265; J. Chem. Soc. 48, 1165; Ber. 18, 651, R.; Chem. Centrbl. 1885, 523; Am. Drug. 1885, 149; Lyons' Manual, p. 120.
- 1885: (18) DIETERICH, E. Estimation of morphine in opium.  
Arch. Pharm. 24, 1023; Pharm. Centrbl. 27, 529; Jsb. d. Pharm. 1885, 109; 1886, 70; J. Chem. Soc. 52, 310.
- 1885: (19) BRITISH PHARMACOPEIA. Assay of cinchona.  
Br. Pharm. 1885, p. 111. [See 1898 (4).]

FOR QUININE AND CINCHONINE.—Mix 200 grains of red cinchona bark, in No. 60 powder, with 60 grs. of calcium hydrate; slightly moisten the powders with half an ounce of water; mix the whole intimately in a small porcelain dish or mortar; allow the moisture to stand for an hour or two, when it will present the characters of a moist dark brown powder, in which there should be no lumps or visible white particles. Transfer this powder to a six ounce flask, add three fluid-ounces of benzolated amyl alcohol, boil them together for about half an hour, decant and drain off the liquid onto a filter, leaving the powder in the flask; add more of the benzolated amyl alcohol to the powder, boil and decant as before, and repeat the operation a third time; then turn the contents of the flask on to the filter, and wash by percolation with more of the benzolated amyl alcohol until the bark is exhausted. If, during the boiling

a funnel be placed in the mouth of the flask, and another flask of cold water be placed in the funnel, this will form a convenient condenser which will prevent the loss of more than a small quantity of the boiling liquid. Introduce the collected filtrate while still warm into a stoppered glass separator; add to it 20 minims of diluted hydrochloric acid, mixed with two fluid drachms of water; shake them well together, and when the acid liquid has separated it may be drawn off, and the process repeated with distilled water slightly acidulated with hydrochloric acid, until the whole of the alkaloids have been removed. The acid liquid thus obtained will contain the alkaloids as hydrochlorates, with excess of acid. It is to be carefully and exactly neutralized with ammonia while warm, and then concentrated to the bulk of three fluid drachms. If now about 15 grs. of tartarated soda (sodium potassium tartrate) be added to the neutral hydrochlorates, and the mixture stirred with a glass rod, insoluble tartrates of quinine and cinchonidine will separate completely in about an hour; and these collected on a filter, washed and dried, will contain eight-tenths of their weight of the alkaloids quinine and cinchonidine, which, divided by 2, represents the percentage of those alkaloids. The other alkaloids will be left in the mother-liquor.

FOR TOTAL ALKALOIDS.—To the mother-liquor from the preceding process add solution of ammonia in slight excess. Collect, wash, and dry the precipitate, which will contain the other alkaloids. The weight of this precipitate, divided by 2, and added to the percentage weight of the quinin and cinchonidine, gives the percentage of total alkaloids.

1885: (20) BRITISH PHARMACOPŒIA. Assay of extract of nux vomica.

Br. Pharm. 1885, p. 163. [See 1898: (6).]

Dissolve 10 grains of the extract in half a fluid ounce of water, heating gently if necessary, and add a drachm of sodium carbonate previously dissolved in half a fluid ounce of water and half a fluid ounce of chloroform, agitate, warm gently, and separate the chloroform. Add to this half a fluid ounce of dilute sulphuric acid, with an equal bulk of water; again agitate, warm and separate the acid liquor from the chloroform. To this acid liquor add now an excess of ammonia, and agitate with half a fluid ounce of chloroform; when the liquors have separated, transfer the chloroform to a weighed dish, and evaporate the chloroform over a water bath. Dry residue for an hour, and weigh.

1885: (21) BRITISH PHARMACOPŒIA. Assay of opium.

Br. Pharm. 1885, p. 296. [See 1898: (7).]

Triturate together in a mortar 140 grains dried powdered opium, 60 grains freshly slaked lime, and 400 grain-measures of distilled water,

2

until a uniform mixture results; then add 1,000 grain-measures of distilled water, and stir occasionally during half an hour. Filter the mixture through a plaited filter about 3 in. in diameter into a wide mouthed bottle or stoppered flask (having the capacity of about 6 fluid ounces, and marked exactly at 1,040 grain-measures) until the filtrate reaches this mark. To the filtered liquid (representing 100 grains of opium) add 100 grain-measures of rectified spirit, and 500 grain-measures of ether, and shake the mixture; then add the chloride of ammonium, shake well and frequently during half an hour, and set it aside for twelve hours—Counterbalance two small filters; place one within the other in a small funnel, and decant the ethereal layer as completely as possible upon the inner filter. Add 200 grain-measures of ether to the contents of the bottle and rotate it; again decant the ethereal layer upon the filter, and afterwards wash the latter with 100 grain-measures of ether slowly added and in small portions. Now let the filter dry in the air and pour upon it the liquid in the bottle in portions, in such a way as to transfer the greater portion of the crystals to the filter. When the liquid has passed through the filter, wash the bottle and transfer the remaining crystals to the filter, with several small portions of distilled water, using not much more than 200 grain-measures in all, and distributing the portions evenly upon the filter. Allow the filter to drain, and dry it, first by pressing between sheets of bibulous paper, and afterwards at a temperature between 131° and 140° F. (55° and 60° C), and finally at 194° to 212° F. (96° to 100° C.) Weigh the crystals in the inner filter, counter-balancing by the other filter. The crystals should weigh ten grains (not less than 9½ nor more than 10½), corresponding to about 10 per cent. of morphine in the dry powdered opium.

1886: (1) HERETH, F. S. Method of titration with Mayers' reagent.

Pharm. Record 1886, p. 209; Am. J. Pharm. 1887, 2;  
Lyons' Manual, p. 29; Ztschr. anal. Chem. 26, 647; J.  
Chem. Soc. 54, 208.

1886: (2) SCHEIBLER. Method of purification of alkaloids, using phosphotungstic acid.  
Lyons' Manual, p. 33.

1886: (3) LYONS, A. B. Estimation of berberine, colorimetric.  
Lyons' Manual, p. 102.

1886: (4) LYONS, A. B. Estimation of hydrastine, gravimetric.  
Lyons' Manual, p. 104.

- 1886: (5) LYONS, A. B. Estimation of physostigmine (eserine) gravimetric.  
Lyons' Manual, p. 122.
- 1886: (6) LYONS, A. B. Approximate estimation of strychnine, gravimetric.  
Lyons' Manual, p. 113.
- 1886: (7) LYONS, A. B. Volumetric estimation of various alkaloids with Mayers' reagent.  
Am. J. Pharm. 1886, 579; 1887, 1; Proc. Mich. S. Ph. A. 1886, 167; Jsb. d. Pharm. 1887, 369; Proc. Am. Ph. A. 1887, 303; Lyons' Manual, p. 23.
- 1886: (8) LYONS, A. B. Estimation of cocaine with Mayers' reagent.  
Pharm. J. Trans. (3) 16, 1094; J. Chem. Soc. 50, 1087.
- 1886: (9) LYONS, A. B. Assay of cinchona, modification of Prollius' method.  
Drug. Circ. 1886, 150.
- 1886: (10) HOOPER, D. Optical method of analysis of quinine sulphate.  
Pharm. J. Trans. 17, 61; J. Chem. Soc. 50, 1086.
- 1886: (11) KASPAR. Assay of cinchona.  
Schweiz. Wochens. ————; Am. J. Pharm. 1886, 492; Proc. Am. Ph. A. 1887, 138.
- 1886: (12) DE VRIJ. Estimation of quinine as chromate.  
Arch. Pharm. (3), 24, 1073; J. Chem. Soc. 52, 404.
- 1886: (13) SNOW, H. W. Estimation of caffeine, gravimetric.  
Am. J. Phar. 1886, 483; Proc. Am. Ph. A. 1886, 107; Lyons' Manual, p. 97.
- 1886: (14) BRAITHWAITE and FARR. Estimation of morphine in opium.  
Pharm. J. Trans. (3) 17, 398; Proc. Am. Ph. A. 1887, 312.
- 1886: (15) VENTURINI, V. Comparison of methods of estimation of morphine in opium.  
Gaz. chim. ital. 16, 239; Ber. de chem. Ges. 19, 882, R.; J. Chem. Soc. 50, 1086.

- 1886: (16) JONES, H. W. Estimation of emetine.  
Pharm. J. Trans. (3) 17, 277; Pharm. Ztg. 1886, 633; J. Chem. Soc. 50, 1086.
- 1886: (17) FLUECKIGER. Assay of ipecac.  
Pharm. Ztg. 1886, 30.
- 1886: (18) WRAMPPELMEIER and MEINERT. As to a true aliquot part, by volume, of the total liquid in opium assay, and as to the solvent effect of ammonium chloride in the same liquid.  
Proc. Mich. State Phar. Assoc., 4, 127; Am. Drugg., 15, 203.
- 1887: (1) PLUGGE, P. C. Alkalimetric estimation of alkaloids.  
Arch. Pharm. (3) 25, 45; Pharm. Post. 20, 369; Chem. Drug. 1887, 388; Fortschritt. 1887, 301; Jsb. d. Pharm. 1887, 369; Chem. Centrbl. 1887, 312 and 971; Ber. d. chem. Ges. 20, 148 R.
- 1887: (2) LILLY. Alkaloidal assay, using Mayers' reagent.  
Pharm. Rundts, 1887, 181; Proc. Am. Ph. A. 1887, 42.
- 1887: (3) KREMEL, A. Assay of extractum aconiti.  
Pharm. Post. 20, 253; Chem. Centrbl. 1887, 733.
- 1887: (4) SQUIBB. Assay of coca.  
Ephemeris, p. 912.
- 1887: (5) RANSOM, F. Assay of ipecacuanha. Volumetric with Mayers' reagent.  
Pharm. J. Trans. (3) 18, 241; J. Chem. Soc. 52, 1147. Chem. Centrbl. 1887, 1410.
- 1887: (6) SNOW, H. W. Assay of ipecac. Volumetric, with Mayers' reagent.  
Proc. Mich. S. Ph. A. 1887, 93.
- 1887: (7) VAN ZWALUWENBERG, A. Bibliography of opium assay, 1817-1887.  
Proc. Mich. S. Ph. A., 1887, 140.
- 1887: (8) STILLWELL, C. M. Opium analysis.  
Am. Chem. J. 8, 295; J. Chem. Soc. 52, 403; Chem. News, 55, 43 and 54; Chem. Centrbl. 1887, 311.



- 1887: (9) BECKURTS, H. Estimation of morphine. Comparison of methods.  
Chem. Centrbl. 1887, 1411.
- 1887: (10) BIRD, F. C. J. Estimation of morphine.  
Pharm. J. Trans. 18, 470.
- 1887: (11) SCHRAUT, S. and BECKURTS, H. Estimation of morphine. Comparison of methods.  
Pharm. Centrbl. 28, 171; Chem. Centrbl. 1887, 605.
- 1887: (12) DIETERICH, E. Simplification and improvement of method for determining morphine.  
Pharm. Centralh. 28, 261 and 479; Ber. d. chem. Ges. 20, 344 and 604, and 21, 147 R.; Chem. Centrbl. 1887, 732 and 1328.
- 1887: (13) PLUGGE, P. C. Separation of the alkaloids of opium, using saturated sodium acetate, potassium ferricyanide, sodium salicylate, potassium sulphocyanide, etc.  
Arch. Pharm. 1887, 343; Niew. Tijdschr. v. Pharm. 1887, 69; Pharm. Ztg. 1887, 376; Pharm. Ztschr. Russ. 1887, 505; Rund. Int. Pharm. 1887, 543; Jsb. d. Pharm. 1887, 422; Proc. Am. Ph. A. 1887, 310.
- 1887: (14) KREMEL. Estimation of morphine in opium.  
Pharm. Post. 20, 661; Chem. Centrbl. 1887, 1529; J. Chem. Soc. 54, 635.
- 1887: (15) SCHLICKUM, O. Modification of Dieterich's method for determination of morphine in opium.  
Arch. Pharm. 25, 13 and 483; Ber. 20, 121 and 749 R.; Proc. Am. Ph. A. 1887, 311; J. Chem. Soc. 52, 622; Chem. Centrbl. 1487, 311 and 971.
1887. (16) ADRIAN and GALLOIS. Assay of opium.  
J. Pharm. Chim. (5) 15, 193; J. Chem. Soc. 52, 622; Chem. Centrbl. 1887, 345.
- 1887: (17) GÖBEL, H. Estimation of morphine. Criticism and improvement of U. S. P. process.  
Chem. Ztg. 11, 508; J. Chem. Soc. 52, 869.
- 1887: (18) WILEY, C. D. Opium assay. Comparison of methods of Squibb, Wainwright and Flückiger. Gravimetric, weighing free morphine.  
Proc. Mich. S. Ph. A. 1887, 138.

- 1887: (19) SQUIBB. Assay of opium.  
Ephemeris, p. 965.
- 1887: (20) JANDONS, A. Estimation of morphine in opium.  
Casop. cesk. lékár. 6, 128; Chem. Centrbl. 1887, 731.
- 1887: (21) BIEL, J. Estimation of morphine in opium.  
Pharm. Ztschr. Russ. 26, 241; Chem. Centrbl. 1887, 602.
- 1887: (22) RUDDIMAN, E. A. Estimation of quinine by Kerner's method.  
J. Anal. Chem. 1, 290; Chem. News 58, pp. 202, 216, and 226; J. Chem. Soc. 56, 323; Chem. Centrbl. 1888, 1595.
- 1887: (23) SCHÄFER, L. Estimation of cinchonidine in quinine sulphate.  
Arch. Pharm. (3) 25, 64 and 1033; J. Chem. Soc. 52, 623; 54, 636; Chem. Centrbl., 1888, 202.  
Pharm. Ztg. 32, 97; Chem. Centrbl. 1887, 314.
- 1887: (24) DIETERICH. Estimation of strychnine, alkalimetric.  
Pharm. Centralh. 1887, 21 and 29; Am. J. Pharm. 1887, 184 and 186; Proc. Am. Ph. A. 1887, 38.
- 1887: (25) BECKURTS and HOLST. Volumetric estimation of strychnine and brucine, using ferrocyanide.  
Pharm. Rund. 1887, 112; Arch. Pharm. 1887, 313; Jsb. d. Pharm. 1887, 434; Am. J. Pharm. 1887, 510; Pharm. Era, 1887, 292.  
Pharm. Centralh. 28, 119; Chem. Centrbl. 1887, 601.
- 1887: (26) BECKURTS and HOLST. Assay of strychnine preparations, alkalimetric.  
Pharm. Centralh. 28, 255; Chem. Centrbl. 1887, 732.
- 1887: (27) COWNLEY and PAUL. Estimation of caffeine in coffee.  
Pharm. J. Trans. 17, 648 and 821; Chem. Ztg. 1887, 55; Jsb. d. Pharm. 1887, 153.
- 1887: (28) COWNLEY and PAUL. Estimation of caffeine in tea.  
Pharm. J. Trans. 18, 419; J. Chem. Soc. 54, 539.
- 1887: (29) SMITH, E. D. Estimation of caffeine in coffee.  
Ztschr. östr. Apoth. Ver. 41, 359; Chem. Centrbl. 1887, 1270; J. Chem. Soc. 54, 539.

- 1887: (30) WAAGE, TH. Estimation of caffeine in tea leaves, gravimetric.  
Jsb. d. Pharm. 1887, 170.
- 1887: (31) LÖSCH, A. Estimation of caffeine in tea leaves.  
Fortschritt 1887, 128; Schweiz. Wochens. 1887, 209; Rund. Int. Pharm. 1887, 330; Arch. Pharm. 25, 443 and 447; Ztschr. Hygiene 1887, 177; Pharm. Centralh. 1887, 301; Jsb. d. Pharm. 1887, 172; J. Chem. Soc. 52, 1002; Pharm. Ztschr. Russ. 26, 177 and 178; Chem. Centrbl. 1887, 584.
- 1887: (32) CRIPPS, R. A. Indirect volumetric estimation of coniine, chlorometric.  
Pharm. Rund. 1887, 210; Rund. Int. Pharm. 1887, 732; Jsb. d. Pharm. 1887, 175.  
Pharm. J. Trans. (3) 18, pp. 12, 511, 820 and 888; Pharm. Era, 1887, 368; J. Chem. Soc. 54, 540.
- 1887: (33) LIGNON. Assay of Ipecac.  
J. Pharm. Chim. 1887, 550; Arch. Pharm. 1887, 642.
- 1887: (34) MACEWEN. Chemistry of calabar beans.  
Chem. and Drugg. 30, 193; Pharm. J. Trans. (3) 17, 641; Allen. Com. Org. Anal. 3, iii, 14.
- 1888: (1) KREMEL. Estimation of hydrastine, berberine, and emetine, gravimetric.  
Pharm. Post, 21, 149; Chem. Ztg. 1888, 84; Pharm. Rund. 1888, 111; Jsb. d. Pharm. 1888, 100; Arch. Pharm. (3) 26, 419; J. Chem. Soc. 54, 1351.
- 1888: (2) KREMEL, A. Estimation of alkaloids in nux vomica.  
Pharm. Post. 21, 534; Arch. Pharm. (3) 26, 899; J. Chem. Soc. 56, 322.
- 1888: (3) WIEDA. Estimation of cinchona alkaloids. Résumé of 25 different processes.  
Pharm. Record 1888, 249 and 264.
- 1888: (4) LENZ, W. Recent processes for testing quinin  
Comparison of methods.  
Ztschr. anal. Chem. 27, 549; J. Chem. Soc. 56, 86.
- 1888: (5) SNOW. Mayer's reagent for the estimation of alkaloids.  
Am. J. Pharm. 1888, 487; Proc. Mich. S. Ph.A. 1888, 107; Proc. Am. Ph. A. 1889, 686.

- 1888: (6) SNOW. Phospho-molybdate for the quantitative analysis of alkaloids, gravimetric and volumetric.  
Pharm. Era, 1888, 380; Pharm. Centralh. 1889, 14; Jsb. d. Pharm. 1889, 379; Am. J. Pharm. 1888, 529; Proc. Am. Ph. A. 1888, 133; J. Pharm. Chim. **19**, 23.
- 1888: (7) KREMEL, A. Estimation of caffeine in guarana.  
Pharm. Post. **21**, 101; Arch. Pharm. (3) **26**, 318; J. Chem. Soc. **54**, 876.
- 1888: (8) BIEL, J. Estimation of nicotine in tobacco extract.  
Pharm. Ztschr. Russ. **27**, 3; Arch. Pharm. (3) **26**, 322; J. Chem. Soc. **54**, 876; Chem. Centrbl. 1888, 425.
- 1888: (9) SQUIBB. Assay of coca and cocaine.  
Ephemeris, pp. 1104 and 1170.
- 1888: (10) SQUIBB. Assay of opium.  
Ephemeris pp. 1113 and 1150.
- 1888: (11) TESCHEMACHER and SMITH, J. D. Estimation of morphine in opium, with comparison of methods.  
Chem. News **57**, 93 and 103; Ber. d. chem. Ges. **21**, p. 548 R; J. Chem. Soc. **54**, 635; Pharm. J. Trans. **19**, 44; Chem. Centrbl. 1888, 562.
- 1888: (12) WILLIAMS, R. Estimation of morphine in opium.  
Chem. News, **57**, 134; J. Chem. Soc. **54**, 635; Chem. Centrbl. 1888, 648; Ber. d. chem. Ges. **21**, 545 R.
- 1888: (13) DIETERICH, E. and BARTHEL, G. Estimation of morphine and its water of crystallization.  
Pharm. Centralh. **29**, 315; Chem. Centrbl. 1888, 1044.
- 1888: (14) CAVENDONI. Alkaloidal estimation, gravimetric.  
L'Orosi, 1888, 301; Rep. de Pharm. 1890, 22; Pharm. Post. 1890, p. 115; Apoth. Ztg. 1898, 16.
- 1888: (15) VAN ITALLIE. Assay of belladonna leaves.  
Pharm. Weekblad. 1888, 883; Apoth. Ztg. 1888, 883
- 1889: (1) DIETERICH, E. Estimation of morphine by Flücker's method.  
Pharm. Centralh. **30**, 602; Chem. Centrbl. 1889, ii, 674 and 942.

- 1889: (2) BECKURTS, H. Separation and estimation of strychnine and brucine, and estimation of alkaloids in extracts containing chlorophyll.  
Pharm. Centralh. 1889, 574; Chem. Centrbl. 1889, ii, 541.
- 1889: (3) GÖBEL, H. Estimation of quinine and cinchonidine in bark, extraction with paraffine oil.  
Chem. Ztg. **13**, 906; Chem. Centrbl. 1889, ii, 521.
- 1889: (4) HOLDERMAN, E. Estimation of morphine by iodometric methods.  
Chem. Centrbl. 1889, ii, 542; Pharm. Centralh. 1889, 579; Pharm. Ztg. 1889, 582; Pharm. Rund. 1889, 258; Jsb. d. Pharm. 1889, 411; Pharm. Ztschr. Russ. 1889, 653.
- 1889: (5) FLUECKIGER, F. A. Determination of morphine in opium, gravimetric.  
Arch. Pharm. **27**, 721 and 769; Pharm. Ztg. 1889, 636; Pharm. Ztg. Russl. 1889, 759; Jsb. d. Pharm. 1889, 80; J. Chem. Soc. **58**, 94; Chem. Centrbl. 1889, ii, 715 and 716; Am. Drug. 1889, 191; Ber. d. Chem. Ges. **22**, 841 R.
- 1889: (6) GEROCK, J. E. Separation of brucine from strychnine, using nitric acid of s. g. 1.056.  
Arch. Pharm. **22**, 158; Am. J. Pharm. 1889, 180; Pharm. Runds. 1889, 116; Proc. Am. Ph. A. 1889, 702; Am. Drug. 1889, 91; Chem. Centrbl. 1889, i, 482.
- 1889: (7) SNOW. Estimation of strychnine and brucine in preparations of nux vomica.  
Proc. Mich. S. Ph. A. 1889, 73.
- 1889: (8) POPOVICI, M. Quantitative determination of nicotine, polariscopic.  
Diss. Erlangen; Zeitschr. f. Physiol. Chem. **13**, 5; Arch. Pharm. 1889, 558; Chem. Ztg. 1889, 148; Pharm. Ztschr. Russ. 1889, 314; Jsb. d. Pharm. 1889, 417; J. Soc. **56**, 802; Chem. Centrbl. 1889, i, 709; 1890, i, 361.
- 1889: (9) KISSLING, R. Estimation of nicotine in tobacco, polariscopic.  
Chem. Ztg. **13**, 1030; J. Chem. Soc. **58**, 430; Chem. Centrbl. 1889, ii, 521.

- 1889: (10) BLUNT, T. P. Assay of emetine in wine of ipecac, volumetric, with Mayer's reagent.  
Pharm. J. Trans. (3) 20, 254; J. Chem. Soc. 58, 310; Rep. de Pharm. 1890, 528.
- 1889: (11) JOHNSTONE, W. Analysis of pepper.  
Analyst. 14, 41; J. Chem. Soc. 58, 95.
- 1889: (12) VAN DER MARK: Estimation of alkaloids in coca leaves.  
Pharm. Ztschr. Russ. 28, 349; Schweiz. Wochens. 27, 174; Rep. de Pharm. 1889, 366; J. Pharm. Chim. (5) 20, 500; J. Chem. Soc. 58, 310.  
Ned. Tijds. Pharm. 89, 116; Pharm. Ztg. 39, 282; Chem. Centrbl. 1889, ii, 206.
- 1889: (13) FAWSETT, T. Estimation of cinchona alkaloids; colorimetric estimation of bromine, after adding excess of bromine water.  
Pharm. J. Trans. (3) 19, 914; J. Chem. Soc. 58, 309.
- 1889: (14) NEUMANN, S. Estimation of quinine in quinine tannate, gravimetric.  
Ztschr. anal. Chem. 1889, 28, 663; Jsb. d. Pharm. 1890, 420; Chem. Centrbl. 1890, i, 742.
- 1889: (15) VAN ITALLIE, L. Alkalimetric estimation of alkaloids in extracts.  
Nieuw Tijds Pharm. 1889, 6; Pharm. Centralh. 1889, 117; Analyst. 1889, 118; Arch. Pharm. 1889, 185; Pharm. Ztg. 1889, 73; Jsb. d. Pharm. 1889, 441; Rep. de Pharm. 1889, 511.
- 1889: (16) VAN DER MARK, B. Assay of nux vomica.  
Ned. Tijds. Pharm. 1889, 4.
- 1889: (17) CRIPPS, R. A. and WITHBY, A. Assay of ipecac.  
Pharm. J. Trans. (3) 19, 721; Pharm. Centralh. 1889, 441.
- 1889: (18) LANDRIN. Assay of cinchona.  
Compt. rend. 108, 750; Rep. de Pharm. 1889, 212.
- 1890: (1) VAN ITALLIE, L. Alkalimetric estimation of alkaloids.  
Nederl. Tijds. Pharm. 2, 209; Pharm. Ztg. 35, 442; Jsb. d. Pharm. 1890, 406; Chem. Centrbl. 1890, ii, 477.

- 1890: (2) SCHWEISSINGER AND SARNOW. Quantitative estimation of alkaloids in narcotic extracts. Volumetric.  
Pharm. Centralh. **31**, 583, 771; Jsb. d. Pharm. 1890, 406; Rep. de Pharm. 1891, 281; Am. J. Pharm. 1891, 96; Chem. Centrbl. 1891, i, 242.
- 1890: (3) EGELING. Gravimetric estimation of alkaloids in *extractum strychni*.  
Nederl. Tijdschr. Pharm. 1890, **2**, 242; Jsb. d. Pharm. 1890, 407; Rep. der Pharm. **1**, 240; Chem. Centrbl. 1891, i, 285.
- 1890: (4) CHRISTIANSEN, A. Determination of alkaloids, and their numerical equivalents. Iodometric.  
Chem. Ztg. 1890, 1346; Jsb. d. Pharm. 1890, 404; Chem. and Drug. 1890, 602; Proc. A. Ph. A. 1891, 609; J. Chem. Soc. **62**, 666; Chem. Centrbl. 1890, ii, 796.
- 1890: (5) BECKURTS, H. Estimation of strychnine in preparations.  
Arch. Pharm. **228**, 330; Chem. Centrbl. 1890, ii, 123 and 477.
- 1890: (6) SEATON and RICHMOND. Estimation of quinine in mixtures. (Double titration, using methyl orange and phenolphthalein.)  
Analyst, **15**, 42; Jsb. d. Pharm. 1890, 416; Chem. Centrbl. 1890, i, 741.
- 1890: (7) LOOFF, G. Estimation of morphine.  
Apot. Ztg. **5**, 271; Am. J. Pharm. **20**, 344; Rep. de Pharm. 1890, 480; Chem. Centrbl. 1890, i, 1082; J. Chem. Soc. **58**, 1349.
- 1890: (8) LOOFF, G. Estimation of morphine.  
Pharm. Centralh. **31**, 688; Chem. Centrbl. 1890, ii, 1034; J. Chem. Soc. **60**, 771.
- 1890: (9) HINSDALE, S. J. Colorimetric estimation of morphine.  
Chem. News, **62**, 77; Jsb. d. Pharm. 1890, 456; J. Chem. Soc. **58**, 1349; Chem. Centrbl. 1890, ii, 531.
- 1890: (10) DIETERICH, E. A shortening of the Helfenberger method of morphine estimation.  
Pharm. Centralh. **31**, 591 and 729; Jsb. d. Pharm. 1890, 455; Ztschr. anal. Chem. **30**, 105; J. Chem. Soc. **60**, 511; Chem. Centrbl. 1890, ii, 857.

- 1890: (11) DIETERICH, E. On morphine and emetine, and the time required for shaking out.  
Pharm. Centralh. 1890, **31**, 741; Jsb. d. Pharm. 1890, 454.
- 1890: (12) CLAASEN, E. On the quantitative estimation of codeine and morphine. Volumetric method, using platinic chloride.  
Pharm. Rund. 1890, **8**, 41; Pharm. Centralh. **31**, 115; Jsb. d. Pharm. 1890, 457; Chem. Centrbl. 1890, **i**, 741; J. Chem. Soc. **58**, 1198.
- 1890: (13) BLUNT, T. P. Alkalimetric estimation of emetine.  
Pharm. J. Trans. (3) **20**, 809; Jsb. d. Pharm. 1890, 442, Chem. Centrbl. 1890, **i**, 881.
- 1890: (14) VITÉ, F. Estimation of caffeine in tea.  
Mitteil. Erlangen, 1890, 113; Chem. Centrbl. 1890, **ii**, 274; J. Chem. Soc. **60**, 372.
- 1890: (15) SPENCER, G. L. Estimation of caffeine in tea.  
J. Anal. Chem. **4**, 158; Chem. Ztg. **14**, Ref. 175; Chem. Centrbl. 1890, **ii**, 172.
- 1890: (16) SUESS, P. Estimation of theobromine in cacao beans. Various methods discussed.  
Ztschr. Anal. Chem. **32**, 57; J. Chem. Soc. **64**, **ii**, 198.
- 1890: (17) THOMPSON, F. A. Assay of conium, modification of Lyons' method.  
Proc. Mich. S. Ph. A., 1890, 19.
- 1890: (18) ARNDT, E. M. Estimation of emetine.  
Apoth. Ztg. 1890, 780.
- 1890: (19) U. S. PHARMACOPŒIA. Assay of cinchona.  
U. S. P. 1890, p. 91.

FOR TOTAL ALKALOIDS.—To 20 gms. of cinchona in very fine powder and contained in a bottle with accurately ground glass stopper, add 200 c. c. of a previously prepared mixture of 19 volumes of alcohol, 5 vols. chloroform, and 1 vol ammonia water, stopper the bottle, and shake thoroughly and frequently during four hours. Then separate the liquid by pouring it into another bottle through a funnel containing a pellet of cotton so that no material loss by evaporation may result. Transfer 100 c. c. of the clear filtrate (representing 10 gms. cinchona) to a beaker, and



evaporate to dryness. Dissolve the residue of crude alkaloids thus obtained in 10 c. c. of water and 4 c. c. normal sulphuric acid, with the aid of gentle heat, filter the cooled solution into a separating funnel, and wash the beaker and filter until filtrate no longer has an acid reaction, using as small a quantity of water as possible. Now add 5 c. c. normal potassium hydrate to render decidedly alkaline, and extract alkaloids by shaking the mixture, first with 20 c. c., then repeatedly with 10 c. c. of chloroform, until a drop of the last chloroform extraction leaves no residue when evaporated on a watch glass. Evaporate the united chloroformic extracts in a tared beaker, dry the residue at 100° C. and weigh. The weight found, multiplied by 10 will give the percentage of total alkaloids in the specimen of cinchona tested.

FOR QUININE.—Transfer 50 c. c. of the clear filtrate remaining from the preceding process (and representing 5 gm. of cinchona) to a beaker, evaporate it to a dryness, and proceed as directed for total alkaloids, using only half the amounts of volumetric acid and alkali there directed. Add the united chloroformic extracts containing the alkaloids in solution, gradually, and in small portions at a time, to about 5 gm. of powdered glass contained in a porcelain capsule placed over a water bath, so that when the contents of the capsule are dry, all or nearly all of the dry alkaloids shall be in intimate admixture with the powdered glass, and the chloroform be completely expelled. Now moisten the residue with ether, and having placed a funnel containing a filter of a diameter of 7 c. m. and well wetted with ether over a small graduated tube (A), transfer the ether-moistened residue from the capsule. Rinse the latter several times, if necessary, with fresh ether, so as to transfer the whole of the residue to the filter, then percolate with ether added drop by drop, until exactly 10 c. c. of percolate have been obtained. Then collect another volume of 10 c. c. by similar slow percolation with ether, in a second graduated tube (B). Transfer the contents of the two tubes completely, (using ether for washing) to two small, tared capsules, properly marked (A and B) so as to avoid confusion, evaporate to a constant weight at 100° C. and weigh them. (The residue in A will contain practically all the quinine, together with a portion of the alkaloids less soluble in ether; the residue B will consist almost entirely of these alkaloids.) From the amount of residue obtained in capsule A deduct that contained in B, and multiply the remainder by 20. The product will represent, approximately, the percentage of quinine (containing 1 molecule of water) in the specimen of cinchona tested.

1890: (20) U. S. PHARMACOPŒIA. Assay of extract of nuxvomica.

U. S. P. 1890, p. 154.

Put 2 gm. of the dried extract of nux vomica into a glass separator, add to it 20 c. c. of a previously prepared mixture of 2 volumes of alcohol, 1 vol. ammonia water (s. g. 0.960), and 1 vol. water, and shake the separator till the extract is dissolved. Then add 20 c. c. of chloroform and agitate during five minutes. Allow the chloroform to separate, remove it as far as possible, pour into the separator a few c. c. of chloroform, and, without shaking, draw this off through the stop-cock to wash the outlet tube. Repeat the extraction with two further portions of chloroform of 15 c. c. each, and wash the outlet tube each time as just directed. Collect all the chloroformic solutions in a wide beaker, expose the latter to a gentle heat, on a water-bath, until the chloroform and ammonia are completely dissipated, add to the residue 10 c. c. of decinormal sulphuric acid measured from a burette, stir gently, and then add 20 c. c. of hot water. When solution has taken place add 2 c. c. of Brazil wood solution, and then carefully run in centinormal potassium hydrate, until a permanent pinkish color is produced. Divide the number of c. c. of centinormal potash used by 10, subtract the number found from the amount of decinormal acid used, multiply the remainder by 0.0364 and that product by 50 (or multiply at once by 1.82), which will give the percentage of total alkaloids in the extract, it being assumed that strychnine and brucine are present in equal proportion, and the above factor being found by taking the mean of their respective molecular weights [ $(334 + 394) \div 2 = 364$ ].

1890: (21) U. S. PHARMACOPŒIA. Assay of opium, crude, extract or tincture.

U. S. P. 1890, pp. 156, 292, and 424.

**EXTRACT OF OPIUM.**—Dissolve 4 grs. in 30 c. c. of water, filter the solution through a small filter, and wash the filter and residue with water, until all soluble matters are extracted, collecting the washings separately. Evaporate in a tared capsule, first the washings to a small volume, then add the first filtrate, and evaporate the whole to a weight of ten grams. Rotate the concentrated solution about in the capsule until the rings of extract are redissolved, pour the liquid into a tared Erlenmeyer flask having a capacity of 100 c. c., and rinse the capsule with a few drops of water at a time, until the entire solution weighs 15 gm. Then add 8.5 c. c. of alcohol, shake well, add 20 c. c. of ether and shake again. Now add 2.2 c. c. of ammonia water, stopper the flask with a sound cork, shake it thoroughly during ten minutes and set aside in a moderately cool place for six hours or over night.

**CRUDE OPIUM.**—Introduce the opium (which if fresh, should be in very small pieces, and if dry, in very fine powder) into a bottle having

a capacity of about 300 c. c. add 100 c. c. of water, cork it well, and agitate frequently during 12 hours. Then pour the whole as evenly as possible upon a wetted filter of 12 cm. diameter, and when the liquid has drained off, wash the residue with water, carefully dropped upon the edges of the filter and the contents, until 150 c. c. of filtrate are obtained. Then carefully transfer the moist opium back to the bottle by means of a spatula, add 50 c. c. of water, agitate thoroughly and repeatedly during 15 minutes and return the whole to the filter. When the liquid has drained off, wash the residue as before, until the second filtrate measures 150 c. c., and finally collect about 20 c. c. more of a third filtrate. Evaporate in a tared capsule, first, the second filtrate to a small volume, then add the first filtrate, rinsing the vessel with the third filtrate, and continue the evaporation until the residue weighs 14 gm. Rotate the concentrated solution about in the capsule until the rings of extract are redissolved, pour the liquid into a tared Erlenmeyer flask of 100 c. c. capacity, and rinse the capsule with a few drops of water at a time, until the entire solution weighs 20 gm. Then add 12.2 c. c. alcohol, shake well, add 25 c. c. of ether and shake again. Then add 3.5 c. c. ammonia water, stopper the flask, shake thoroughly during ten minutes and set aside in a moderately cool place for six hours or over night.

**TINCTURE OF OPIUM.**—Evaporate 100 c. c. to about 20 c. c., add 40 c. c. of water, mix thoroughly, and set aside for an hour, occasionally stirring, and disintegrating the resinous flakes adhering to the capsule. Then filter, and wash the filter and residue with water, until all soluble matters are extracted, collecting the washings separately. Evaporate in a tared capsule, first the washings to a small volume, then add the first filtrate, and evaporate the whole to a weight of 14 gm. Then proceed exactly as with crude opium from this point, using same quantities.

After standing for 6 hours or over night, remove the stopper carefully, and should any crystals adhere to it, brush them into the flask. Place in a small funnel two rapidly acting filters, of 7 cm. diameter, plainly folded, one within the other (the triple fold of the inner being laid against the single fold of the outer), wet them well with ether, and decant the ethereal solution as completely as possible upon the inner filter. Add 10 c. c. of ether to the contents of the flask, rotate it, and again decant the ethereal layer upon the inner filter. Repeat this operation with another portion of 10 c. c. of ether. Then pour into the filter the liquid in the flask, in portions, in such a way as to transfer the greater portion of the crystals to the filter, and, when this has passed through, transfer the remaining crystals to the filter by washing the flask with several portions of water, not using more than about 10 c. c.

in all. Allow the double filter to drain, then apply water to the crystals drop by drop, until they are practically free from mother-water, and afterwards wash them, drop by drop, from a pipette, with a saturated alcoholic solution of morphine. When this has passed through, displace the remaining alcohol by ether, using 10 c. c. or more if necessary. Allow the filter to dry in a moderately warm place, at a temperature not exceeding 60°, to constant weight, then carefully transfer the crystals to a weighed watch-glass, and weigh. The weight found represents percentage of morphine (crystallized) from tincture, multiplied by 10 gives percentage from crude opium, and multiplied by 25 percentage from extract.

1890. (22) NAGELVOORT, J. B. Rapid estimation of morphine in opium.  
Am. J. Pharm. 1890, 407; *Terapia moderna*, 1891, 80; *Rep. de Pharm.* 1891, 283.
- 1891: (1) LAMBERT, A. Polariscopic estimation of morphine.  
J. Pharm. Chim. (5) 3, 593; *Jsb. d. Pharm.* 1891, 550; J. Chem. Soc. 60, 1043; *Chem. Centrbl.* 1891, ii, 233.
- 1891: (2) LLOYD, J. U. Assay of galenical preparations. (Using ferric chloride and sodium bi-carbonate, and extracting with chloroform.)  
*Proc. A. Ph. A.*, 1891, 124;
- 1891: (3) THOMPSON, F. A. Method for the assay of preparations of belladonna, stramonium and henbane, cinchonas, colchicum, coca, and conium. (Using oak sawdust).  
*Proc. Mich. St. Ph. A.* 1891, 67.
- 1891: (4) HAUBENSAK, W. Estimation of total alkaloids in cinchona.  
*Schweiz. Wochens.* 29, 147; *Chem. Centrbl.* 1891, i, 1006; *J. Chem. Soc.* 60, 1402.
- 1891: (5) SPENCER, G. L. Estimation of caffeine in tea.  
J. Anal. Chem. 4, 390; *Chem. Centrbl.* 1891, i, 373; J. Chem. Soc. 60, 964; *Chem. News*, 63, 18.
- 1891: (6) PAUL, B. H. Estimation of caffeine in tea, etc.  
*Pharm. J. Trans.* 21, 882; *J. Chem. Soc.* 60, 1403; *Rev. intern. falsific.* 4, 210; *Chem. Centrbl.* 1891, ii, 595.

- 1891: (7) PEZZOLATO, A. Determination of nicotine in presence of ammonia.  
Gaz. chim. ital. **20**, 780; Ber. d. chem. Ges. **24**, 222, R; J. Chem. Soc. **60**, 771; Chem. Centrbl. 1891, i, 287; Einf. Stud. Alk. 293.
- 1891: (8) ALLEN, A. H. Assay of aconite preparations.  
Pharm. J. Trans. (3) **22**, 230; J. Chem. Soc. **62**, 392.
- 1891: (9) BECKURTS, H. Estimation of alkaloids in extracts of *nux vomica*, aconite, belladonna, and hyoscyamus, etc.  
Pharm. Rund. 1891, 240; Apoth. Ztg. 1891, 537; Am. J. Pharm. 1892, 26.
- 1891: (10) KREMEL, A. Estimation of coniine.  
Ann. di Chim. e di Farm. **14**, 149; Einf. Stud. Alk. p. 542.
- 1892: (1) NICHOLS, H. T. and NORTON, T. H. Examination of the Lloyd method.  
J. Anal. Chem. **6**, 162; Chem. Centrbl. 1892, ii, 188.
- 1892: (2) ALLEN, A. H. Methyl orange as the best indicator for titration of alkaloids.  
Chem. and Drug. 1892, 104; Proc. Am. Ph. A. 1892, 1035.
- 1892: (3) ALLEN, A. H. Volumetric estimation of alkaloids.  
Chem. News **66**, 259; Ber. **26**, 389, R; J. Chem. Soc. **64**, ii, 252; Chem. Centrbl. 1893, i, 137.
- 1892: (4) PARTHEIL, A. Use of ethereal iodo-eosin as an indicator for alkaloid.  
Apot. Ztg. 1892, 435; Am. J. Pharm. 1892, 522; Chem. Centrbl. 1892, ii, 638.
- 1892: (5) BARTHE, L. Volumetric estimation of alkaloids. (Double titration, with phenolphthalein and litmus.)  
Compt. rend. 1892, **115**, 512; Chem. News, 1892, 223; Jsb. d. Pharm. 1892, 491; Proc. A. Ph. A. 1893, 817; Am. J. Pharm. 1892, 368; J. Chem. Soc. **64**, ii, 146.
- 1892: (6) BARTHE, L. Analysis of quinine sulphate and estimation of quinine in presence of other cinchona alkaloids.  
Compt. rend. **115**, 1085; J. Chem. Soc. **64**, ii, 199; Chem. Centrbl. 1893, i, 233.

- 1892: (7) SCHMIDT, J. H. Estimation of quinine in cinchona barks.  
Pharm. Centralh. **33**, 594; Chem. Centrbl. 1892, ii, 946; J. Chem. Soc. **64**, ii, 310.
- 1892: (8) KUERSTEINER, C. Assay of cinchona, gravimetric.  
Pharm. Ztg. 1892, 750; Am. J. Pharm. 1893, 74; Proc. A. Ph. A. 1893, 404.  
Schweiz. Wochens **30**, 473; Chem. Centrbl. 1893, i, 137; Rep. de Pharm. 1894, 75.
- 1892: (9) PINETTE, J. Estimation of nicotine,  
Chem. Ztg. **16**, 1072; Chem. Centrbl. 1892, ii, 560.
- 1892: (10) BIRD, F. C. J. Estimation of emetine with Mayer's reagent.  
L'Union pharm. 1892, No. 10; Jsb. d. Pharm. 1892, 519.
- 1892: (11) KOTTMAYER, G. Estimation of emetine in ipecacuanha root, volumetric.  
Pharm. Post, 1892, 913 and 933; Jsb. d. Pharm. 1892, 182; Chem. Centrbl. 1892, ii, 638.
- 1892: (12) KELLER, C. C. Estimation of emetine.  
Schweiz Wochens. **30**, 501 and 509; Chem. Centrbl. 1893, i, 235 and 322; J. Chem. Soc. **64**, ii, 397; Jsb. d. Pharm. 1893, 171; Am. J. Pharm. 1893, 78.
- 1892: (13) TRAUB. Estimation of alkaloids, iodometric.  
Schweiz. Wochens. 1892, 22; Apoth. Ztg. 1892, —.
- 1892: (14) DOMERGUE and NICOLAS. Estimation of caffeine in tea.  
J. Pharm. Chim. (5) **25**, 302; Rep. de Pharm. 1892, 155.
- 1892: (15) CAZENEUVE and BIÉTRIX. Estimation of caffeine in tea.  
Moniteur scientif. 1892, April; Rep. de Pharm. 1892, 203.
- 1892: (16) GUILLOT. Rapid estimation of eaffeine.  
Arch. de Méd. et de Pharm. milit; Rep. de Pharm. 1892, 374; Apothek. Ztg. 1893, 132; Jsb. d. Pharm. 1893, 721; Chem. Centrbl. 1893, i, 865; J. Chem. Soc. **64**, ii, 608.
- 1892: (17) BECK, C. R. Assay of ipecac.  
Pharm. J. Trans. (3) **22**, 891; Rep. de Pharm. 1892, 273.

- 1893: (1) KELLER, C. C. Alkalimetric estimation of alkaloids in cinchona bark.  
Festschr. Stiftungsfeier Schweiz. Apoth. Ver; Pharm. Ztg. 1893, 786; Jsb. d. Pharm. 1893, 165.
- 1893: (2) KELLER, C. C. Estimation of emetin.  
Schweiz. Wochens. **31**, 470 and 485; Jsb. de Pharm. 1893, 175; Chem. Centrbl. 1894, i, 236.
- 1893: (3) KELLER. Estimation of emetin in ipecacuanha root.  
Pharm. Post, **25**, 913; J. Pharm. Chim. (5) **27**, 465; J. Chem. Soc. **64**, ii, 560.
- 1893: (4) KELLER, C. C. Estimation of alkaloids in strychnos seeds. Titration using iodo-eosin.  
Schweiz. Wochens. 1893, —; Pharm. Ztg. 1893, No. 96 and 101; Apoth. Ztg. **8**, 542; Ztschr. anal. Chem. **35**, 491; J. Chem. Soc. **66**, ii, 491.
- 1893: (5) MEYER, A. Valuation of ipecacuanha root, volumetric.  
Apoth. Ztg. 1893, pp. 167, 178; Jsb. d. Pharm. 1893, 173.
- 1893: (6) GRANDVAL and LAJOUX. Estimation of caffeine.  
J. Pharm. Chim. **27**, 545; Rep. de Pharm. 1893, 301; Rev. intern. falsific. 1893, 168; Jsb. d. Pharm. 1893, 722. J. Chem. Soc. **64**, ii, 559; Chem. Centrbl. 1893, ii, 166.
- 1893: (7) KELLER, C. C. Assay of alkaloidal drugs, volumetric.  
Oestr. Ztschr. Pharm. 1893, 563 and 586; Am. J. Pharm. 1894, 42; Proc. A. Ph. A. 1894, 532.
- 1893: (8) SOKOLOFF. Estimation of caffeine in tea.  
Chem. and Drug. 1893, 210; Proc. A. Ph. A., 1894, 514. J. Russ. Chem. Soc. **24**, 130; J. Chem. Soc. **64**, ii, 352.
- 1893: (9) SUESS, P. Estimation of theobromine.  
Ztschr. anal. Chem. **32**, 57; Chem. Centrbl. 1893, i, 322.
- 1893: (10) BRUNNER, H. and LEINS, H. Quantative separation of caffeine and theobromine, using ammonio silver nitrate.  
Schweiz. Wochens. **31**, 85; Jsb. d. Pharm. 1893, 720; Chem. Centrbl. 1893, i, 802; J. Chem. Soc. **64**, ii, 608.
- 1893: (11) DE WIJS, B. J. C. Estimation of morphine in opium.  
Nederl. Tijds. Pharm. **5**, 282; Chem. Centrbl. 1893, ii, 738.

- 1893: (12) CANNEPIN and VAN EIJK. Titration of morphine, cochineal as indicator.  
Bull. Soc. Chim., (3) 9, 437; Ber. 26, 837, R; J. Chem. Soc. 64, ii, 607; Chem. Centrbl. 1893, ii, 299; Rep. de Pharm. 1893, 249.
- 1893: (13) HEUT, G. Estimation of coniine and nicotine.  
Arch. Pharm. 231, 376 and 658. J. Chem. Soc. 64, ii, 607; Chem. Centrbl. 1894, i, 307.
- 1893: (14) VEDRÖDI. Analysis of tobacco. Study of methods.  
Ztschr. anal. Chem. 32, 277; J. Chem. Soc. 64, ii, 504; Chem. Centrbl. 1893, ii, 164.
- 1893: (15) THOMPSON, F. A. Estimation of hydrastine, and separation from berberine. Volumetric.  
Am. J. Pharm. 1893, 371; Proc. A. Ph. A., 1893, 691.
- 1893: (16) RANWEZ. Estimation of alkaloids in preparations of aconite, hyoscyamus, stramonium and cicuta.  
J. Pharm. Chim. (5) 28, 208 and 264; Rep. de Pharm. 1893, 344; Chem. Centrbl. 1893, ii, 737 and 890.
- 1893: (17) CASPARI, C. and DOHME, R. L. Titration as a means of assaying drugs. Discussion of methods.  
Am. J. Pharm. 1893, 473 and 478. Proc. A. Ph. A. 1894, 555.
- 1893: (18) GRANDVAL and LAJOUX. Estimation of alkaloids with potassium mercuric iodide.  
J. Pharm. Chim. 1893, Aug. 15; Rep. de Pharm. 1893, 436; Rev. inter. falsific. 6, 186 and 207, Chem. Centrbl. 1893, ii, 597 and 668.
- 1893: (19) GRANDVAL, A. and LAJOUX, H. Estimation of alkaloids in drugs.  
J. Pharm. Chim. 28, 99 and 152; Jsb. d. Pharm. 1893, 483; J. Chem. Soc. 64, ii, 608; Apoth. Ztg. 1893, 90; Rep. de Pharm. 1893, 392.
- 1893: (20) VITALI, D. Volumetric estimation of certain metals, and salts of the alkaloids. Applies to sulphates and halides.  
L'Orosi, 16, 181; Pharm. Ztg. 1893, 321; Jsb. d. Pharm. 1893, 207; Pharm. Post, 1893, 297; Am. J. Pharm. 1893, 384; J. Chem. Soc. 64, ii, 607.



- 1893: (21) VAN LEDDEN-HULSEBOSCH. Determination of alkaloids in extracts, gravimetric.  
Pharm. Centralh. 1893, 101 and 289; Jsb. d. Pharm. 1893, 593; Am. Drug. 1893, 372; Proc. A. Ph. A., 1893, 404.
- 1893: (22) LIEUNNIGH, F. Criticism of Hulsebosch's process.  
Pharm. Centralh. 1893, 591; Chem. Centrbl. 1893, ii, 890; Proc. A. Ph. A., 1894, 1092; J. Chem. Soc. 66, ii, 126.
- 1894: (1) KELLER, C. C. Alkaloidal assaying.  
Schweiz. Wochens. 1894, Nos. 5, 6, 7, 12, 13; Pharm. Ztg. 1894, 344 and 361; Jsb. d. Pharm. 1894, 11; Proc. A. Ph. A. 1895, 537.
- 1894: (2) FARR and WRIGHT. Preference given to gravimetric rather than volumetric methods.  
Pharm. J. Trans. (3) 25, 121; Proc. A. Ph. A. 1895, 623; Chem. Centrbl. 1894, ii, 602.
- 1894: (3) SCHWICKERATH, K. Alkaloidal assaying of drugs.  
Pharm. Rund. 1894, 137; Jsb. d. Pharm. 1894, 24; Drug. Bull. 1894, 246; Chem. Centrbl. 1894, ii, 302.
- 1894: (4) BECKURTS. Estimation of alkaloids in drugs.  
Apoth. Ztg. 1894, 661 and 750; Pharm. Ztg. 1894, 676; Pharm. Centralh. 1894, 566; Jsb. d. Pharm. 1894, 26.
- 1894: (5) DIETERICH. Discussion of methods for determination of alkaloids.  
Helfenb. Annal. 1894, 15; Pharm. Centralh. 1895, 163; Jsb. d. Pharm. 1895, 500; Proc. A. Ph. A. 1895, 545; Chem. Centrbl. 1895, i, 814.
- 1894: (6) KUNZE, W. E. Separation and estimation of the cacao alkaloids. Volumetric for theobromine.  
Ztschr. anal. Chem. 33, 1; Pharm. Centralh. 34, 108; Rep. de Pharm. 1895, 225; J. Chem. Soc. 66, ii, 168; Chem. Centrbl. 1894, i, 398.  
Schweiz. Wochens. 32, 334; Chem. Centrbl. 1894, ii, 179.
- 1894: (7) THIEL, J. Estimation of theobromine.  
Forschungsber. Lebensm. 1, 108; Chem. Centrbl. 1894, i, 878.
- 1894: (8) BRUNNER, H. Quantitative separation of theobromine and caffeine.  
Schweiz. Wochens. 32, 126; Chem. Centrbl. 189 932.

- 1894: (9) HILGER and EMINGER. Estimation of theobromine in cocoa.  
Forschungsber. Lebensm. **1**, 292; Chem. Centrbl. 1894, ii, 462; J. Chem. Soc. **68**, ii, 542.
- 1894: (10) STÆDER, W. Estimation of alkaloids in cortex granati.  
Nederl. Tijds. Pharm. **6**, 39; Chem. Centrbl, 1894, i, 606.
- 1894: (11) CUSHMAN, A. S. Detection and estimation of strychnine in corpses.  
Trans. Acad. Sci. St. Louis, **6**, 17; Chem. Centrbl. 1894, ii, 461; J. Chem. Soc. **68**, ii, 542.
- 1894: (12) DOTT, D. B. Assay of opium. Acid titration.  
Pharm. J. Trans. (3) **24**, 847; J. Chem. Soc. **70**, ii, 283; Apoth. Ztg. **4**, 795; Rep. de Pharm. 1894, 509.
- 1894: (13) KELLER, C. C. Estimation of hydrastine, gravimetric.  
Apoth. Ztg. 1894, 52 and 133; Am. J. Pharm, 1894, 201; Proc. A. Ph. A. 1894, 546.
- 1894: (14) LEINS, H. Estimation of caffeine and theobromine.  
Inaug.-Dissert. Lausanne, 1894, pp. 12 and 44; Guareschi, Einführung in das studium der Alkaloide. p. 405.
- 1895: (1) KEBLER, L. F. Estimation of morphine in opium.  
J. Soc. Chem. Ind. **14**, 464; Ber. d. Chem. Ges. **29**, 146, R; J. Chem. Soc. **70**, ii, 403; Chem. Centrbl. 1895, ii, 257.
- 1895: (2) VEDRÜDI, V. Estimation of nicotine and ammonia in tobacco.  
Ztschr. anal. Chem. **34**, 413; J. Chem. Soc. **68**, ii, 541.
- 1895: (3) KISSLING, R. Estimation of nicotine and ammonia in tobacco.  
Ztschr. anal. Chem. **34**, 731; J. Chem. Soc. **70**, ii, 401.
- 1895: (4) SOKOLOFF. Estimation of caffeine in tea.  
Rev. intern. falsific. **8**, 84; Chem. Centrbl. 1895, i, 450.

- 1895: (5) CRIPPS, R. A. Assay of ipecac.  
Pharm. J. Trans. (3) 25, 1093; J. Chem. Soc. 70, ii, 284.
- 1895: (6) UMNEY, J. C. Volumetric estimation of aconitine in pharmaceutical preparations; saponification and estimation of acid produced.  
Pharm. J. Trans. (3) 25, 860; Proc. A. Ph. A. 1895, 539; J. Chem. Soc. 70, ii, 283.
- 1895: (7) KEBLER and LA WALL. Volumetric estimation of alkaloids, with comparison of various indicators.  
Am. J. Pharm. 1895, 499; Jsb. d. Pharm. 1895, 408; Proc. A. Ph. A. 1896, 798  
J. Am. Chem. Soc. 17, 822; J. Chem. Soc. 70, ii, 551; Chem. Centrbl, 1895, ii, 1090.
- 1895: (8) VAN LEDDEN-HULSEBOSCH. Estimation of alkaloids.  
Pharm. Centralh. 16, 205; Ber. d. chem, Ges. 28, 1066, R.
- 1895: (9) KELLER, C. C. Estimations of alkaloids in nux vomica.  
Schweiz. Wochens. 33, 452; Chem. Centrbl. 1896, i, 228; J. Chem. Soc. 72, ii, 84; Chem. Ztg. 19, Ref. 392; Einf. Stud. Alk. p. 504.
- 1895: (10) MENDINI, A. Estimation of emetine.  
Boll. chim.-farm. 1895, 590; Rep. de Pharm. 1895, 515.
- 1895: (11) KELLER, C. C. Estimation of cocaine.  
Schweiz. Wochens. 33, 453; Guareschi Einf. Stud. Alk. 1896, 492.
- 1895: (12) GERMAN PHARMACOPCEIA: Assay of cinchona.  
Arzneibuch f. d. deutsche Reich, 1895, p. 79.

20 gm. of finely powdered bark are shaken vigorously and repeatedly with 10 c. c. ammonia, 20 c. c. alcohol and 170 c. c. ether, and after 24 hours 100 c. c. of the clear liquid is decanted. Add to this 3 c. c. normal hydrochloric acid and 27 c. c. water, and distill off the ether and alcohol, and if necessary, add enough normal hydrochloric acid to make slightly acid. When cold, add 3 c. c. normal caustic potash, or enough to redden phenolphthalein, and filter. The precipitate is washed repeatedly with small amounts of water until the washings are no longer alkaline (to phenolphthalein). Press gently with filter paper, and dry, first over sulphuric acid, then at 100° and weigh.

1895: (13) GERMAN PHARMACOPŒIA. Assay of opium.

Arzneibuch. f. d. deutsche Reich, 1895, p. 236.

Rub up in a mortar 6 gms. powdered opium with 6 c. c. water; dilute, and wash the mixture with water into a weighed flask, and make up to weight of 54 gms. Shake repeatedly during an hour, and filter through a plaited filter of 10 c. m. diameter. To 42 gm. of the filtrate add 2 gm. of a mixture of 17 parts ammonia and 83 parts water, mix well but avoid excessive shaking, and filter at once through plaited filter. 36 gms. of this filtrate is placed in a tared flask, 10 gm. ether is added, the flask is shaken, and 4 gm. of the ammonia diluted as above, and the shaking continued until the liquid has cleared. It is then allowed to stand 6 hours. Pour off the ethereal layer as completely as possible on to a plain filter of 8 c. m. diameter, to the aqueous liquid add 10 gm. ether, agitate a few minutes and again pour off the ether. Then pour the water on the filter, at the same time transferring the crystals; wash twice with 5 gm. water saturated with ether. Dry the filter at 100°, transfer crystals completely to the flask. Dry to constant weight and weigh.

1895: (14) VAN ITALLIE. Valuation of narcotic extracts.

Runds. für Pharm. 21, 327; Rep. de Pharm. 1895, 359.

1896: (1) PRESCOTT, A. B. Periodides of the alkaloids as molecular forms for estimation.

Pharm. Rund. 1896, 123; Proc. A. Ph. A. 1896, 800; J. Am. Chem. Soc 17, 859.

1896: (2) BECKURTS, H. and FRERICHS, G. Estimation of alkaloids in pharmaceutical extracts.

Apoth. Ztg. 11, 916; Chem. Centrbl. 1897, i, 205.

1896: (3) KELLER. Process for alkaloidal assay.

Oest. Ztschr. Pharm. 1896, 154.

1896: (4) KIPPENBERGER, K. New method for quantitative isolation of alkaloids from ptomains, using periodide, acetone, etc.

Ztschr. anal. Chem. 35, 407; Ber. d. Chem. Ges. 22, 1009, R; J. Chem. Soc. 70, ii, 681; Chem. Centrbl. 1896, ii, 761.

1896: (5) KIPPENBERGER, K. Titration of alkaloids with iodine solution.

Ztschr. anal. Chem. 34, 294; and 35, 10 and 422; Ber. d. chem. Ges. 28, 796; 29, 598 and 1010, R; J. Chem. 68, ii, 467, and 70, ii, 282 and 682; Chem. Centrbl. 1896, i, 515 and ii, 762.

- 1896: (6) BECKURTS, H. Estimation of hydrastine and berberine in extractum hydrastis canadensis.  
Apoth. Ztg. **11**, 552; Chem. Centrbl. 1896, ii, 565; Rep. de Pharm. 1896, 413.
- 1896: (7) HEGLAND, J. M. A. Separation and estimation of hydrastine.  
Nederl. Tijds. Pharm. **8**, 197; Chem. Centrbl. 1896, ii, 454.
- 1896: (8) ALLEN, A. H. Titration of quinine, and its relation to indicators.  
Chem. and Drug. 1896, 22; Analyst, **21**, 85; J. Chem. Soc. **70**, ii, 584; Chem. Centrbl. 1896, i, 1146.
- 1896: (9) KRAMERS, I. G. Estimation of quinine salts by means of sodium nitroprusside.  
Rec. Trav. chim., **15**, 138; J. Chem. Soc. **72**, ii, 83; Chem. Centrbl. 1896, ii, 602.
- 1896: (10) HOWARD, D. Estimation of quinine.  
Pharm. J. Trans. (4) **3**, 505; J. Chem. Soc. **72**, ii, 391.
- 1896: (11) VEDRÖDI, V. Estimation of nicotine and ammonia in tobacco.  
Ztschr. anal. Chem. **35**, 309; J. Chem. Soc. **70**, ii, 631; Chem. Centrbl. 1896, ii, 367.
- 1896: (12) LOOFF, G. Estimation of morphine in opium, gravimetric.  
J. Pharm. Chim. (6) **4**, 312; J. Chem. Soc. **72**, ii, 293; Chem. Centrbl. 1896, i, 1033; Rep. de Pharm. 1896, 310.
- 1896: (13) GUNN, A. Rapid and accurate estimation of total alkaloids in coca leaves.  
Pharm. J. Trans. (4) **3**, 249; J. Chem. Soc. **72**, ii, 294; Chem. Centrbl. 1896, ii, 1012; Rep. de Pharm. 1896, 512.
- 1896: (14) PUCKNER, W. A. Estimation of caffeine.  
J. Am. Chem. Soc. **18**, 978; J. Chem. Soc. **72**, ii, 390.
- 1896: (15) VAN LEDDEN-HULSEBOSCH. Estimation of caffeine in tea, gravimetric.  
Pharm. Centralh. **36**, 742; Chem. Centrbl. 1896, i, 332; J. Chem. Soc. **72**, ii, 83.

- 1896: (16) GANE. Caffeine in tea.  
J. Soc. Chem. Ind. **15**, 95; J. Chem. Soc. **74**, ii, 100; Chem. Centrbl. 1896, i, 873.
- 1896: (17) GOMBERG, M. Use of Wagner's reagent for estimation of caffeine.  
J. Am. Soc. **18**, 331 and 347; J. Chem. Soc. **72**, i, 129; Chem. Centrbl. 1896, i, 1075.
- 1896: (18) PETIT, A. and TERRAT, P. Estimation of caffeine in tea.  
J. Pharm. Chim. **3**, 529; J. Chem. Soc. **70**, ii, 629; Bull. Soc. Chim. Paris, **15**, 811; Chem. Centrbl. 1896, ii, 214.
- 1896: (19) DELACOUR. Estimation of caffeine in tea, coffee, etc. Gravimetric.  
J. Pharm. Chim. (6) **4**, 490; J. Chem. Soc. **72**, ii, 293.
- 1896: (20) EMINGER, A. Estimation of theobromine in cacao preparations.  
Forschungsber. Lebensm. **3**, 275; Chem. Centrbl. 1896, ii, 808.
- 1896: (21) DOTT, D. B. Opium assay.  
J. Soc. Chem. Ind. **15**, 91; J. Chem. Soc. **74**, ii, 100.
- 1896: (22) ————. Assay of ipecac.  
Pharm. Centralh., 1896, 763.
- 1896: (23) SMITH, C. E. Assay of nux vomica. Volumetric.  
Am. J. Pharm. **68**, No. 4; Apoth. Ztg. 1896, 386.
- 1896: (24) ESCHENBURG, H. Assay of cinchona.  
Apoth. Ztg. 1896, 147; Rep. de Pharm. 1896, 215.
- 1896: (25) GEORGES. Estimation of caffeine.  
J. Pharm. Chim. 1896, July 15; Rep. de Pharm. 1896, 401; Viertelj. Fortsch. Chem. Nahrungsm. **12**, 47; Ztschr. österr. Apot. V. 1897, 91; Chem. Centrbl. 1897, ii, 506.
- 1896: (26) KEBLER, L. F. Assay of cinchona, comparison of methods.  
Am. J. Pharm. 1896, 79; Rep. de Pharm. 1896, 508.

- 1896: (27) DUNSTAN and TICKLE. Estimation of aconitine. Decomposition, distillation, and titration of acetic acid formed.  
Pharm. J. Trans. (4) 2, 121; J. Chem. Soc. 72, ii, 294.
- 1897: (1) TASSILLY, E. Estimation of caffeine in coffee.  
Bull. Soc. Chim. (3) 17, 761; Chem. Centrbl. 1897, ii, 643.
- 1897: (2) FORSTER, A. and RIECHELMAN, R. Estimation of caffeine in coffee.  
Ztschr. f. öffentl. Chem. 3, 129; Chem. Centrbl. 1897, i, 1259.
- 1897: (3) KELLER, C. C. Estimation of caffeine in tea.  
Ber. d. pharm. Ges. 7, 105; Chem. Centrbl. 1897, i, 1134.
- 1897: (4) HILGER and JUCKENACK. Estimation of caffeine in tea and coffee.  
Apoth. Ztg. 12, 145 and 422; J. Pharm. Chim. (6) 6, 184 and 190; J. Chem. Soc. 72, ii, 611.
- 1897: (5) MAUPY, L. Estimation of theobromine in cocoa and chocolate.  
J. Pharm. Chim. (6) 5, 329; J. Chem. Soc. 72, ii, 531; Chem. Centrbl. 1897, i, 1077.
- 1897: (6) FARR, E. H. and WRIGHT, R. Estimation of alkaloids.  
Pharm. J. Trans. 1897, 58, 202; J. Chem. Soc. 74, 101.
- 1897: (7) FOUQUET, L. Separation of codeine and morphine, with anisole.  
J. Pharm. Chim. (6) 5, 49; J. Chem. Soc. 72, ii, 390; Chem. Centrbl. 1897, i, 342.
- 1897: (8) MONTEMARTINI and TRASCIATI. Estimation of morphine in opium.  
Gaz. chim. ital. 27, ii, 302; Chem. Centrbl. 1897, ii, 1165.
- 1897: (9) GRANDVAL and LAJOUX. Estimation of morphine in opium and opiates.  
J. Pharm. Chim. (6) 5, 153; J. Chem. Soc. 72, ii, 391.

1898: (1) WENTZKY, O. Assay of drugs. Resumé of processes since 1880.

Apoth.-Ztg. 13, pp. 6, 15, 29, 36, 43 and 75.

1898: (2) TRILLICH and GOECKEL. Estimation of caffeine.

Pharm. Ztg. 42, 53; Pharm. J. Trans. (4) 6, 316.

1898: (3) BRITISH PHARMACOPŒIA. Assay of belladonna.

Pharm. J. Trans. (4) 6, 368.

Shake 10 c. c. of the fluid extract with 10 c. c. of chloroform, 50 c. c. of water, and a decided excess of ammonia; separate the chloroformic solution and twice repeat the agitation and separation with chloroform. Shake the mixed chloroformic solutions with 5 c. c. of diluted sulphuric acid, mixed with twice its volume of warm water, separate, and repeat the operation. Wash the mixed acid liquids with 3 c. c. of chloroform, then agitate with 10 c. c. of chloroform and an excess of ammonia. Separate the chloroformic solution, twice repeat the agitation and separation with chloroform, wash the mixed chloroformic solutions with 5 c. c. of water containing 1 drop of solution of ammonia, evaporate on a water bath, dry the residue at 100° C., and weigh. Next dissolve the residue in 10 c. c. of decinormal hydrochloric acid, neutralize with centinormal sodium hydroxide, using tincture of cochineal as an indicator, deduct the measure of soda solution from 100 c. c., and multiply the remainder by .00287. The product will be the weight in grams of alkaloids present in 10 c. c. of fluid extract.

1898: (4) BRITISH PHARMACOPŒIA. Assay of cinchona, tincture or fluid extract.

Pharm. J. Trans. (4) 6, 368.

Place 5 c. c. of the fluid extract in a stoppered glass separator, together with 25 c. c. of water, 30 c. c. of benzolated amyl alcohol, and 15 c. c. of solution of potassium hydroxide. After shaking repeatedly, the lower dark-colored alkaline layer that separates is run off and again shaken with 30 c. c. of benzolated amyl alcohol, the lower layer that now separates being rejected. The two upper layers containing the alkaloids in solution are then mixed, washed with a little water, and agitated thoroughly with 30 c. c. of a warm mixture of 1 volume of diluted hydrochloric acid and 5 volumes of water. This operation is repeated with the alcoholic liquor that separates, and the two acid layers are subsequently mixed. The alkaloids are then shaken out with 10 c. c. of chloroform and sufficient solution of ammonia to impart a strongly alkaline reaction; the agitation and separation repeated with two successive



quantities of 10 c. c. of chloroform, the chloroformic liquids mixed, evaporated slowly, the residue dried at 110° C., and the residual alkaloïds weighed.

1898: (5) BRITISH PHARMACOPŒIA. Assay of ipecac, fluid extract.

Pharm. J. Trans. (4) 6, 369.

Dilute 20 c. c. of the extract with an equal bulk of water, remove the alcohol by the aid of a water bath, add to the warm solution an excess of solution of lead subacetate, filter, wash the precipitate with water, and add the washings to the filtrate. Remove excess of lead from filtrate by precipitation with dilute sulphuric acid, filter, wash, and add the washings to the filtrate. Transfer filtrate to a separator, add excess of solution of ammonia (s. g. 0.959), and agitate with 25 c. c. of chloroform. Separate and set aside the chloroformic solution, and twice repeat the agitation with chloroform and the subsequent separation. Then mix the chloroformic solutions, evaporate, dry at a temperature below 80° C., and weigh the residue of total alkaloids.

1898: (6) BRITISH PHARMACOPŒIA. Assay of nux vomica, fluid extract.

Pharm. J. Trans. (4) 6, 369.

Evaporate 10 c. c. to a thick syrup on a water-bath, dissolve the residue in 20 c. c. of water, heating if necessary, place the solution in a separator, and shake out with 5 gm. of sodium carbonate dissolved in 25 c. c. of water, together with 10 c. c. of chloroform. Separate, and twice repeat the operation with chloroform only. Then mix 6 c. c. of dilute sulphuric acid with 25 c. c. of water, divide the mixture into three parts, and shake the mixed chloroformic solutions with each in turn. Mix the acid liquids, dilute with water to 175 c. c. and shake well and frequently in a stoppered flask during half an hour with 25 c. c. of potassium ferrocyanide solution (10 gms. in 200 c. c.) The precipitate formed is transferred to a filter, the flask rinsed with water containing one-fortieth its volume of dilute sulphuric acid and the precipitate washed till the washings are free from bitterness. Next shake the precipitate in a separator with 5 c. c. of solution of ammonia, add 15 c. c. of chloroform in two successive solutions, evaporate in a counterpoised dish in a current of warm air, the dish being covered to avoid loss of strychnine. The residue is dried for 1 hour on a water-bath, and then weighed.

1898: (7) BRITISH PHARMACOPŒIA. Assay of opium tincture.  
Pharm. J. Trans. (4) 6, 380.

Evaporate 80 c. c. to 30 c. c. mixing the residue with 3 gm. of freshly-slaked lime, adding water to 85 c. c., and setting aside for half an hour, stirring occasionally. Next, filter off 50 c. c. of the tincture, add 5 c. c. of 90% alcohol and 30 c. c. of ether, shake, and add 2 gm. of ammonium chloride. Shake well and frequently during half an hour, set aside for 12 hours for the morphine to separate, filter, wash with morphinated water, and dry the crystals first by gentle pressure between filter paper, then at 55° or 60° C., and finally at 110° C. for two hours. Weigh the crystals, and titrate 0.3 gm. with decinormal solution of sulphuric acid until the liquid, after boiling, slightly reddens blue litmus paper. Add to the weight of anhydrous morphine, thus indicated, 0.05 gm. to cover loss.

## PART II.—INDEX OF AUTHORS.

- ADRIAN and GALLOIS. 1887. (16) Opium.  
ALESSANDRI. 1882. (5) Cinchona alkaloids.  
ALLEN, A. H. 1891. (8) Aconite.  
“ 1892. (2) Use of methyl orange.  
“ 1892. (3) Volumetric estimation.  
“ 1896. (8) Titration of quinine.  
ARNDT, E. M. 1890. (18) Emetine.  
  
BARTHE L. 1892. (5) Volumetric estimation.  
“ 1892. (6) Quinine and cinchona alkaloids.  
BARTHEL, G. and DIETERICH, E. 1888. (13) Morphine.  
BECK, C. R. 1892. (17) Ipecac.  
BECKURTS, H. 1887. (9) Morphine.  
“ 1889. (2) Strychnine, brucine, and general process.  
“ 1890. (5) Strychnine.  
“ 1891. (9) Nux vomica, aconite, belladonna and hyos-  
cyamus.  
“ 1894. (4) General process.  
“ 1896. (6) Hydrastine and berberine.  
“ and FRERICHS. 1896. (2) General process.  
“ “ HOLST. 1887. (25) Strychnine and brucine.  
“ “ “ 1887. (26) Strychnine, alkalimetric.  
“ “ SCHRAVT. 1887 (11) Morphine.  
BERNHARDT, W. 1884. (2) Morphine.  
BIEL, J. 1882. (6) Cinchona.

- BIEL, J. 1887. (21) Morphine.  
 " 1888. (8) Nicotine.
- BIÉTRIX and CAZENEUVE. 1892. (15) Caffeine.
- BIRD, F. C. J. 1887. (10) Morphine.  
 " 1892. (10) Emetine.
- BLUNT, T. P. 1889. (10) Emetine.  
 " 1890. (13) Emetine, alkalimetric.
- BLYTH, A. W. 1877. (1) Caffeine.  
 " 1881. (7) Quinine.
- BRAITHWAITE and FARR. 1886. (14) Morphine.
- BRUNNER, H. 1894. (8) Caffeine and theobromine.  
 " and LEINS, H. 1893. (10) Caffeine and theobromine.
- CANNEPIN and VAN ELJK. 1893. (12) Morphine.
- CARLES. 1872. (2) Quinine.
- CASPARI, C. and DOHME, R. L. 1893. (17) Assay by titration.
- CAVENDONI. 1888. (14) General process.
- CAZENEUVE and CAILLOL. 1877. (2) Piperine.  
 " " BIÉTRIX. 1892. (15) Caffeine.
- CHRISTIANSEN, A. 1890. (4) General process, iodometric.
- CLAASEN, E. 1890. (12) Codeine and morphine.
- COBLENTZ. 1885. (2) Atropine.
- CONROY, M. 1883. (4) Strychnine.  
 " 1884. (1) Morphine.
- COWNLEY and PAUL. 1887. (27) Caffeine in coffee.  
 " " " 1887. (28) Caffeine in tea.
- CRIPPS, R. A. 1887. (32) Coniine.  
 " 1895. (5) Emetine.  
 " and WITHBY, A. 1889. (17) Emetine.
- CUSHMAN, A. S. 1894. (11) Strychnine in corpses.
- DELACOUR. 1896. (19) Caffeine.
- DE VRIJ, J. E. 1880. (6) Quinine as herapathite.  
 " 1882. (2) General process for barks.  
 " 1882. (3) Quinine as herapathite.  
 " 1885. (6) Quinine as oxalate.  
 " 1885. (7) Cinchonidine in quinine sulphate.  
 " 1885. (8) Quinine in sulphate.  
 " 1885. (9) Cinchona alkaloids.  
 " 1886. (12) Quinine as chromate.
- DE WIJS, B. C. J. 1893. (11) Morphine.
- DIETERICH, E. 1885. (18) Morphine.

- DIETERICH, E. 1887. (12) Morphine.  
 “ 1887. (24) Strychnine.  
 “ 1889. (1) Morphine.  
 “ 1890. (10) Morphine.  
 “ 1890. (11) Morphine and emetine.  
 “ 1894. (5) Discussion of methods.  
 “ and BARTHEL. 1888. (13) Morphine.
- DOHME, R. L. and CASPARI, C. 1893. (17) Assay by titration.
- DOMERGUE and NICOLAS. 1892. (14) Caffeine.
- DOTT, D. B. 1894. (12) Morphine, alkalimetric.  
 “ 1896. (21) Opium assay.
- DRAGENDORFF. 1874. (2) Strychnine.  
 “ 1878. (4) Theobromine.
- DUNSTAN, W. R. and SHORT. 1883. (5) Strychnine.  
 “ “ 1883. (6) Strychnine and brucine.  
 “ and RANSOM. 1884. (7) Atropine.  
 “ “ 1885. (1) Atropine.  
 “ and TICKLE. 1896. (27) Aconitine.
- DWARS. 1878. (5) Quinine  
 ! 1879. (1) Quinine and strychnine.
- EGELING. 1890. (3) Strychnine.
- EMINGER, A. 1896. (20) Theobromine.  
 “ and HILGER. 1894. (9) Theobromine.
- ESCHENBURG, H. 1896. (24) Cinchona.
- EYKMAN. 1881. (8) Cinchona.
- FAIRTHORNE. 1882. (13) Cinchona.
- FARR, E. H. and BRAITHWAITE. 1886. (14) Morphine.  
 “ and WRIGHT, R. 1894. (2) Discussion of methods.  
 “ “ 1897. (6) Estimation of alkaloids.
- FAWSSETT. 1889. (13) Cinchona.
- FLEURY, G. 1879. (8) Morphine.
- FLUECKIGER, F. A. 1879. (9) Morphine.  
 “ 1885. (17) Morphine.  
 “ 1886. (17) Emetine.  
 “ 1889. (5) Morphine.
- FORSTER and RIECHELMANN. 1897. (2) Caffeine.
- FOUQUET, L. 1897. (7) Codeine and morphine.
- FRERICHS, G. and BECKURTS. 1896. (2) General process.
- GALLOIS and ADRIAN. 1887. (16) Morphine.
- GANE. 1896. (16) Caffeine.

- GEORGES. 1896. (25) Caffeine.  
 GEROCK, J. E. 1889. (6) Brucine and strychnine.  
 GERRARD, A. W. 1882. (12) Atropine.  
 GOEBEL, H. 1887. (17) Morphine.  
 " 1889. (3) Quinine and cinchonidine.  
 GÖCKEL and TRILLICH. 1898. (2) Caffeine.  
 GOMBERG, M. 1896. (17) Caffeine.  
 GRANDVAL, A., and LAJOUX, H. 1893. (6) Caffeine.  
 " " 1893. (18) General process.  
 " " 1893. (19) Emetine, cocaine, and cin-  
 chona.  
 " " 1897. (9) Morphine.  
 GROVES, T. B. 1864. General process.  
 GUILLOT. 1892. (16) Caffeine.  
 GUNN, A. 1896. (13) Coca alkaloids.  
  
 HAGER. 1880. (4) Cinchona alkaloids.  
 " 1881. (2) Alkaloids as picrates.  
 " 1883. (1) Method of extraction.  
 HAUBENSAK, W. 1891. (4) Cinchona.  
 HEGLAND, J. M. A. 1896. (7) Hydrastine.  
 HEINTZ, E. 1878. (2) Method of extraction.  
 HERETH, F. S. 1886. (1) Method of titration.  
 HESSE, O. 1880. (7) Quinine.  
 " 1881. (6) Cinchonidine.  
 HEUT, G. 1893. (13) Coniine and nicotine.  
 HILBIG, C. 1880. (5) Cinchona.  
 HILGER and EMINGER. 1894. (9) Theobromine.  
 " " JUCKENACK. 1897. (4) Caffeine.  
 HINSDALE, S. J. 1890. (4) Morphine.  
 HOLDERMAN. 1889. (9) Morphine, iodometric.  
 HOLST and BECKURTS. 1887. (25) Strychnine and brucine.  
 " " 1887. (26) Strychnine.  
 HOOPER, D. 1886. (10) Quinine.  
 HOWARD, D. 1896. (10) Quinine.  
  
 JANDONS, A. 1887. (20) Morphine.  
 JOHNSTONE, W. 1889. (11) Piperine.  
 JONES, H. W. 1886. (16) Emetine.  
 JUCKENACK and HILGER. 1897. (4) Caffeine.  
  
 KASPAR. 1886. (11) Cinchona.  
 KEBLER, L. F. 1895. (1) Morphine.

- KEBLER, L. F. 1896. (26) Cinchona.  
 " and LA WALL. 1895. (7) Volumetric process.
- KELLER, C. C. 1892. (12) Emetine.  
 " 1893. (1) Cinchona.  
 " 1893. (2) Emetine.  
 " 1893. (3) Emetine.  
 " 1893. (4) Strychnine.  
 " 1893. (7) Strychnine.  
 " 1894. (1) General process.  
 " 1894. (13) Hydrastine.  
 " 1895. (9) Strychnine.  
 " 1895. (11) Cocaine.  
 " 1896. (3) General process.  
 " 1897. (3) Caffeine.
- KERNER, G. 1880. (8) Quinine.
- KIPPENBERGER, K. 1896. (4) Separation of alkaloids from ptomaines.  
 " 1896. (5) General iodometric process.
- KISSEL. 1882. (4) Cinchona.
- KISSLING, R. 1882. (10) Nicotine.  
 " 1882. (11) Nicotine.  
 " 1883. (2) Nicotine.  
 " 1889. (9) Nicotine, polariscopic.  
 " 1895. (3) Nicotine and ammonia.
- KOTTMEYER, G. 1892. (11) Emetine.
- KRAMERS, I. G. 1896. (9) Quinine.
- KREMEL, A. 1887. (3) Aconite.  
 " 1887. (14) Morphine.  
 " 1888. (1) Emetine, hydrastine and berberine.  
 " 1888. (2) Strychnine.  
 " 1888. (7) Caffeine.  
 " 1891. (10) Coniine.
- KUNZE, W. E. 1894. (6) Caffeine and theobromine.
- KUERSTEINER, C. 1892. (8) Cinchona.
- LAJOUX, H. and GRANDVAL, A. 1893. (6) Caffeine.  
 " " 1893. (18) General process.  
 " " 1893. (19) Emetine, cocaine, and cin-  
 chona.  
 " " 1897. (9) Morphine.
- LALIEN. 1885. (15) Morphine.
- LAMBERT, A. 1891. (1) Morphine.
- LANDRIN. 1889 (18) Cinchona.

- LANGLOIS and PORTES. 1881. (1) Morphine.
- LA WALL and KEBLER. 1895. (7) Comparison of indicators.
- LEINS, H. 1894. (14) Caffeine and theobromine.
- “ and BRUNNER, H. 1893. (10) Caffeine and theobromine.
- LENZ, W. 1888. (4) Quinine, comparison of methods.
- LIEUNNIGH, F. 1893. (22) General process.
- LIGNON. 1887. (33) Emetine.
- LILLY. 1887. (2) Atropine, and general process.
- LLOYD, J. U. 1885. (11) Berberine.
- “ 1891. (2) General process.
- LOOFF, G. 1890. (7) Morphine.
- “ 1890. (8) Morphine.
- “ 1896. (12) Morphine.
- LCESCH, A. 1879. (5) General process.
- “ 1879. (6) General process.
- “ 1887. (31) Caffeine.
- LYNN, J. 1876. (3) Morphine.
- LYONS, A. B. 1884. (6) General process.
- “ 1885. (12) Emetine.
- “ 1885. (13) Strychnine.
- “ 1886. (3) Berberine.
- “ 1886. (4) Hydrastine.
- “ 1886. (5) Physostigmine.
- “ 1886. (6) Strychnine.
- “ 1886. (7) Discussion of methods.
- “ 1886. (8) Cocaine.
- “ 1886. (9) Cinchona.
- MACEWEN. 1887. (34) Physostigmine.
- MARKOWNIKOFF, B W. 1876. (2) Caffeine.
- MASING, E. 1877. (3) Veratrine and physostigmine.
- MAUPY, L. 1897. (5) Theobromine.
- MEINERT and WRAMPELMEIER. 1886. (18) Opium.
- MENDINI, A. 1895. (10) Emetine.
- MEYER, H. 1882. (7) Cinchona.
- MEYER, A. 1893. (5) Emetine.
- MONTEMARTINI and TRASCIATTI. 1897. (8) Morphine.
- MYLIUS, E. 1880. (9) Opium alkaloids.
- “ 1881. (10) Morphine.
- NAGELVOORT, J. B. 1890. (22) Morphine.
- NEUMAN, S. 1889. (14) Quinine.

- NICHOLS, H. T., and NORTON, T. H. 1892. (1) Examination of Lloyd's method.
- NICOLAS and DOMERGUE. 1892. (14) Caffeine.
- PARTHEIL, A. 1892. (4) Iodo-eosin as indicator.
- PATRONOUILLARD. 1880. (3) Caffeine.
- PAUL, B. H. 1891. (6) Caffeine.
- “ and COWNLEY. 1887. (27) Caffeine in coffee.
- “ “ 1887. (28) Caffeine in tea.
- PETIT, A. 1879. (7) Morphine.
- “ and TERRAT, P. 1896. (18) Caffeine.
- PEZZOLATO, A. 1891 (7) Nicotine.
- PHARMACOPŒIA, British. 1885. (19) Cinchona.
- “ “ 1885. (20) Nux vomica.
- “ “ 1885. (21) Opium.
- “ “ 1898. (3) Belladonna.
- “ “ 1898. (4) Cinchona.
- “ “ 1898. (5) Ipecac.
- “ “ 1898. (6) Nux vomica.
- “ “ 1898. (7) Opium.
- “ U. S. 1890. (19) Cinchona.
- “ “ 1890. (20) Nux vomica.
- “ “ 1890. (21) Opium.
- “ German. 1895. (12) Cinchona.
- “ “ 1895. (13) Opium.
- PINETTE, J. 1892. (9) Nicotine.
- PLUGGE, P. C. 1887. (1) General volumetric process.
- “ 1887. (13) Separation of opium alkaloids.
- POEHL. 1881. (9) Pilocarpine.
- POPOVICI, M. 1889. (8) Nicotine.
- PORTES and LANGLOIS. 1881. (1) Morphine.
- PRESCOTT, A. B. 1877. (5) Quinine.
- “ 1879. (11) Morphine.
- “ 1880. (1) General process.
- “ 1896. (1) Periodides.
- “ and THUM. 1878. (1) Cinchona, comparison of methods.
- PROLLIUS. 1881. (4) Cinchona.
- PRUNIER, L. 1879. (3) Cinchona.
- PUCKNER, W. A. 1896. (14) Caffeine.
- RANSOM, F. 1887. (5) Emetine.
- “ and DUNSTAN. 1884. (7) Belladonna.



- RANSOM, F. and DUNSTAN. 1885. (1) Belladonna.  
 RANWEZ. 1893. (16) Aconite, hyoscyamus and cicuta.  
 REDWORD. 1884. (8) Belladonna.  
 RICHMOND and SEATON. 1890. (6) Quinine.  
 RIECHELMANN and FORSTER. 1897. (2) Caffeine.  
 ROTHER. 1880. (11) Opium.  
 RUDDIMAN, E. A. 1887. (22) Quinine.  
  
 SARNOW and SCHWEISSINGER. 1890. (2) General process.  
 SCHACHT, C. 1881. (5) Quinine.  
 SCHLÄFER, L. 1887. (23) Cinchonidine.  
 SCHEFFER, E. 1885. (5) Nicotine.  
 SCHEIBE, E. 1883. (7) Morphine.  
 SCHEIBLER. 1886. (2) General process.  
 SCHLICKUM, O. 1887. (15) Morphine.  
 SCHMIDT, J. H. 1892. (7) Quinine.  
 SCHMITT, E. 1880. (10) Opium.  
 SCHRAGE, F. 1878. (3) Cinchona.  
 SCHRAUT, A., and BECKURTS, H. 1887. (11) Morphine.  
 SCHWEISSINGER. 1885. (14) Strychnine and brucine.  
     "    and SARNOW. 1890. (2) General process.  
 SCHWICKERATH, K. 1894. (3) General process.  
 SEATON and RICHMOND. 1890. (6) Quinine.  
 SHIMOYANO, Y. 1883. (3) Cinchona.  
     "    1885. (10) Cinchona.  
 SHORT and DUNSTAN. 1883. (5) Strychnine.  
     "    "    1883. (6) Strychnine and brucine.  
 SKALWEIT, J. 1881. (3) Nicotine.  
     "    1882. (9) Nicotine.  
 SMITH, E. D. 1887. (29) Caffeine.  
 SMITH, C. E. 1896. (23) Nux vomica.  
 SMITH, J. D., and TESCHEMACHER, E. F. 1888. (11) Morphine.  
 SNOW, H. W. 1886. (13) Caffeine.  
     "    1887. (6) Emetine.  
     "    1888. (5) General process, Mayer's reagent.  
     "    1888. (6) General process, phospho-molybdic acid.  
     "    1889. (7) Strychnine and brucine.  
 SOKOLOFF, N. 1876. (1) Caffeine.  
     "    1893. (8) Caffeine.  
     "    1895. (4) Caffeine.  
 SPENCER, G. L. 1890. (15) Caffeine.  
     "    1891. (5) Caffeine.

- SQUIBB, E. R. 1882. (1) Morphine.  
 " 1882. (8) Cinchona.  
 " 1884. (4) Cocaine.  
 " 1884. (5) Caffeine.  
 " 1885. (3) Atropine.  
 " 1885. (4) Cocaine.  
 " 1887. (4) Cocaine.  
 " 1887. (19) Morphine.  
 " 1888. (9) Cocaine.  
 " 1888. (10) Morphine.
- STILWELL, C. M. 1887. (8) Morphine.
- STEDER, W. 1894. (10) Cortex granati.
- SUESS, P. 1890. (16) Theobromine.  
 " 1893. (9) Theobromine.
- TASSILLY, E. 1897. (1) Caffeine.
- TERRAT, P., and PETIT, A. 1896. (18) Caffeine.
- TESCHEMACHER, E. F. 1877. (4) Morphine.  
 " and SMITH, J. D. 1888. (11) Morphine.
- THIEL, J. 1894. (7) Theobromine.
- THOMPSON, F. A. 1890. (17) Coniine.  
 " 1891. (3) General process.  
 " 1893. (15) Hydrastine and berberine.
- THRESH, J. C. 1880. (2) Volumetric method.
- THUM and PRESCOTT. 1878 (9) Cinchona.
- TICKLE and DUNSTAN. 1896. (27) Aconitine.
- TRACCIATI and MONTEMARTINI. 1897. (8) Morphine.
- TRAUB. 1892. (13) General iodometric process.
- TRILLICH and GÖCKEL. 1898. (2) Caffeine.
- UMNEY, J. C. 1895. (6) Aconitine.
- VAN DER MARK. 1889. (12) Cocaine.  
 " 1889. (16) Strychnine.
- VAN ELK and CANNEPIN. 1893. (12) Morphine.
- VAN ITALLIE. 1888. (15) Atropine.  
 " 1889. (15) Alkalimetric process.  
 " 1890. (1) Alkalimetric process.  
 " 1895. (14) General process.
- VAN LEDDEN-HULSEBOSCH. 1893. (21) General process.  
 " " 1895. (8) General process.  
 " " 1896. (15) Caffeine.

- VAN ZWALUWENBERG, A. 1887. (7) Bibliography of opium assay.  
 VEDRČEDI, V. 1893. (14) Nicotine, comparison of methods.  
 " 1895. (2) Nicotine and ammonia.  
 " 1896. (11) Nicotine and ammonia.  
 VENTURINI, V. 1886. (15) Morphine, comparison of methods.  
 VITALI, D. 1893. (20) General process for chlorides and sulphates.  
 VITÉ, F. 1890. (14) Caffeine.  
 VON PERGER. 1884. (3) Morphine.
- WAAGE, TH. 1887. (30) Caffeine.  
 WAGNER. 1861. Periodide method.  
 WAINWRIGHT. 1885. (16) Morphine.  
 WENTZKY, O. 1898. (1) Resumé of processes.  
 WIEDA. 1888. (3) Cinchona, resumé of 25 processes.  
 WILEY, C. D. 1887. (18) Morphine.  
 WILLIAMS, R. 1888. (12) Morphine.  
 WITHBY, A., and CRIPPS, R. A. 1889. (17) Emetine.  
 WOLFRAM, G. 1879. (4) Theobromine.  
 WRAMPELMEIER and MEINERT. 1886. (18) Opium.  
 WRIGHT, R., and FARR, E. H. 1894. (2) Discussion of methods.  
 " " 1897. (6) General process.
- YVON. 1879. (2) Morphine.
- ZINOFFSKY. 1872. (1) General volumetric method.  
 " 1874. (1) Coniine.

---

 PART III.—SUBJECT INDEX.

- ACONITINE. 1887. (3) Kremel.  
 " '91. (8) Allen.  
 " '95. (6) Umney.  
 " '96. (27) Dunstan and Tickle.
- ATROPINE. 1882. (12) Gerrard.  
 " '84. (7) Dunstan and Ransom.  
 " '84. (8) Redword.  
 " '85. (1) Dunstan and Ransom.  
 " '85. (2) Coblenz.  
 " '85. (3) Squibb.  
 " '88. (15) Van Itallie.  
 " '98. (3) British Pharmacopolia.

BERBERINE.	1885.	(11)	Lloyd.
"	'86.	(3)	Lyons.
"	'88.	(1)	Kremel.
"	'93.	(15)	Thompson.
"	'96.	(6)	Beckurts.
BRUCINE.	1883.	(6)	Dunstan and Short.
"	'85.	(14)	Schweissinger.
"	'87.	(25)	Beckurts and Holst.
"	'89.	(2)	Beckurts.
"	'89.	(6)	Gerock.
"	'89.	(7)	Snow.
"			See also strychnine.
CAFFEINE.	1876.	(2)	Markownikoff.
"	'77.	(1)	Blyth.
"	'80.	(3)	Patronouillard.
"	'84.	(5)	Squibb.
"	'86.	(13)	Snow.
"	'87.	(27)	Cownley and Paul.
"	'87.	(28)	Cownley and Paul.
"	'87.	(29)	E. D. Smith.
"	'87.	(30)	Waage.
"	'87.	(31)	Löesch.
"	'88.	(7)	Kremel.
"	'90.	(14)	Vitè.
"	'90.	(15)	Spencer.
"	'91.	(5)	Spencer.
"	'91.	(6)	Paul.
"	'92.	(14)	Domergue and Nicolas.
"	'92.	(15)	Cazeneuve and Biètrix.
"	'92.	(16)	Guillot.
"	'93.	(6)	Grandval and Lajoux.
"	'93.	(8)	Sokoloff.
"	'93.	(10)	Brunner and Leins.
"	'94.	(6)	Kunze.
"	'94.	(8)	Brunner.
"	'94.	(14)	Leins, A.
"	'95.	(4)	Sokoloff.
"	'96.	(14)	Puckner.
"	'96.	(15)	Van Ledden-Hulsebosch.
"	'96.	(16)	Gane.
"	'96.	(17)	Gomberg.

CAFFEINE.	1896.	(18)	Petit and Terrat.
"	'96.	(19)	Delacour.
"	'96.	(25)	Georges.
"	'97.	(1)	Tassilly.
"	'97.	(2)	Forster and Riechelmann.
"	'97.	(3)	Keller.
"	'97.	(4)	Hilger and Juckenack.
"	'98.	(2)	Trillich and Gœckel.

CEPHAELIN, see IPECAC.

CINCHONA ALKALOIDS:

CINCHONIDINE.	1881.	(6)	Hesse.
"	'85.	(7)	De Vrij.
"	'87.	(23)	Schæfer.
"	'89.	(3)	Gœbel.
QUININE.	1872.	(2)	Carles.
"	'77.	(5)	Prescott.
"	'78.	(5)	Dwars.
"	'79.	(1)	Dwars.
"	'79.	(10)	—
"	'80.	(6)	De Vrij.
"	'80.	(7)	Hesse.
"	'80.	(8)	Kerner.
"	'81.	(7)	Blyth.
"	'82.	(3)	De Vrij.
"	'85.	(6)	De Vrij.
"	'85.	(8)	De Vrij.
"	'85.	(19)	British Pharmacopœia.
"	'86.	(10)	Hooper.
"	'86.	(12)	De Vrij.
"	'87.	(22)	Ruddiman.
"	'88.	(4)	Lenz.
"	'89.	(3)	Gœbel.
"	'89.	(14)	Neumann.
"	'90.	(6)	Seaton and Richmond.
"	'90.	(19)	U. S. Pharmacopœia.
"	'92.	(7)	J. H. Schmidt.
"	'95.	(12)	German Pharmacopœia.
"	'96.	(8)	Allen.
"	'96.	(9)	Kramers.
"	'96.	(10)	Howard.
TOTAL ALKALOIDS.	1878.	(1)	Prescott and Thum.
"	'78.	(3)	Schrage.

## CINCHONA, Continued:

TOTAL ALKALOIDS.	1879.	(3)	Prunier.
"	'80.	(4)	Hager.
"	'80.	(5)	Hilbig.
"	'81.	(4)	Prollius.
"	'81.	(5)	Schacht.
"	'81.	(8)	Eykman.
"	'82.	(2)	De Vrij.
"	'82.	(4)	Kissel.
"	'82.	(5)	Alessandri.
"	'82.	(6)	Biel.
"	'82.	(7)	H. Meyer.
"	'82.	(8)	Squibb.
"	'82.	(13)	Fairthorne.
"	'83.	(3)	Shimoyana.
"	'85.	(9)	De Vrij.
"	'85.	(10)	Shimoyana.
"	'86.	(9)	Lyons.
"	'86.	(11)	Kaspar.
"	'88.	(3)	Wieda.
"	'89.	(13)	Fawsett.
"	'89.	(18)	Landrin.
"	'91.	(4)	Haubensak.
"	'92.	(6)	Barthe.
"	'92.	(8)	Kuersteiner.
"	'93.	(1)	Keller.
"	'96.	(24)	Eschenburg.
"	'96.	(26)	Kebler.
"	'98.	(4)	British Pharmacopœia.
COCAINE.	1884.	(4)	Squibb.
"	'85.	(4)	Squibb.
"	'86.	(8)	Lyons.
"	'87.	(4)	Squibb.
"	'88.	(9)	Squibb.
"	'89.	(12)	Van der Mark.
"	'95.	(11)	Keller.
"	'96.	(13)	Gunn.
CODEINE.	1887.	(13)	Plugge.
"	'90.	(12)	Claasen.
"	'97.	(7)	Fouquet.
CONIINE.	1874.	(1)	Zinoffsky.
"	'76.	(1)	Sokoloff.
"	'87.	(32)	Cripps.

- CONINE 1890. (17) Thompson.  
 " '91. (10) Kremel.  
 " '93. (13) Heut.

EMETINE, see IPECAC.

ESERINE, see PHYSOSTIGMINE.

- GENERAL PROCESSES. 1861. Wagner.  
 " '64. Groves.  
 " '78. (2) Heintz.  
 " '79. (5) Lœsch.  
 " '79. (6) Lœsch.  
 " '80. (1) Prescott.  
 " '81. (2) Hager.  
 " '83. (1) Hager.  
 " '84. (6) Lyons.  
 " '86. (2) Scheibler.  
 " '87. (2) Lilly.  
 " '88. (5) Snow.  
 " '88. (14) Cavendoni.  
 " '89. (2) Beckurts.  
 " '91. (2) Lloyd.  
 " '91. (3) Thompson.  
 " '91. (9) Beckurts.  
 " '92. (1) Nichols and Norton.  
 " '93. (16) Ranwez.  
 " '93. (18) Grandval and Lajoux.  
 " '93. (19) Grandval and Lajoux.  
 " '93. (21) Van Ledden-Hulsebosch.  
 " '93. (22) Lieunnigh.  
 " '94. (1) Keller.  
 " '94. (2) Farr and Wright.  
 " '94. (3) Schwickerath.  
 " '94. (4) Beckurts.  
 " '94. (5) Dieterich.  
 " '95. (8) Van Ledden-Hulsebosch.  
 " '95. (14) Van Itallie.  
 " '96. (1) Prescott.  
 " '96. (2) Beckurts and Frerichs.  
 " '96. (3) Keller.  
 " '96. (4) Kippenberger.  
 " '97. (6) Farr and Wright.  
 " '98. (1) Wentzky.

GENERAL PROCESSES, VOLUMETRIC.	'72.	(1)	Zinoffsky.
"	"	'80.	(2) Thresh.
"	"	'86.	(1) Hereth.
"	"	'86.	(7) Lyons.
"	"	'87.	(1) Plugge.
"	"	'88.	(6) Snow.
"	"	'89.	(15) Van Itallie.
"	"	'90.	(1) Van Itallie.
"	"	'90.	(2) Schweissing and Sarnow.
"	"	'90.	(4) Christiansen.
"	"	'92.	(2) Allen.
"	"	'92.	(3) Allen.
"	"	'92.	(4) Partheil.
"	"	'92.	(5) Barthe.
"	"	'92.	(13) Traub.
"	"	'93.	(7) Keller.
"	"	'93.	(17) Caspari and Dohme.
"	"	'93.	(20) Vitali.
"	"	'95.	(7) Kebler and LaWall.
"	"	'96.	(5) Kippenberger.

HYDRASTINE.	1886.	(4)	Lyons.
"	'88.	(1)	Kremel.
"	'93.	(15)	Thompson.
"	'94.	(13)	Keller.
"	'96.	(6)	Beckurts.
"	'96.	(7)	Hegland.

HYOSCYAMINE, see ATROPINE.

IPECAC.	1885.	(12)	Lyons.
"	'86.	(16)	H. W. Jones.
"	'86.	(17)	Flückiger.
"	'87.	(5)	Ransom.
"	'87.	(6)	Snow.
"	'87.	(33)	Lignon.
"	'88.	(1)	Kremel.
"	'89.	(10)	Blunt.
"	'89.	(17)	Cripps and Withby.
"	'90.	(11)	Dieterich.
"	'90.	(13)	Blunt.
"	'90.	(18)	Arndt.
"	'92.	(10)	Bird.



IPECAC.	1892.	(11)	Kottmayer.
"	'92.	(12)	Keller.
"	'92.	(17)	Beck.
"	'93.	(2)	Keller.
"	'93.	(3)	Keller.
"	'93.	(5)	A. Meyer.
"	'95.	(5)	Cripps.
"	'95.	(10)	Mendini.
"	'96.	(22)	—
"	'98.	(5)	British Pharmacopœia.
MORPHINE	1877.	(4)	Teschemacher.
"	'76.	(3)	Lynn.
"	'79.	(2)	Yvon.
"	'79.	(7)	Petit.
"	'79.	(8)	Fleury.
"	'79.	(9)	Flückiger.
"	'79.	(11)	Prescott.
"	'80.	(9)	Mylius.
"	'80.	(10)	E. Schmitt.
"	'80.	(11)	Rother.
"	'81.	(1)	Portes and Langlois.
"	'81.	(10)	Mylius.
"	'82.	(1)	Squibb.
"	'83.	(7)	Scheibe.
"	'84.	(1)	Conroy.
"	'84.	(2)	Bernhardt.
"	'84.	(3)	Von Perger.
"	'85.	(15)	Lalien.
"	'85.	(16)	Wainwright.
"	'85.	(17)	Flückiger.
"	'85.	(18)	Dieterich.
"	'85.	(21)	British Pharmacopœia.
"	'86.	(14)	Braithwaite and Farr.
"	'86.	(15)	Venturini.
"	'86.	(18)	Wrampelmeier and Meinert.
"	'87.	(7)	Van Zwaluwenberg.
"	'87.	(8)	Stillwell.
"	'87.	(9)	Beckurts.
"	'87.	(10)	Bird.
"	'87.	(11)	Beckurts and Schraut.
"	'87.	(12)	Dietrich.

- MORPHINE. 1887. (13) Plugge.  
 " '87. (14) Kremel.  
 " '87. (15) Schlickum.  
 " '87. (16) Adrian and Gallois.  
 " '87. (17) Gœbel.  
 " '87. (18) C. D. Wiley.  
 " '87. (19) Squibb.  
 " '87. (20) Jandons.  
 " '87. (21) Biel.  
 " '88. (10) Squibb.  
 " '88. (11) Teschemacher and Smith.  
 " '88. (12) Williams.  
 " '88. (13) Dietrich and Barthel.  
 " '89. (1) Dietrich.  
 " '89. (4) Holderman.  
 " '89. (5) Flückiger.  
 " '90. (7) Loeff.  
 " '90. (8) Loeff.  
 " '90. (9) Hinsdale.  
 " '90. (10) Dietrich.  
 " '90. (11) Dietrich.  
 " '90. (12) Claasen.  
 " '90. (21) U. S. Pharmacopœia.  
 " '90. (22) Nagelvoort.  
 " '91. (1) Lambert.  
 " '93. (11) de Wijs.  
 " '93. (12) Cannepin and Van Eijk.  
 " '94. (12) Dott.  
 " '95. (1) Kebler.  
 " '95. (13) German Pharmacopœia  
 " '96. (12) Loeff.  
 " '96. (21) Dott.  
 " '97. (7) Fouquet.  
 " '97. (8) Montemartini and Trasciatti.  
 " '97. (9) Grandval and Lajoux.  
 " '98. (7) British Pharmacopœia.
- NARCOTINE. 1897. (13) Plugge.
- NICOTINE. 1881. (3) Skalweit.  
 " '82. (9) Skalweit.  
 " '82. (10) Kissling.  
 " '82. (11) Kissling.

- NICOTINE. 1883. (2) Kissling.  
 " '85. (5) Scheffer.  
 " '88. (8) Biel.  
 " '89. (8) Popovići.  
 " '89. (9) Kissling.  
 " '91. (7) Pezzolato.  
 " '92. (9) Pinette.  
 " '93. (13) Heut.  
 " '93. (14) Vedrödi.  
 " '95. (2) Vedrödi.  
 " '95. (3) Kissling.  
 " '96. (11) Vedrödi.
- OPIUM ALKALOIDS, separation. 1887. (13) Plugge.
- PHYSOSTIGMINE. 1877. (3) Masing.  
 " '86. (5) Lyons.  
 " '87. (34) MacEwan.
- PILOCARPINE. 1881. (9) Pöehl.
- PIPERINE. 1877. (2) Cazeneuve and Caillol.  
 " '89. (11) Johnstone.
- POMEGRANATE (cortex granati). 1894. (10) Stöder.
- QUININE, see CINCHONA.
- STRYCHNINE. 1874. (2) Dragendorff.  
 " '79. (1) Dwars.  
 " '83. (4) Conroy.  
 " '83. (5) Dunstan and Short.  
 " '83. (6) Dunstan and Short.  
 " '85. (13) Lyons.  
 " '85. (14) Schweissinger.  
 " '85. (20) British Pharmacopœia.  
 " '86. (6) Lyons.  
 " '87. (24) Dieterich.  
 " '87. (25) Beckurts and Holst.  
 " '87. (26) Beckurts and Holst.  
 " '88. (2) Kremel.  
 " '89. (2) Beckurts.  
 " '89. (6) Gerock.  
 " '89. (7) Snow.  
 " '89. (16) Van der Mark.

STRYCHNINE.	1890.	(3)	Egeling.
"	'90.	(4)	Keller.
"	'90.	(5)	Beckurts.
"	'90.	(20)	U. S. Pharmacopœia.
"	'93.	(4)	Keller.
"	'94.	(11)	Cushman.
"	'95.	(9)	Keller.
"	'96.	(23)	C. E. Smith.
"	'98.	(6)	British Pharmacopœia.

THEBAINE, see OPIUM.

THEINE, see CAFFEINE.

THEOBROMINE.	1878.	(4)	Dragendorff.
"	'79.	(4)	Wolfram.
"	'90.	(16)	Süss.
"	'93.	(9)	Süss.
"	'93.	(10)	Brunner and Leins.
"	'94.	(6)	Kunze.
"	'94.	(7)	Thiel.
"	'94.	(8)	Brunner.
"	'94.	(9)	Hilger and Eminger.
"	'94.	(14)	Leins.
"	'96.	(20)	Eminger.
"	'97.	(5)	Maupy.

VERATRINE. 1877. (3) Masing.



UNIVERSITY OF CALIFORNIA LIBRARY  
BERKELEY

Return to desk from which borrowed.  
This book is DUE on the last date stamped below.

5 Aug 58

3 1953

19 Jul '60 JC

REC'D LD

JUL 27 1960

JUL 22 1970 82

REC'D LD JUL 28 70-1 PM 821

Due end of FALL Quarter  
subject to recall after — DEC 15 '70 8

REC'D LD JUL 7 70-1 PM 421

QD421  
M8

~~QD~~

237438

~~QD~~

