

Praise 10 18 vol 10 1870 Aug 10 1870


RB59278



Library
of the
University of Toronto



Digitized by the Internet Archive
in 2017 with funding from
University of Toronto



THE
ANIMAL KINGDOM,
ARRANGED ACCORDING TO ITS ORGANIZATION,
Serving as a
Foundation for the
NATURAL HISTORY OF ANIMALS,
and an
Introduction to Comparative Anatomy.
BY
BARON CUVIER,

Great Officer of the Legion of Honour, Commissor of State, &c. Member of the Royal Council of Public Instruction, One of the Forty of the French Academy, Perpetual Secretary to the Academy of Sciences, Member of the Academies & Royal Societies of London, Berlin, Pittsburgh, Stockholm, Copenhagen, &c. Gottingen, Bonn, &c. Muenchen, the Netherlands, Calcutta & of the Indian, Society of London, &c. &c.

WITH FIGURES DESIGNED AFTER NATURE:
the
Crustacea, Arachnides & Insecta,
by
M. Latreille.

Member of the Legion of Honour, Member of the Institute, Royal Academy of Sciences & of the greater portion of other learned Societies in Europe, America, &c.

Translated from the latest French Edition.

(with)
ADDITIONAL NOTES,
and
Illustrated by nearly 800 Coloured Plates.

IN FOUR VOLUMES.

VOL. III.

REPTILES - FISHES.

LONDON.

G. Henderson, 2 Old Bailey, Ludgate Hill.

AND SOLD BY ALL BOOKSELLERS.

1837.



TABLE OF CONTENTS TO PLATES.

VOLUME II.

REPTILIA—PISCES.

		Vol. II. Page
	REPTILIA.	
Plate 1. Fig. 1.— <i>HEMIDACTYLUS MARGINATUS</i> , Cuv.	.	36
Fig. 2.— <i>UROPELTIS CEYLANICUS</i> , Nob.	.	50
Fig. 3.— <i>DACTYLETHRA CAPENSIS</i> (<i>The Thimble or Cape Frog</i>)	.	69
Plate 2. Fig. 1.— <i>TESTUDO DEPRESSA</i> , Cuv.	.	6
Fig. 2.—Head of the <i>TESTUDO CLAUSA</i>	.	6
Fig. 3.— <i>EMYS CONCINNA</i> , Le Conte. <i>Taken from a Young Specimen</i>	.	6
Fig. 4.— <i>CHELONIA VIRGATA</i> , Nob.	.	8
Fig. 5.— <i>CHELYS FIMBRIA</i> , Gm. (<i>The Fringed Tortoise</i>)	.	9
Fig. 6.— <i>TRYONIX GANGETICUS</i> , Duv. (<i>The Tysre of the Ganges</i>)	.	10
Fig. 7.— <i>TRYONIX AEGYPTIACUS</i> , Geoff. (<i>The Soft-shelled Tortoise of the Nile</i>)	.	9
Plate 3. Fig. 1.— <i>CROCODILUS BIPORCATUS</i> , Cuv. (<i>The Double-crested Crocodile</i>)	.	14
Fig. 2.—Head of <i>ALLIGATOR SCLEROPS</i> , Cuv. (<i>The Spectacle Alligator</i>)	.	15
Fig. 3.—Head of <i>GAVIALIS TENUIROSTRIS</i> , Cuv.	.	13
Plate 4. Fig. 1.— <i>TESTUDO SERRATA</i> , Daud.	.	6
Fig. 2.— <i>TESTUDO MYDAS</i> , Lin. (<i>The Common Turtle</i>)	.	8
Fig. 3.— <i>TESTUDO CEPEDIANA</i> , Daud.*	.	8
Fig. 4.— <i>TESTUDO CARETTA</i> , Gm.	.	8
Fig. 5.— <i>TESTUDO CORIACEA</i> , Lin. (<i>The Coriacous Turtle</i>)	.	9
Fig. 6.— <i>TESTUDO FEROX</i> , Penn. (<i>The Soft-shelled Tortoise of America</i>)	.	9
Plate 5. Fig. 1.— <i>TESTUDO FIMBRIA</i> , Gm.	.	9
Fig. 2.— <i>TESTUDO SERPENTINA</i> , Lin. (<i>The Snapper</i>)	.	7
Fig. 3.— <i>TESTUDO LUTEOLA</i> , Daud.†	.	
Fig. 4.— <i>TESTUDO STELLATA</i> , Schœpf.	.	6
Fig. 5.— <i>TESTUDO GEOMETRICA</i> , Lin. (<i>The Geometrica</i>)	.	5
Fig. 6.— <i>TESTUDO IMBRICATA</i> , Lin. (<i>The Hawk-billed Tortoise</i>)	.	8
Fig. 7.— <i>TESTUDO RADIATA</i> , Shaw. (<i>The Coui</i>)	.	6
Plate 6. Fig. 1.— <i>AMEIVA</i> (<i>The American Lizard</i>)	.	19
Fig. 2.—The Scales of the Belly of a <i>Sauvegarde</i>	.	18
Fig. 3.— <i>CENTROPYX CALCARATUS</i> (<i>The Spurred Centropyx</i>)	.	19
2nd. Plate 6. Figs. 1, 2, 3.—Head of the <i>Great Python of Java</i>	.	53
Figs. 4, 5, 6.—Head of a <i>Rattle-snake</i>	.	57
Plate 7. Fig. 1.— <i>MONITA CROCODILINUS</i> , Merr. (<i>The Great Dragon of Lacep.</i>)	.	18
Fig. 2.— <i>TUPINAMBIS ALBIGULARIS</i> , Daud. (<i>The White-throated Monitor</i>)	.	17
Fig. 3.— <i>LACERTA MAGNIFICUS</i> , Seba, (<i>The Beautiful Monitor</i>)	.	17
Plate 8. Fig. 1.— <i>LACERTA OCELLATA</i> , Daud. (<i>The Ocellated Lizard</i>)	.	20
Fig. 2.— <i>LACERTA VIRIDIS</i> , Daud. (<i>The Green Lizard</i>)	.	20
Fig. 3.— <i>LACERTA STIRPIUM</i> , Daud.	.	20

* This species figured by Daudin, closely approximates to the young of the *Testudo Imbricata*.

† See Daudin Hist. Nat. des Rep., Tom. 2, p. 277.

	REPTILIA.	Vol. II. Page
Plate 9. Fig. 1.— <i>TRAPELUS AEGYPTIUS</i> (<i>The Mutable Agama</i>) Young specimen	.	24
Fig. 2.— <i>TRAPELUS AEGYPTIUS</i> , Geoff. N. S.	.	24
Fig. 3.— <i>LACERTA CALOTES</i> , Lin.	.	25
Plate 10. Fig. 1.— <i>MONITOR GEMMATUS</i> ,* In the collection of the French Mu- seum	.	17
Fig. 2.— <i>DRACENA BICARINATA</i> ; <i>Lacerta Bicarinata</i>	.	15
Plate 11. Fig. 1.— <i>LACERTA AGILIS</i> , Lin. (<i>The Common European Lizard</i>)	.	20
Fig. 2.— <i>ALGYRA BARBARICA</i> , Cuv.; <i>Lacerta algryra</i> , Lin.	.	20
Fig. 3.— <i>TACHYDROMUS OCELLATUS</i> , Daud. (<i>The Ocellated Swift Lizard</i>)	.	20
Plate 12. Fig. 1.— <i>CORDYLUS MICROLEPIDOPUS</i>	.	21
Fig. 2.—Head of the <i>STELLIO VULGARIS</i> , (<i>The Common Stellio</i>)	.	22
Fig. 3.— <i>DORYPHORUS BREVICAUDATUS</i> , (<i>The Short-tailed Dorypho- rus of Daud</i>)	.	22
Fig. 4.— <i>UROMASTIX RETICULATUS</i> , (<i>The Reticulated Stellio of Ben- gal</i>)	.	22
Plate 13. Fig. 1.— <i>LACERTA VITTATA</i> , Gm. (<i>The White-banded Gecko</i>)	.	35
Fig. 2.— <i>STELLIO FIMBRIATUS</i> , Sch. (<i>The Fringed Gecko</i>)	.	37
Fig. 3.— <i>STELLIO GECKO</i> , Sch. (<i>The White-spotted Gecko</i>)	.	35
Fig. 4.— <i>GECKO LÆVIS</i> , D. (<i>The Smooth Gecko</i>)	.	36
Plate 14. Fig. 1.— <i>ANOLIUS VELIFER</i> , Cuv. (<i>The Great-crested Anolius</i>)	.	31
Fig. 2.— <i>ECPHIMOTUS TUBERCULATA</i> , (<i>The Tuberculated Ecpimo- tes</i>)	.	30
Fig. 3.— <i>OPHIRUS TORQUATUS</i> , Cuv. (<i>The Black-collared Grey Quetz- paleo</i>)	.	30
Plate 15. Fig. 1.— <i>ANOLIUS VELIFER</i> , (<i>The Great-crested Anolius</i>) Another view see Pl. 14.	.	31
Fig. 2.— <i>ANOLIUS EQUESTRIS</i> , (<i>The Equestrian Anolius</i>)	.	31
Fig. 3.— <i>GECKO INUNGUIS</i> , (<i>The Nailless Gecko</i>)	.	34
Fig. 4.— <i>GECKO OCELLATUS</i> , (<i>The Ocellated Gecko</i>)	.	34
Fig. 5.— <i>GECKO CEPEDIANUS</i> , (<i>The Cepedian Gecko</i>)	.	34
Plate 16. Fig. 1.— <i>PHYSIGNATHUS COCINCINUS</i> , Nob. (<i>The Physignath</i>)	.	26
Fig. 2.— <i>SITANA PONTICERIANA</i> , Cuv. (<i>The Sitan</i>)	.	28
Fig. 3.— <i>IGUANA CYCHLURA</i> , Cuv. (<i>The Carolina or Arm-tailed Igu- ana</i>)	.	29
Plate 16. Fig. 1.— <i>AGAMA JACKSONIENSIS</i> , (<i>The Bachelor</i>) In the collection of the French Museum	.	23
Fig. 2.— <i>LEIOPELIS GUTTATUS</i> , Cuv. (<i>The Striped Leiolepis</i>)	.	24
Fig. 3.— <i>CALOTES GUTTUROSA</i> (<i>The Crested Agama</i>)	.	25
Plate 18. Fig. 1.— <i>BRACHYLOPHUS FASCIATUS</i> (<i>The Banded Aguana</i>)	.	26
Fig. 2.— <i>ISTIURUS COCHINSINENSIS</i> , Vale. (<i>The Istiurus or Lophura of Cochin China</i>) In the collection of the French Museum	.	27
Plate 19. Fig. 1.— <i>OPHYRESSA SUPERCILIOSA</i>	.	29
Fig. 2.— <i>LYRIOCEPHALUS MARGARITACEUS</i> , Mer.	.	26

* This is a species which approaches in its peculiarities of conformation the *Lacerta Bivittata* of Kuhl.

REPTILIA.

	Vol. II. Page
Plate 20. Figs. 1, 2, 3.—Head of a CÆCILIA	65
Figs. 4, 5, 6.—Head of an AMPHISBÆNA (<i>The Double-walker</i>)	48
Figs. 7, 8, 9.—Head of an OPHISAURUS	46
Plate 21. Fig. 1.—DRACO FIMBRIATUS (<i>The Dragon</i>)	27
Fig. 2.—SITANA PONTICERIANA (<i>The Sitan</i>)	27
Plate 22. Fig. 1.—IGUANA NUDICOLLIS, Cuv. (<i>The Naked Aguana</i>)	29
Fig. 2.—BASILISCUS; <i>Lacerta basiliscus</i> , Lin. (<i>The Basilisk</i>)	30
Fig. 3.—POLYCHRUS MARMORATUR, Lin. (<i>The Marbled Lizard</i>)	30
Plate 23. Fig. 1.—HEMIDACTYLUS TUBERCULOSUS, Daud. (<i>The Spotted Hemidactylus</i>)	36
Plate 24. Fig. 1.—PHYLLURUS PLATURUS, White	38
Fig. 2.—Foot of the PLATYDACTYLUS INUNGUIS	34
Fig. 3.—Foot of the HEMIDACTYLUS TRIHEDRUS	36
Fig. 4.—Foot of the THECADACTYLUS	36
Fig. 5.—Foot of the PTYODACTYLUS LOBATUS	37
Fig. 6.—Foot of the SPHEROIDACTYLUS	38
Fig. 7.—Foot of the STENODACTYLUS GUTTATUS	38
Plate 25. Fig. 1.—TORTRIX MACULATA, Lin. (<i>The Variegated Tortrix of America</i>)	50
Fig. 2.—BOA CANINA, Lin. (<i>The Canine Boa</i>)	52
Fig. 3.—Tail of the UROPELTIS CEYLANICUS, Cuv. (<i>The Uropeltis of Ceylon</i>)	50
Plate 25. bis. Fig. 1.—CHAMÆLEO AFRICANUS (<i>The Common Chameleon</i>)	39
Fig. 2.—SCINCUS CÆLESTINUS, Valenc. (<i>The Blue Scinc</i>) In the collection of the French Museum	41
Fig. 3.—SEPS STRIATA (<i>The Striped Seps</i>)	42
Plate 26. Fig. 1.—ERIX BENGALENSIS, Merr. (<i>The Hairy Serpent of Bengal</i>)	52
Fig. 2.—HEAD of the SCYTALE CORONATA, Merr. (<i>The Crowned Syc-tale or False Boa</i>)	52
Fig. 3.—ERPETON TENTACULATUS, Lacep. (<i>The Tentaculated Serpent</i>)	53
Plate 26. bis. Fig. 1.—BIPES LEPIDOPODUS, Lacep. (<i>The Two-footed Lepidopodus</i>)	43
Fig. 2.—CHALCIS TETRADACTyla, Lacep.	44
Fig. 3.—CHIROTES CANALICULATUS, Lacep.	44
Plate 27. Fig. 1.—PSEUDOPUS DURVILLII (<i>The False-foot of Durville</i>)	46
Fig. 2.—HEAD of the ANGUIS FRAGILIS, Lin. (<i>The Common Blind worm of Europe</i>)	47
Fig. 3.—ACONTIAS MELEAGRIS, Lin. (<i>The Darting Serpent of the Cape of Good Hope</i>)	47
Plate 28. Fig. 1.—AMPHISBÆNA FULIGINOSA, Lin. (<i>The Double-walker</i>)	48
Fig. 2.—TYPHLOPS CINEREUS, Schn. (<i>The Blind Serpent of Schneider</i>)	49
Plate 29. Fig. 1.—VIPERA BERUS, Lin. (<i>The common Viper</i>)	59
Fig. 2.—CROTALUS HORRIDUS, Lin. (<i>The Diamond Rattle-snake</i>)	59
Plate 30. Fig. 1.—TRIMERESURUS TORTOR, Less. (<i>The Winding Trimeresurus</i>)	62
Fig. 2.—ACANTHOPHIS CERASTINUS, Daud. (<i>The Venomous Acanthophis</i>)	62
Plate 31. Fig. 1.—PYTHON SCHNEIDERII, Merr. (<i>The Great Python of Schneider</i>)	53
Fig. 2.—DIPSAS CYANODON, Cuv. (<i>The Fatal Thirst-causing Serpent of India</i>)	54

TABLE OF THE PLATES.

REPTILIA.

	Vol. II. Page
Plate 31. Fig. 3.—Head of the <i>XENOPELTIS CONCOLOR</i> , Reinwardt. (<i>The Many-coloured Xenopeltis</i>)	53
Plate 32. Fig. 1.— <i>DENDROPHIS AHÆTIILLA</i> (<i>The Serpent of the Woods</i>)	54
Fig. 2.— <i>DRYINUS NASUTUS</i> , Russel (<i>The Hog-nosed Dryinus or Coluber</i>)	54
Plate 33. Fig. 1.— <i>BUNGARUS ANNULARIS</i> , Daud.	63
Fig. 2.— <i>ANGUIS PLATURUS</i> , Lin.	63
Plate 34. Fig. 1.— <i>HYDROPHIS CLORIS</i> , Russel	63
Fig. 2.— <i>CÆCILIA LOMBRICOIDÆA</i> , Daud.	65
Fig. 3.— <i>CÆCILIA ALBIVENTRIS</i> , Daud.	65
Plate 35. Fig. 1.— <i>HYDROPHIS FASCIATUS</i> , Russel (<i>The Bengal Water Serpent</i>)	63
Fig. 2.— <i>CÆCILIA BIVITTATA</i> , Cuv. (<i>The Blind Serpent or The Cæcilia of America</i>)	68
Fig. 3.—.	65
Plate 36. Fig. 1.— <i>RANA PALUSTRIS</i> , Le Conte.	68
Fig. 2.— <i>CERATOPHRIS GRANOSA</i> (<i>The American Horned Frog</i>)	69
Fig. 3.— <i>HYLA BICOLOR</i> , Daud. (<i>The South American Two-coloured Tree Frog</i>)	69
Plate 37. Fig. 1.— <i>BUFO VULGARIS</i> (<i>The Common Toad</i>)	70
Fig. 2.— <i>OXYRHYNCHUS BICOLOR</i> , Val. (<i>The Angular-muzzle Toad</i>) In the collection of the French Museum	72
Fig. 3.— <i>ENGYSTOMA MARMORATUM</i> (<i>The Marbled Toad</i>)	72
Plate 38. Fig. 1.— <i>SALAMANDRA ATRA</i> , Laurenti (<i>The Black Salamander</i>).	74
Fig. 2.— <i>TRITON VITTATUS</i> , Valenc. (<i>The Triton</i>) In the collection of French Museum.	75
Fig. 3.— <i>MONOPOMA GIGANTEA</i> , Barton (<i>The Great Salamander of North America</i>)	76
Plate 39. Fig. 1.— <i>AMPHIUMA MEANS</i> , Gard. and Harl. of the French Museum (<i>The Two-toed Amphiuma</i>)	76
Fig. 2.— <i>AMPHIUMA TRYDACTYLUM</i> (<i>The Three-toed Amphiuma</i>)	76
Fig. 3.— <i>AXOLOTL PISCIFORMIS</i> , Shaw (<i>The Fish Serpent of Mexico</i>)	76
Fig. 4.— <i>MENOBRANCHUS LATERALIS</i> , Harl.	77
Plate 40. Fig. 1.— <i>PROTEUS ANGUINUS</i> , Laur. (<i>The Blind Proteus</i>)	77
Fig. 2.— <i>SIREN STRIATA</i> , Le Conte (<i>The Streaked Siren</i>)	78
Fig. 3.— <i>SIREN INTERMEDIA</i> , Le Conte (<i>The Intermediate Siren of Le Conte</i>)	77



P I S C E S.

Plate 1. Fig. 1.— <i>ETELIS CARBUNCULUS</i> (<i>The Etelis</i>)	87
Fig. 2.— <i>DIPLOPRION BIFASCIATUM</i> (<i>The Bifasciate Diploprion</i>)	87
Fig. 3.—Head of the <i>HURO NIGRICANS</i> (<i>The Black Basse</i>)	87
Fig. 4.—Head of the <i>NIPHON SPINOSUS</i> (<i>The Spiny Niphon</i>)	87
Fig. 5.—Head of the <i>ENOPLOSSUS ARMATUS</i> (<i>The Enoplosus or Armed Perch of Japan</i>)	87
Plate 2. Fig. 1.— <i>PERCA ACUTA</i> , Cuv. and Val.	85
Fig. 2.— <i>LATES NOBILIS</i> , Cuv. and Val.	86
Fig. 3.— <i>CENTROPOMUS UNDECIMALIS</i> , Cuv. (<i>The Eleven-spined Centropomus</i>)	86

TABLE OF THE PLATES.

v

PISCES.

	Vol. II. Page
Plate 3. Fig. 1.— <i>APOGON TRIMACULATUS</i> (<i>The Mediterranean Apogon</i>) .	87
Fig. 2.— <i>CHEILODIPTERUS ARABICUS</i> (<i>The Arabian Cheilodipterus</i>) .	88
Fig. 3.— <i>AMBASSIS COMMERSONII</i> (<i>Commerson's Ambassus</i> ; or, <i>The Dugol</i>)	88
Plate 4. Fig. 1.— <i>LUCIO-PERCA SANDRA</i> , Cuv. (<i>The Common Pike Perch</i>) .	89
Fig. 2.— <i>LUCIO-PERCA AMERICANA</i> (<i>The American Perch</i>) .	89
Fig. 3.— <i>HURO NIGRICANS</i> , Cuv. (<i>The Black Basse of the Huron</i>) .	87
Plate 5. Fig. 1.— <i>LUCIO-PERCA VOLGENSIS</i> , Pallas (<i>The Russian Perch</i>) .	89
Fig. 2.— <i>NIPHON SPINOSUS</i> , Cuv. (<i>The Spined Niphon</i>) .	87
Fig. 3.— <i>ENOPLOSSUS ARMATUS</i> , Lacep. (<i>The Armed Enoplossus</i>) .	87
Plate 6. Fig. 1.— <i>SERRANUS ANTHIAS</i> (<i>The Spined Serranus</i>) .	89
Fig. 2.— <i>SERRANUS CABRILLA</i> , Nob. (<i>The Common Serranus</i>) .	89
Fig. 3.— <i>SERRANUS GIGAS</i> , Nob. (<i>The Merra of the Mediterranean</i>) .	90
Plate 7. Fig. 1.— <i>SERRANUS PHAETON</i> , Nob. (<i>The Long-tailed Serranus</i>) .	91
Fig. 2.— <i>SERRANUS FASCICULARIS</i> , Nob. (<i>The Banded Serranus</i>) .	89
Fig. 3.— <i>SERRANUS OCULATUS</i> , Nob. (<i>The Large-eyed Serranus</i>) .	90
Plate 8. Fig. 1.— <i>MESOPRION UNINOTATUS</i> , Cuv. (<i>The One-spotted Mesoprion</i>) .	91
Fig. 2.— <i>MESOPRION CHRYSURUS</i> , Nob. (<i>The Golden-tailed Mesoprion</i>) .	91
Fig. 3.— <i>DIACOPE RIVULATA</i> , Nob. (<i>The Banded Diacope</i>) .	91
Plate 8. bis. Fig. 1.— <i>LABRAX MUGRONATUS</i> , Cuv. and Val. .	86
Fig. 2.— <i>LABRAX LUPUS</i> , Cuv. (<i>The Basse Perch</i>) .	85
Fig. 3.— <i>SERRANUS SCRIBA</i> , Cuv. (<i>The Mediterranean Sea Perch</i>)	89
Plate 9. Fig. 1.— <i>STERNOPTYX DIAPHANA</i> , Herm. . . .	199
Fig. 2.— <i>STERNOPTYX OLfersii</i> , Cuv. . . .	199
Fig. 3.— <i>GYNMARCHUS SENEGALENSIS</i> ; or <i>Niloticus</i> , Cuv. . . .	226
Plate 10. bis. Fig. 1.— <i>PLECTROPOMA LEOPARDINUS</i> , Cuv. (<i>The Leopard-spotted Plectropoma</i>)	91
Fig. 2.— <i>PLECTROPOMA PUELLA</i> , Cuv. (<i>The Beautiful Plectropoma</i>)	91
Fig. 3.— <i>SERRANUS ALTIVELIS</i> , Cuv. (<i>The Large-finned Serranus</i>)	91
Plate 11. Fig. 1.— <i>PENTACEROS CAPENSIS</i> , Nob. . . .	93
Fig. 2.— <i>GRYSTES SALMOIDES</i> , Nob. (<i>The Salmon-formed Growler</i>) .	93
Fig. 3.— <i>CENTROPISTES NIGRICANS</i> , Nob. (<i>The American Black Bate, or Perch</i>)	93
Plate 12. Fig. 1.— <i>PERCA SERRATO-GRANULATA</i> , Cuv. (<i>The Granulated Perch</i>)	85
Fig. 2.— <i>GRAMMICHTHYS ORIENTALIS</i> , Cuv. (<i>The Oriental Grammist</i>) .	86
Fig. 3.— <i>ASPRO VULGARIS</i> , Cuv. (<i>The Common Aspro</i>) .	87
Fig. 4.—Head of the <i>LABRAX VULGARIS</i> ; <i>Labrax lupus</i> , Cuv. (<i>The Basse Perch</i>)	85
Fig. 5.—Head of the <i>LATES NOBILIS</i> , Cuv. (<i>The American Pike</i>) .	86
Plate 12. bis. Fig. 1.— <i>HERLIASES INSOLATUS</i> , Cuv. . . .	114
Fig. 2.— <i>POMACENTRUS FASCIATUS</i>	113
Fig. 3.— <i>GYPHISODON CÆLESTINUS</i>	114
Plate 13. Fig. 1.— <i>CENTROPRISTES AURORUBENS</i> , Nob. (<i>The Golden Red Perch</i>)	93
Fig. 2.— <i>GRYSTES MACQUARIENSIS</i> (<i>The Growler of New South Wales</i>)	93
Fig. 3.—Head of the <i>ACERINA VULGARIS</i> (<i>The Common Gremille</i>) .	92
Fig. 4.—Head of the <i>POLYPRION CERNUM</i> (<i>The Mediterranean Cer-nier</i>)	92

PISCES.	Vol. II. Page
Plate 13. Fig. 5.—Head of the <i>RYPTICUS ARENATUS</i> , Cuv. (<i>The Soap-wort</i>)	92
Fig. 6.—Fin of the <i>CIRRHITES FASCIATUS</i>	93
Plate 13. bis. Fig. 1.— <i>DULES MARGINATUS</i> , Nob. (<i>The Bordered Doule</i>)	94
Fig. 2.— <i>DATNIA ARGENTEA</i>	94
Fig. 3.— <i>THERAPON THERAPS</i> , Cuv.	94
Plate 14. Fig. 1.— <i>APOGON REX MULLORUM</i> , Cuv. (<i>The Red Surmullet</i>)	88
Fig. 2.— <i>POMATOMUS TELESCOPIUM</i> , Risso (<i>The Telescope Pomatomus</i>)	88
Fig. 3.— <i>LUCIO-PERCA SANDRA</i> , Cuv. (<i>The Common Sandre</i>). For another view, see plate 4. fig. 1.	89
Fig. 4.—Head of the <i>CHEILODIPTERUS ARABICUS</i> , Cuv. and Val. (<i>The Tabah of the Arabs</i>)	88
Fig. 5.—Head of the <i>AMBASSIS COMMERMORONII</i> , (<i>The Dugol</i>)	88
Plate 14. bis. Fig. 1.— <i>DULES AURIGA</i> , Nob.	94
Fig. 2.— <i>POMOTIS VULGARIS</i> , Nob. (<i>The Common Pomatis or American Pond Perch</i>)	93
Fig. 3.— <i>PRIACANTHUS JAPONICUS</i> , Nob. (<i>The Priacanth of Japan</i>)	94
Plate 15. Fig. 1.— <i>SERRANUS HEXAGONATUS</i> , Cuv. (<i>The Hexagon Sea Perch</i>)	90
Fig. 2.— <i>MESOPRION AYA</i> , Cuv. (<i>The Garanha or Red Mesopriion</i>)	91
Fig. 3.—Head of the <i>PLECTROPOMUS PUELLA</i> , Cuv.	91
Fig. 4.—Head of the <i>SERRANUS ANTHIUS</i> , Cuv. (<i>The Mediterranean Barber</i>)	90
Fig. 5.—Head of the <i>DIACOPE RIVULATA</i> , Cuv. (<i>The Coromandal Orati</i>)	91
Plate 16. Fig. 1.— <i>CENTRARCHUS SPAROIDES</i> , Lacep.	94
Fig. 2.— <i>DULES RUPESTRIS</i> , Cuv. (<i>The Doule</i>)	94
Fig. 3.—Head of the <i>POMOTIS VULGARIS</i> (<i>The American Pond Perch</i>)	93
Fig. 4.—Head of the <i>PRIACANTHUS JAPONICUS</i> , Nob. (<i>The Priacanthus of Japan</i>)	94
Plate 16. bis. Fig. 1.— <i>PELOR FILAMENTOSUM</i> , Cuv.	107
Fig. 2.— <i>PLATYCEPHALUS ASPER</i> , Cuv. and Val.	105
Fig. 3.— <i>DACTYLOPTERUS ORIENTALIS</i> , Cuv. (<i>The Flying Gurnard</i>)	103
Plate 17. Fig. 1.— <i>THERAPON PUTA</i> , Cuv. (<i>The Therapon</i>)	94
Fig. 2.— <i>PELATES QUADRI-LINEATUS</i> (<i>The Four-lined Pelates</i>)	94
Fig. 3.— <i>TRICHODON STELLERI</i> , Cuv. (<i>The Trichodon</i>)	95
Fig. 4.—Head of the <i>HELOTES SEX-LINEATUS</i> , Cuv. (<i>The Six-lined Helotes</i>)	95
Plate 17. ter. Fig. 1.— <i>HELOTES SEX-LINEATUS</i> , Cuv.	95
Fig. 2.— <i>MYRIPRISTIS JAPONICUS</i> , Cuv. and Val.	96
Fig. 3.— <i>HOLOCENTRUM HASTATUM</i>	96
Plate 18. Fig. 1.— <i>TRACHINUS ARANEUS</i> Riss. (<i>The Weever</i>)	97
Fig. 2.— <i>PERCIS CANCELLOTA</i> , Cuv.	98
Fig. 3.— <i>URANOSCOPIUS GUTTATUS</i> , Cuv. (<i>The Star-gazer</i>)	97
Fig. 4.—Head of the <i>PINGUIPES BRASILIANS</i> , Cuv.	97
Fig. 5.—Head of the <i>PERCOPHIS BRASILIANS</i> , Cuv.	97
Plate 18. bis. Fig. 1.— <i>SILLAGO ERYTHREA</i> ,	95
Fig. 2.— <i>MYRIPRISTES SEYCHELLENSIS</i> ,	96
Fig. 3.— <i>BERYX DECADACTYLUS</i> ,	96

PISCES.

	Vol. II. Page
2nd. Plate 18. bis. Fig. 1.— <i>PERCIS SEMI-FASCIATA</i> , Cuv. and Val. (<i>The Half-banded Percis</i>).	97
Fig. 2.— <i>PERCOPHIS BRASILIANS</i> , (<i>The Brazilian Percophis</i>)	97
Fig. 3.— <i>URANOSCOPUS INERMIS</i> , Cuv. and Val. (<i>The Un-armed Star-gazer</i>)	98
 Plate 18. ter. Fig. 1.— <i>BERYX LINEATUS</i> , Cuv. and Val.	96
Fig. 2.— <i>TRACHINUS RADIATUS</i> , Cuv	97
Fig. 3.— <i>PINGUIPES BRASILIANS</i> , Cuv.	97
 Plate 19. Fig. 1.— <i>POLYNEMUS ENNEADACTYLUS</i> , Vahl.	99
Fig. 2.— <i>SPYHRÆNA VULGARIS</i> , Lin.	99
Fig. 3.— <i>PARALEPIS COREGONOIDES</i> , Risso.	99
Fig. 4.— <i>UPENEUS LATEUS</i> Cuv. and Val.	100
Fig. 5.—Head of the <i>MULLUS BARBATUS</i> , Lin. (<i>The Red Surmullet</i>)	100
 Plate 19. bis. Fig. 1.— <i>POLYNEMUS QUADRI-FILIS</i> (<i>The Four-filimented Paradise Fish</i>)	99
Fig. 2.— <i>PARALEPIS SPHYRÆNOIDES</i> , Risso	99
Fig. 3.— <i>SPHYRÆNA BARRACUDA</i> , Cuv. and Val.	99
 Plate 20. Fig. 1.—Head of <i>GADUS MORRHUA</i> , Lin. (<i>The Cod</i>)	209
 Plate 20. bis. Fig. 1.— <i>SILLAGO DOMINA</i> , Cuv.	95
Fig. 2.— <i>MULLUS BARBATUS</i> , Lin. (<i>The Red Surmullet</i>)	100
Fig. 3.— <i>UPENEUS VLAMINGII</i> , Cuv. and Val.	100
 Plate 21. bis. Fig. 1.— <i>AGRIOPUS VERRUCOSUS</i>	107
Fig. 2.— <i>PELOR JAPONICUM</i>	107
Fig. 3.— <i>SYNANCHIA EROSA</i> , Cuv.	107
 Plate 22. bis. Fig. 1.— <i>SCORPÆNA GRANDICORNIS</i>	105
Fig. 2.— <i>BEMBRAS JAPONICUS</i>	105
Fig. 3.— <i>HEMILEPIDOTUS TILESII</i>	104
 Plate 23. Fig. 1.— <i>TRIGLA PÆCILOPTERA</i> , Cuv. and Val.	101
Fig. 2.— <i>CEPHALACANTHUS SPINARELLA</i> , Cuv.	103
Fig. 3.— <i>COTTUS POROSUS</i> , Cuv.	103
Fig. 4.— <i>COTTUS OCTODECIM SPINOSUS</i> , Cuv.	103
 Plate 23. bis. Fig. 1.— <i>PRIONOTUS TRIBULUS</i> , Cuv.	102
Fig. 2.— <i>TRIGLA LUCERNA</i> , Brünn.	102
Fig. 3.— <i>PERISTEDION CATAPHRACTA</i> , Lacep.	102
 Plate 24. Fig. 1.— <i>ASPIDIPHORUS SEGALIENSIS</i> , Cuv.; <i>Agonus lavigatus</i> , Tiles	104
Fig. 2.—Head of the <i>ASPIDIPHORUS QUADRICORNIS</i> , Cuv.	104
Fig. 3.— <i>HEMITRIPTERUS AMERICANUS</i> , Cuv.	104
Fig. 4.— <i>HEMILEPIDOTUS TILESII</i> , Cuv.	104
Fig. 5.—Head of the <i>PLATYCEPHALUS ASPER</i> , Cuv.	105
 Plate 24. bis. Fig. 1.— <i>TRIGLA PAPILIO</i> , Cuv. (<i>The Butterfly Gurnard</i>)	102
Fig. 2.— <i>COTTUS BUBALIS</i> , Euphr. (<i>The Bull Head</i>)	103
Fig. 3.— <i>OPLICHTHYS LANGSDORFII</i> , Cuv. and Val., Hist. Nat. des Poissons, Vol. IV.	102
 Plate 24. ter. Fig. 1.— <i>SCIÆNA PAMA</i>	109
Fig. 2.— <i>OTOLITHUS RUBER</i> , Cuv.	109
Fig. 3.— <i>OTOLITHUS TEROE</i> , Cuv. and Vall.; or <i>Lutjan de Cayenne</i> , Lacep.	109

PISCES.	Vol. II.	Page
Plate 25. Fig. 1.— <i>PLATYCEPHALUS GRANDISPINIS</i>	.	105
Fig. 2.— <i>SCORPÆNA NESOGALLICA</i> , Cuv.	.	105
Fig. 3.— <i>TÆNIANOTUS TRIACANTHUS</i> , Cuv.	.	105
2nd. Plate 25. Fig. 1.— <i>CHEILODACTYLUS ANTONII</i>	.	112
Fig. 2.— <i>AMPHIPRION CHRYSOGASTER</i>	.	113
Fig. 3.— <i>HELIASES FRENATUS</i>	.	114
Fig. 4.—Head of the <i>SCOLOPSIDES LYCOGENIS</i>	.	113
Fig. 5.—Head of the <i>PREMNAS SEMICINCTUS</i>	.	113
Fig. 6.—Head of the <i>DASCYLLUS MARGINATUS</i>	.	114
Fig. 7.—Head of the <i>POMACENTRUS FASCIATUS</i>	.	113
Fig. 8.—Head of the <i>GLYPHISODON CÆLESTINUS</i>	.	114
Plate 25. bis. Fig. 1.— <i>SEBASTES NORVEGICUS</i> , Cuv.	.	106
Fig. 2.— <i>BLEPSIAS VILLOSUS</i> , Cuv.	.	106
Fig. 3.— <i>PTEROIS VOLITANS</i> , Cuv.	.	106
Plate 25. ter. Fig. 1.— <i>OTOLITHUS GUATUCUPA</i> , Marcg.	.	109
Fig. 2.— <i>ANCYLODON PARVIPINNIS</i> , Cuv. and Val.	.	110
Fig. 3.— <i>CORVINA SEMILUCTUOSA</i> , Cuv.	.	110
Plate 26. Fig. 1.— <i>SEBASTES VARIABILIS</i> , Cuv.	.	106
Fig. 2.— <i>BLEPSIAS TRILOBUS</i> , Cuv.	.	105
Fig. 3.— <i>APISTES MARMORATUS</i> , Cuv.	.	107
2nd. Plate 26. Fig. 1.— <i>PAGELLUS CENTRODONTUS</i>	.	116
Fig. 2.— <i>BOOPS SALPA</i>	.	117
Fig. 3.—Head and teeth of the <i>SARGUS RONDELETHII</i> (<i>The Sargus of Rondelet</i>)	.	115
Fig. 4.—Head and Dentition of the <i>CHRYSOPHYS AURATA</i> , Lin. (<i>The Golden Eye-brow</i>)	.	115
Fig. 5.—Head and Dentition of the <i>PAGELLUS ERYTHRINUS</i> , Lin. (<i>The Pagel</i>)	.	116
Plate 26. bis. Fig. 1.— <i>EQUES PUNCTATUS</i>	.	111
Fig. 2.— <i>BORIDIA GROSSIDENS</i> , Cuv. and Val.*	.	111
Fig. 3.— <i>ELEGINUS MACLOVINUS</i> , Cuv. and Val. †	.	111
Plate 26. ter. Fig. 1.— <i>HÆMULON QUADRILINEATUM</i> , Cuv. and Val.	.	111
Fig. 2.— <i>HÆMULON HETERODON</i> , Desm.	.	111
Fig. 3.— <i>PRISTIPOMA BILINEATUM</i> , Cuv.	.	112
Plate 26. quar. Fig. 1.— <i>PREMNAS SEMICINCTUS</i>	.	113
Figs. 2, 3.—The Brain of the <i>SCIÆNA PAMA</i> ; <i>Bola Pama</i> , Buchanan	.	109
Fig. 4.— <i>DASCYLLUS MARGINATUS</i> , Ehr.	.	114
Plate 27. Fig. 1.— <i>AGRIOPUS PERUVIANUS</i> , Cuv.	.	107
Fig. 2.— <i>PELOR MACULATUM</i> , Cuv.	.	107
Fig. 3.— <i>SYNANCEYA ELONGATA</i> , Cuv. ‡	.	107

* A new species of *Sciænoides* with two dorsals, which possesses some of the characters of the *Corvina*, with the addition of numerous large teeth in both jaws.

† Another new species of *Sciænoides* with two dorsals, slightly different from the preceding, and not possessing those large teeth, which is their chief characteristic.

‡ A new species described by Cuvier and Vall. in the fourth volume of their *Ichthyology*.

PISCES.

		Vol. II.	Page
Plate 27. ter. Fig. 1.— <i>LOBOTES SOMNOLENTUS</i> , Cuv.	.	.	112
Fig. 2.— <i>DIAGRAMMA ORIENTALE</i> , Cuv.	.	.	112
Fig. 3.— <i>DIAGRAMMA CARVIFRONS</i> , Cuv.	.	.	112
2nd. Plate 27. ter. Fig. 1.— <i>GLYPHISODON KORUSCHI</i> , Cuv.	.	.	114
Fig. 2.— <i>AMPHIPRION LATICLAVIUS</i> , Cuv.	.	.	113
Fig. 3.— <i>AMPHIPRION TUNICATUS</i> , Cuv.	.	.	113
Plate 27. quar. Fig. 1.— <i>GASTEROSTEUS TRACHURUS</i> , Cuv. and Val.	.	.	108
Fig. 2.— <i>GASTEROSTEUS GYMNRURUS</i> , Cuv. (<i>The Stickle-back</i>).	.	.	108
Fig. 3.— <i>GASTEROSTEUS NOVEBORACENSIS</i> , Cuv. and Val.	.	.	108
Fig. 4.— <i>GASTEROSTEUS LEIURUS</i> , Cuv. and Val.	.	.	108
Plate 28. Fig. 1.— <i>MONOCENTRIS JAPONICA</i> , Cuv.	.	.	107
Fig. 2.— <i>GASTEROSTEUS BRACHYCENTRUS</i> , Cuv. (<i>The Three-spined Stickle-back</i>).	.	.	108
Fig. 3.— <i>OREOSOMA CONIFERUM</i> , Cuv.	.	.	108
Plate 28. bis. Fig. 1.— <i>LARIMUS BREVICEPS</i> , Cuv. (<i>The Larime</i>)*			

* After all the groups that have been described, and which compose the regular series of *Sciænoides* with two dorsals, there still remains some fish of this family which escape all the characters assigned to any of them, and which, should each become the type of a peculiar group to which we ought to give a generic name, we will call them if you please anomalous *Sciænas*; but remembering that nature acknowledges nothing anomalous in what exists, and that there is no anomaly, but relatively to the incomplete abstractions of our mind, Cuvier has given the name of *Larimus* to some of these fish. They are *Sciænoides* with two dorsals, teeth as the pile on velvet; the front of the cranium does not project forward, and the snout is very short; their preoperculum is slightly indented.

The *L. breviceps*, Cuv., resembles the *Corvina* in almost everything; but far from having the snout swelled and prominent, it has it flat and extremely short as in the *Barber*, (*Serranus anthias*); so that its eye, which is large, is in the anterior half of the length of the head; nevertheless, the jaws are cavernous, and there are no teeth in the palate. The length of the head measures nearly the fourth of the whole length, and the height of the body slightly exceeds the length of the head. The opening of the mouth descends backward; the openings of the nostril are all near the anterior edge of the eye. There is no pore to the lower jaw. The teeth are soft, on a very narrow band. The under orbital is not more distinguished than in the other *Sciænas*, and gives the same shelter to a part of the maxillary; all the head, except the lips and the maxillary, is scaly. There are scales even on the lower jaw, but not on the membrane of the gills, which are well cut, and have seven rays; of which the upper are flat, and the under very small. The preoperculum has the border wide and almost without any apparent indentations. The bone of the operculum terminates in two points, but concealed, obtuse, and so frail as scarcely to be felt. The first dorsal is triangular and is not half the height of the body; it has ten spiny rays disposed as usual. The second has one spiny and twenty-eight soft rays. The scales ascend between the rays, even to the half of its height. The pectorals are long and pointed; the ventrals wide, and almost equal to the pectorals in length. The anal less in length than in height, has only seven soft rays, but its second spine is strong as in the corvines. This fin is placed under the middle of the second dorsal, and much behind the anus, which is exactly between the anal and the ventrals, under the commencement of the second dorsal. The caudal is rhomboidal; it has a line of scales on its middle. The scales on the body are very large: there are about forty-five on a longitudinal line, and fifteen on a vertical. There are some between the bases of the dorsal and the anal. The lateral line is parallel to the back, and one third above, as in the greater part of the *Sciænas*. The

PISCES.	Vol. II. Page
Plate 28. bis. Fig. 2.— <i>LEPIPTERUS FRANCISCI</i> , Cuv. (<i>The Lepipterus of the Island of St. Francis</i>)*	111
Fig. 3.— <i>CORVINA OCELLATA</i> , Cuv. (<i>The Spotted Corvina</i>)	110
Fig. 4.—Anatomical details of the Brain of the <i>ANABAS SENNAL</i> .	144
Fig. 5.—Anatomical details of the Brain of the <i>OPHICEPHALE</i> .	146
2nd. Plate 28. bis. Fig. 1.— <i>POGONIAS FASCIATUS</i> , Cuv. (<i>The Banded Pogonias</i>)	110
Fig. 2.— <i>UMBIRINA COROIDES</i> , Cuv.	110
Fig. 3.— <i>MICROPOGON LINEATUS</i> , Cuv.; <i>Umbrina fourneiri</i> , Desmar.	111

whole of this fish is silvered; its back, tinted with grey brown, has brownish oblique lines, which descend in advancing towards the tail; on the flank these lines become longitudinal; a little lower down they fade away; and there are none on the lower part. The membrane of the first dorsal is blackish, with a white triangular spot at its base in each interval of the rays. The individual from which our drawing was taken, is seven to eight inches in length. It was M. Delalande who brought the first Larimes to Europe from Brazils, but there have been recently several received from St. Domingo through M. Ricord. It is called in that island *silver-fish*, on account of its colour, which in its fresh state is as *brilliant* (according to our traveller) as that of the purest mercury. M. Ricord adds that this fish is a foot long, that its flesh is bad, and only eaten by the poor.

* The name of *Lepipterus* designates *Sciænoides* with two dorsals, teeth as the pile on velvet, prolonged snout, the front of the cranium rather concave, and the vertical fins of which, are very scaly. We know only one—the *Lepipterus Francisci*. This fish seems also to resemble the *Corvina*, by its soft teeth and its large anal spine: but it is a *Corvina* with the front of the cranium lengthened, and even a little concave; besides, the scales which cover its second dorsal, and its caudal, approximates it to the *Eques* and to the *Polynemus*. Its general appearance is similar to the *Otolithus tørœ*, Cuv.; but, with a little attention, it is easily distinguished from it. It is a fish altogether of an elongated form, head long, front of the cranium rather concave. Its height is only the sixth of its length, and its head is almost the fourth; but its head is a third less in height than in length. The under orbital, covered by scales, which do not permit us to distinguish it from the cheek, forms a border, under which the upper jaw retires, as in the *Ombrinæ*; the mouth is very small, and it has very fine soft teeth, like the pile on velvet, in the two jaws only. All the opercular pieces are scaly, like the cranium and the snout. The preoperculum has some indentations slightly marked towards its angle, which is rounded. The bone of the operculum ends by a single flat point. The pectoral is moderate and pointed; the ventral proceeds a little further back than it, and passes it. The first dorsal rises in the middle of the pectoral; it has ten spiny rays, very weak and little elevated, the first of which is very short; the second dorsal has a spiny ray and thirty-three soft ones; it is nearly of equal height everywhere. The caudal is rounded. They are both completely scaly, so that even in the caudal it is difficult to count the rays. The anal only occupies a small space in length, under the middle of the second dorsal. It is twice as high as it is long. Its first spine is very short; the second, compressed and arched like a sabre, is wider and stronger than in any other *Sciæna*; there are afterwards seven soft rays and some scales between them. The scales of the body are smooth, fine, and slightly striated on the edges. There is one stout spine, rather free, but not scaly, formed by the coracoidien, in the middle of the pectoral. The lateral line is nearly parallel to the back, and marked by simple but continuous tubercles.

This fish is nineteen or twenty inches long, is entirely of silver colour, with very numerous oblique, grey, or bluish lines throughout its entire length. There is a continuation of brown lines in the intervals of the rays of the first dorsal, and brown dots on those of the second. The other fins have only spots. This is one of the numerous acquisitions to natural history, procured by M. Augustine St. Hilaire. He took it in the river of St. Francis, in Brazils.

PISCES.

Vol. II. Page

3rd. Plate 28. bis. Fig. 1.— <i>MÆNA VOMERINA</i> , Cuv. ; A new species from the Mediterranean.	.	.	.	118
Fig. 2.— <i>BOOPS VULGARIS</i> , Cuv.	117
Fig. 3.— <i>APHAREUS CÆRULESCENS</i> , Cuv.*	.	.	.	119
Fig. 4.—A fragment of the intermaxillary jaw of a <i>Daurades</i> , with the Molars detached; <i>Chrysophys auratus</i> , Cuv.	.	.	.	115
Fig. 5.—The superior intermaxillary jaw of a <i>Sargus. a.</i> The incisors; b. The germ of an incisor tooth; c. The molars.	.	.	.	115
Plate 28. ter. Fig. 1.— <i>LATILUS DOLIATUS</i> †	112
Fig. 2.— <i>CHEILODACTYLUS ZONATUS</i> , Cuv.	112
Fig. 3.— <i>MACQUARIA AUSTRALASIÆ</i> ,‡ Cuv.	112
2nd. Plate 28. ter. Fig. 1.— <i>APSILUS FUSCUS</i> , Cuv. (<i>The Brown Apsilus</i>)§	93
Fig. 2.— <i>CHÆTODON STRIGATUS</i> , Langsd.	120
Fig. 3.— <i>CHÆTODON RETICULATUS</i>	120
Plate 28. quar. Fig. 1.— <i>CENTRARCHUS SPAROIDES</i> ; <i>Labrus sparoides</i> , Lacep.	.	.	.	94
Fig. 2.— <i>RYPTICUS ARENATUS</i> , Cuv. and Val.	92
Fig. 3.— <i>CIRRHITES FASCIATUS</i> (<i>The Banded Cirrhite</i>). A new species very nearly allied to the <i>Cirrh. aprinus</i>	93
2nd. Plate 28. quar. Fig. 1.— <i>TÆNIANOTUS TRIACANTHUS</i> , Lacep. For another view, see pl. 25, fig. 3	105
Fig. 2.— <i>APISTUS TRACHINOIDES</i> 	136

* Labre fourche et Caranxomore Sacrestin of Lacep. A fish of the family Menides but not classed by Cuvier; its place should be after the Gerries.

† A new species brought by MM. Quoy and Gaimard from the Isle of France. It is a neighbouring species to the *Cheilodactylus zonatus*.

‡ A new species discovered in Macquarie River, New Holland. It approaches closely to the *Latilus doliatus* and the *Coryphéne chinoise*, Lacep.

§ It is a new species found by MM. Quoy and Gaimard during their second voyage, and closely approximates to the *Grystes Macquariensis*.

|| This is one of the most remarkable among all the little Apistes. It was sent from Java by Messers Kuhl and Van Hasselt. The three first thorny rays of its dorsal, placed on the neck, are so distant from the others, and the membrane that unites them to the rest of the fin is so low, that they form, it may be said, a distinct fin, so that at the first glance we are inclined to take the fish for a *Trachinus*; but we are quickly undeceived when we remark that the rays following are also thorny, on almost the whole length of the dorsal; in short its figure is nearly that of the *Scorpæna scrofa*. The head resembles that of a *Scorpæna*; the profile scarcely descending at all. The eyes of middling size, are separated by an interval of double their diameter; the lower jaw ascends obliquely to meet the other: there are soft teeth in the jaws, before the vomer, and in the palate; the orbit and the cranium have only slight bones without points. The under anterior orbital has two sharp points, of which the upper, rather longer than the other, does not go beyond the under part of the middle of the orbit. The great sub-orbital is only marked by some projecting wrinkles. The preoperculum is rounded, has a very short spine, followed by three small flat teeth; the operculum has two bones and two little points: there are only six rays to the gills, the membrane of which is much sloped. The first dorsal spine ascends above the edge of the preoperculum, and the two following proceed almost from the same points; the fourth is on the anterior third of the pectoral; the fifth and last on the anterior third of the anal; all are straight, very strong, and nearly uniform. The end of the dorsal is only composed of four branchial rays, of which the fourth is attached above the tail the whole of its length, without, however, quite extending to the caudal. The Anal has three spines, and also four branchial rays, of which the last attaches itself like that of the dorsal,

PISCES.	Vol. II. Page
2nd. Plate 28. quar. Fig. 3.— <i>APISTUS MONODACTYLUS</i> , Cuv.; <i>Scorpæna mono-</i> <i>dactyla</i> , Bl. and Schn.	106
Plate 29. Fig. 1.— <i>SCIENA AQUILLA</i> ; <i>S. umbra</i> , Cuv.	109
Fig. 2.— <i>CORVINA FURCA</i> , Cuv.	110
Fig. 3.— <i>JOHNII SENEGALENSIS</i> , Cuv.	110
Fig. 4.—Head of the <i>OTOLITHUS RUBER</i> , Cuv.	109
Fig. 5.—Head of the <i>ANCYLODON PARVIPINNIS</i> . A new species from the Coast of Cayenne.	110
Fig. 6.—Head of the <i>UMBRINA COROIDES</i> . A new species from Brazil. 110	
Fig. 7.—Head of the <i>POGONIAS FASCIATUS</i>	111
2nd. Plate 29. Fig. 1.— <i>NOMEUS PERONII</i> (<i>The Shepherd Fish</i>)	131
Fig. 2.— <i>CARANX BOOPS</i> (<i>The False Mackerel or Carungul</i>)	132
Fig. 3.— <i>GALLUS AEGYPTIACUS</i> (<i>The Egyptian Gallus</i>)	133
Plate 29. bis. Fig. 1.— <i>OTOLITHUS TEROE</i> , Cuv.	109
Fig. 2.— <i>OTOLITHUS GUATUCUPA</i> ; <i>Otolithus striatus</i> , Cuv. (<i>The Striped Otolithe</i>)	109
Fig. 3.— <i>CORVINA LOBATA</i> , Cuv.	109
Plate 29. ter. Fig. 1.— <i>CHEILODACTYLUS CARPONENNIS</i> , Cuv.	112
Fig. 2.— <i>SCOLOPSIDES LYCOGENIS</i> , Cuv.	113
Fig. 3.— <i>DIAGRAMMA GATERINA</i> , Cuv.	112
2nd. Plate 29. ter. Fig. 1.— <i>CHÆTODON STRIGANGULUS</i> , Sol.	120
Fig. 2.— <i>CHÆTODON LUNULA</i>	120
Fig. 3.— <i>CHÆTODON EPHIPPUM</i> , Cuv.*	120
Plate 30. Fig. 1.— <i>EQUES BALTEATUS</i>	111
Fig. 2.— <i>LOBOTES ERATE</i>	112
Fig. 3.—Head of the <i>HÆMULION HETERODON</i>	111
Fig. 4.—Head of the <i>PRISTIPOMA BILINEATUM</i>	112
Fig. 5.—Head of the <i>DIAGRAMMA ORIENTALE</i>	112
Plate 30. bis. Fig. 1.— <i>LEIOSTOMUS HUMERALIS</i> ; or <i>Johnius humeralis</i> , Cuv.. 110	
Fig. 2.— <i>NEBRIS MICROPS</i> , Cuv. (<i>The Small-eyed Nebris</i>)†	110
Fig. 3.— <i>CORVINA DENTEX</i> (<i>The Many-tooth Corvina</i>)	110

but remains rather further from the caudal: the latter is rounded and composed of twelve entire rays, with some shorter ones above and below. The pectoral, very sharp, of less than a quarter of the total length, has twelve rays, all branched. The ventral composed as usual, and half attached, is one fourth shorter.

The body of this little fish when it begins to dry, shows some very small scales, as if absorbed in the skin; its epidermis forms very fine vertical wrinkles. The lateral line is straight, and on the upper fifth, is marked by oblique tubercles. The general colour appears to be a red brown, dotted with white on the sides of the head and under the jaw, which changes to a silvery white under the throat and breast. The dorsal is crossed almost vertically by four or five irregular blackish bands, among which are small grey lines. The anal has two or three of these bands and the same lines between them. There is one at the base of the caudal; on the rest of which are lines of brown dots, slightly marked on a white ground. The pectoral is also crossed by lines of dots or little brown spots; the ventral is whitish, and at the end black; sometimes it has also a blackish spot crosswise. The individual from which our figure is taken, measured two inches and a half long.

* A new species found by M.M. Lesson and Garnot, near one of the Society Islands.

† A new species from the neighbourhood of Surinam, which differs in a few sim-

PISCES.	Vol. II. Page
2nd Plate 30. bis. Fig. 1.— <i>SARGUS ANNULARIS</i> , Cuv.	115
Fig. 2.— <i>SARGUS RHOMBOIDES</i> , Cuv.	115
Fig. 3.— <i>CHARAX PUNTAZZO</i> , Cuv.; <i>Sparus Puntazzo</i> Gmelin, (<i>The Common Puntazzo</i>)	115
Plate 30. ter. Fig. 1.— <i>ZANCLUS CORNUTUS</i> ; <i>Chætodon cornutus</i> , Lin.	121
Fig. 2.— <i>HENIOCHUS MONOCEROS</i> , (<i>The Coachman</i>)	121
2nd. Plate 30. ter. Fig. 1.— <i>CHRYSOPHRYS GIBBICEPS</i> , Cuv.*	115
Fig. 2.— <i>CHRYSOPHRYS CRASSIROSTRIS</i> , Cuv.†	115
Fig. 3.— <i>CHRYSOPHRYS AURATA</i> , Cuv. (<i>The Golden eye-brow</i>)	115
Plate 30. quar. Fig. 1.— <i>SCATOPHAGUS ORNATUS</i> , Cuv.‡	121
Fig. 2.— <i>TAURICHTHYS VARIUS</i> , (<i>The Horseman of India</i>)	121
Fig. 3.— <i>HOLACANTHUS TRIMACULATUS</i> ,	122

ple particulars from the Corvina. It forms the type of a new subdivision in the Ichthyology of our author.

* This is a new species of Dory, brought from the Cape by M. Delalande, and is strikingly separated from the others, by the extreme elevation of the crest of the cranium. A very excellently preserved specimen of this fish, may be seen among the anatomical collection of the Royal College of Surgeons.

† Individuals of this species were taken in the Mediterranean on the coast of Corsica, near Ajaccio, by M. Payraudeau, and on the coasts of the Peloponesus by M. Bory St. Vincent. We find a figure of them also among some plates engraved in Spain, and which have not yet, we believe, been published; so that though more rare than the *Sparus Auratus*, Lin., it seems to inhabit the same parts. It is known by the thickness of its snout, as well as by the greater length of its body, its height being contained three times and a half in its length; the neck is much more elevated in proportion; the eyes are larger; the space between the eyes wider and projects more; the lower edge of the under orbital is more arched; the preoperculum more rounded; its border straighter, but equally bare. The interoperculum is much larger, and crosses under the throat with that of the opposite side; the jaws are much swelled; there are six large incisors, and four rows of grinders in the upper jaw, and three in the under; they are larger and wider apart; the oval tooth is more rounded; the dorsal is lower; the pectoral rather wider. The ventral rays are very strong, and extend themselves in the form of a large fan. The number of rays are as follows: D. 11—13; A. 3—11, C. 17; P. 15; V. 1—5. The scales are close, higher than wide; we have counted more than eighty, from the gill to the end of the tail; the lateral line is almost straight. The individual we saw, was nearly as fresh as when taken out of the water; the colour is deep blue on the back with bright gold reflexions; these reflexes proceeded from a golden streak traced on each scale. On the shoulder, and on the top of the operculum, is seen a large black spot. The front of the cranium is blue; the face is bluish, with copper reflexes. Between the eyes there is a crescent more arched than that of the common Dory, and of the most beautiful colour of polished gold. On each temple in front of the superscupular, is a very beautiful spot, as brilliant as the crescent. The under part of the orbit is also gilt, as well as the border of the preoperculum. The lower part of the operculum, and the lower jaw, are similar to the red of polished and brilliant copper; the fins appear bluish grey. The individual obtained near the coast of Corsica, was about eighteen inches long; that of the Morea only a foot, and yet its large oval tooth was already out. None of the authors, we have consulted, appear to have any knowledge of this beautiful Dory, which is equal to the other in the brilliancy of its colours.

‡ A new species found by MM. Quoy and Gaimard near Amboyna, and figured by Cuvier and Val., in their Ichthyology under this name. It forms part of a new subdivision, in which is embraced the *Chatodon tetricanthus* of Lacepede.

PISCES.	Vol. II. Page
Plate 31. Fig. 1.— <i>SMARIS ALCEDO</i> , Riss. (<i>The Blue-spotted Picarel</i>)	118
Fig. 2.— <i>CÆSIO TILE</i> , Cuv.	118
Fig. 3.—Head of the <i>MÆNA VOMERINA</i> , Cuv. (<i>Vomerine Mendole</i>) A new species from the Mediterranean	118
Fig. 4.—Head of the <i>GERRES PLUMIERI</i> , Cuv.	119
 Plate 31. bis. Fig. 1.— <i>PAGELLUS LITHOGNATHUS</i> , Cuv.; <i>Pagrus lithognathus</i>	116
Fig. 2.— <i>PAGELLUS ERYTHRINUS</i> , Cuv. (<i>The Pagel</i>)	116
Fig. 3.— <i>PAGELLUS CALAMUS</i> , Cuv.	116
 Plate 31. ter. Fig. 1.— <i>LETHRYNUS GENIVITTATUS</i> , Cuv.	117
Fig. 2.— <i>CANTHARUS VULGARIS</i> , Cuv.	117
Fig. 3.— <i>BOOPS VULGARIS</i> , Cuv.	117
 Plate 31. quar. Fig. 1.— <i>HOLACANTHUS SEMI-CIRCULATUS</i> , Cuv.	122
Fig. 2.— <i>HOLACANTHUS LAMARCKII</i> , Lacep.	122
Fig. 3.— <i>POMACANTHUS CINGULATUS</i> , Cuv.	122
 Plate 32. Fig. 1.— <i>CAZETODON SETIFER</i> , Cuv.	121
Fig. 2.— <i>EPHISSUS ORBIS</i> , Cuv. (<i>The Horseman</i>)	121
Fig. 3.—Head of the <i>EPHISSUS LONGIROSTRIS</i> ,	121
 Plate 32. bis. Fig. 1.— <i>DENTEX PERONII</i> , Cuv. (<i>Peron's Dentex</i>)	116
Fig. 2.— <i>DENTEX FILAMENTOSUS</i> , Cuv.	116
Fig. 3.— <i>DENTEX VULGARIS</i> , Cuv. (<i>The Common Dentex</i>)	116
 Plate 32. ter. Fig. 1.— <i>CHELMON LONGIROSTRIS</i> , Cuv. (<i>The Long-mouthed Chelmon</i>)	121
Fig. 2.— <i>ASPIDOPHORUS MONOPTERYGIUS</i> ; <i>Cottus Monopterygius</i> , Bl. (<i>The Aspidophorus with a single Dorsal</i>)	104
Fig. 3.— <i>APRION VIRESSENS</i> , Cuv.*	104
 2nd. Plate 32. ter. Fig. 1.— <i>SPIROBRANCHUS CAPENSIS</i> , Cuv.	145
Fig. 2.— <i>TRICHOPUS TRICHOPTERUS</i> , Lacep.	145
Fig. 3.—Anatomy of the Brain of the <i>EPHISSUS GIGAS</i> , (<i>The Gigantic Horseman</i>)	121
Fig. 4.—Shoulder bone,	121
Figs. 5, 6.—Interspinal and Anal bones of the <i>EPHISSUS GIGAS</i>	121
 Plate 32. quar. Fig. 1.— <i>ANABAS SCANDENS</i> , Cuv.; <i>Pereis scandens</i> , Dald.	144
Fig. 2.— <i>PSETTUS SEBÆ</i> , Cuv.	122
Fig. 3.—The Labyrinthinal Organ of the <i>OPHICEPHALUS STRIATUS</i> , Bl.	146
Fig. 4.—Anatomy of the Brain of the <i>OSPHROMENUS GOURAMI</i> , Lacep.	145
 Plate 33. Fig. 1.— <i>HOLACANTHUS CILIARIS</i> , Cuv.	122
Fig. 2.— <i>POMACANTHUS AUREUS</i> , Cuv.	122
 2nd. Plate 33. Fig. 1.— <i>SCOMBER SCOMBRUS</i> , (<i>The Common Mackerel</i>)	124
Fig. 2.— <i>THYNNUS VULGARIS</i> , (<i>The Tunny</i>)	125
Fig. 3.— <i>AUXIS PELEI</i> , (<i>The Auxis</i>)	126

* This is a new species which combines several of the characteristics of the great family of Perches with a single dorsal, with many of the peculiarities of the *Spa-roides*. It is fully described by Cuvier and Val., in their *Ichthyology*, vol. vi, p. 544, &c. Its place in this arrangement, is immediately after *Aspidophoræ*.

PISCES.	Vol. II. Page
Fig. 4.—The mouth with the teeth of the <i>AUXIS</i> ,	126
Fig. 5.—The mouth of <i>PELAMYS SARDA</i> ,	126
 Plate 33. bis. Fig. 1.— <i>PENTAPUS AUROLINEATUS</i> , Cuv. (<i>The Golden-striped Pentapode</i>)	117
Fig. 2.— <i>PENTAPUS POROSUS</i> , Cuv.	117
Fig. 3.— <i>LETHRYNUS ESCULENTUS</i> , Cuv.	117
 Plate 33. ter. Fig. 1.— <i>OPHICEPHALUS LIMBATUS</i> ; or <i>Marginatus</i> , Cuv.	146
Fig. 2.— <i>OPHICEPHALUS STRIATUS</i> , Bl.	146
Fig. 3.— <i>OPHICEPHALUS GRANDINOSUS</i> , Cuv.	146
 Plate 33. quar. Fig. 1.— <i>COLISA VULGARIS</i> ; <i>Trichopodus Colisa</i> , Buchanan	144
Fig. 2.— <i>MACROPODUS VENUSTUS</i> , Cuv. A new species,	144
Fig. 3.— <i>PLATAX GUTTULATUS</i> , Cuv.	122
Fig. 4.— <i>PLATAX PUNCTULATUS</i> , Cuv.	122
 Plate 34. Fig. 1.— <i>PLATAX GAIMARDI</i> , Cuv.; a new species found by M. Gaimard, near New Guinea,	122
Fig. 2.— <i>PSETTUS RHOMBEUS</i> , Cuv.	122
 Plate 34. bis. Fig. 1.— <i>EPHISSUS GOREENSIS</i> , Cuv. A new species nearly allied to the <i>Gigas</i>	121
Fig. 2.— <i>DREPANE PUNCTATA</i> , Cuv.; <i>Chætodon punctata</i> , Lin.	121
Fig. 3.—Head of the <i>SPIROBRANCHE</i> ,	145
Fig. 4.—Head of the <i>MACROPODE</i> ,	144
Fig. 5.—Head of the Common <i>COLISA</i> ; <i>Trichopodus Colisa</i> , Buchanan,	144
 Plate 34. ter. Fig. 1.— <i>THYNNUS PELAMYS</i> , Cuv. (<i>The Tropical Bonita, or Ray-bellied Tunny</i>)	125
Fig. 2.— <i>PELAMIS SARDA</i> , Cuv. (<i>The Common Pelamis</i>)	126
Fig. 2 a.—Dentition of the <i>PELAMIS SARDA</i> ,	125
Fig. 3.— <i>THYNNUS THUNNINA</i> , Nob. (<i>The Thonine</i>)	125
Fig. 4.—Head of the <i>XIPHIAS GLADIUS</i> , Lin. (<i>The Sword-fish</i>) a young subject	127
Fig. 5.—The same as fig. 4., but taken from a full grown Fish,	127
 Plate 34. quar. Fig. 1.— <i>PIMELEPTERUS BOSCII</i> , Lacep. (<i>The Fat-fin</i>)	123
Fig. 2.— <i>POLYACANTHUS HASSELTI</i> , Cuv. A new species discovered by Van Hasselt, on the coast of Java.	144
Fig. 3.— <i>BRAMA RAI</i> , Bl. and Schn. (<i>The Sea Bream</i>)	123
 Plate 35. Fig. 1.— <i>PIMELEPTERUS ALTIPINNIS</i> .*	123
Fig. 2.— <i>DIPTERODON CAPENSIS</i> , (<i>The Cape Dipteronodon</i>)	123
Fig. 3.—Head of the <i>PIMELEPTERUS BOSCII</i> ; <i>Pimeleptere bosquier</i> , Lacep.	123
 Plate 35. bis. Fig. 1.— <i>THYRSITES ATUN</i> , Cuv.; <i>Scomber atun</i> , Euph. and Lacep.	127
Fig. 2.— <i>CYBIUM TRITOR</i> , Cuv. A new species which approaches near to the <i>Cybum Commersoni</i> ,	127
Fig. 3.— <i>THYRSITES LEPIDOPOIDES</i> . A new species from the coast of Brazils	127

* A new species discovered by MM. Quoy and Gaimard, near the coast of New Guinea, which is remarkable for the height of the spines of the dorsal and anal fins; in other respects it is similar to the *P. Boscii*.

PISCES.		Vol. II. Page
Plate 35. ter. Fig. 1.— <i>GEMPYLUS PROMETHEUS</i> , Cuv.	.	127
Fig. 2.— <i>LEPIDOFUS ARGYREUS</i> , Cuv.	.	127
Fig. 3.— <i>GEMPYLUS COLUBER</i> , Cuv.	.	127
Plate 35. quar. Fig. 1.— <i>HELOSTOMA TEMMINCKII</i> , Cuv.	.	144
Fig. 2.— <i>PEMPHERIS TAITENSIS</i> , Cuv.	.	124
Fig. 2.— <i>TOXOTES JACULATOR</i> , Cuv. (<i>The Archer Fish</i>)	.	125
Plate 36. Fig. 1.— <i>BRAMA RAI</i> , Bl. (<i>The Sea Bream</i>). For another view, see Pl. 34. quar. fig. 3.	.	123
Fig. 2.— <i>PEMPHERIS MEXICANA</i> , Cuv.	.	123
Fig. 3.— <i>TOXOTES JACULATOR</i> , Sch. (<i>The Archer Fish</i>). For another view, see Pl. 35. quar. fig. 3.	.	125
Plate 36. ter. Fig. 1.— <i>GERRES PLUMIERI</i> . A new species discovered in the Atlantic, by Plumier.	.	119
Fig. 2.— <i>CÆSIO ERYTHROGASTER</i> ,*	.	118
Fig. 3.— <i>SMARIS CHRYSSELIS</i> ,†	.	118
Plate 37. ter. Fig. 1.— <i>ELACATE ATLANTICA</i> ; <i>Centronotus spinosus</i> , Mitchell	129	
Fig. 2.— <i>LICHIA GLAYCOS</i> ; or, <i>L. gluaca</i> , Cuv.	.	129
Fig. 3.— <i>LICHIA VADIGO</i> , Rondel	.	129
Plate 37. quar. Fig. 1.— <i>THYNNUS ALALONGA</i> , (<i>The Germon</i>)	.	125
Fig. 2.— <i>THYNNUS BRACHYPTERUS</i> , Cuv.	.	125
Fig. 3.— <i>TAYNNUS BREVIPINNIS</i> ,‡	.	125
Plate 38. Fig. 1.— <i>CYBİUM LINEOLATUM</i> , Cuv. (<i>The long Tassard</i>)	.	126
Fig. 2.— <i>THYRSITES CHILENSIS</i> , (<i>The White Thysites</i>)	.	127
Fig. 3.— <i>GEMPYLUS PROMETHEUS</i> , (<i>The Gempyle</i>).§. For another view, see Pl. 35. ter. fig. 1.	.	127
Plate 38. bis. Fig. 1.— <i>OLISTUS MALABARICUS</i> , Cuv. and Val.	.	133
Fig. 2.— <i>SCYRIS INDICA</i> , Cuv. and Val.	.	133
Plate 38. ter. Fig. 1.— <i>APOLECTUS STROMATEUS</i> , Cuv.	.	129
Fig. 2.— <i>CHORINEMUS SANCTI PETRI</i> , Cuv.¶	.	129

* A new species discovered by MM. Kuhl and Van Hasselt, in the Indian Ocean. It is very similar to the *C. sinaris*, but not the same.

† Another species, which has been found in the Mediterranean, near to the coast of Sicily.

‡ The Mediterranean produces this new species of Tunny, which resembles the common one in every respect, except that the pectorals are much shorter.

§ A new species found near St. Helena, by MM. Quoy and Gaimard. It approaches closely to the *G. serpens* of Cuvier.

|| A small fish, specimens of which have been brought from Pondicherry by M. Sonnerat, and from the coast of Java by MM. Kuhl and Van Hasselt, which very much resembles a *Trichinotus*; except that the ventrals are attached to the under part of the throat, a circumstance which Cuvier, in his work on fishes, has thought sufficient to rank it as the type of a distinct genera. Its place should, therefore, be immediately after the *Trichinotus*, page 129.

¶ M. de Lacepede has called by the name of *Scomberoides*, those *Lichia* resembling the rest in the greater part of their organs; but of which the rays of the second dorsal or of the anal are either entirely detached, or united by a membrane so fragile, that it often entirely disappears; and they seem to form false fins, like those we have observed in the Mackerel and Tunnies. Not being able to retain the name of *Scomberoides*, since it would be so similar to that of the family, Cuvier has affixed to them, that of *Chorinemus*; one of which, the—*C. Sancti Petri*, a beautiful and large fish, was brought by M. Dussumier from the coast of Malabar, and is remarkable by a double range of spots on each flank. Its height is the

PISCES.	Vol. II.	Page
Plate 38. ter. Fig. 3.— <i>TRACHINOTUS PAMPANUS</i> , Cuv.*	.	129
Plate 38. quar. Fig. 1.— <i>TETRAPTERUS INDICUS</i> , Cuv.	.	128
Fig. 2.—Skeleton of the <i>TETRAPTERUS BELONE</i> , Ra.	.	128
Plate 39. Fig. 1.— <i>XIPHIAS GLADIUS</i> , Lin. (<i>The Common Sword-Fish</i>)	.	127
Fig. 2.—Same Fish, but of a younger age,	.	127
Fig. 3.— <i>TETRAPTURUS BELONE</i> , Raf.	.	128
Fig. 4.— <i>HISTIOPHORUS PULCELLUS</i> .	.	128
Plate 39. ter. Fig. 1.— <i>RHYNCHICHTYS PELAMIDIS</i> , Cuv.†	.	94
Fig. 2.— <i>NANDUS MARMORATUS</i> , Cuv.‡	.	94
Fig. 3.— <i>SCOMBER COLIAS</i> , Gm. (<i>The Mediterranean Mackerel</i>)	.	125
Plate 40. Fig. 1.— <i>NAUCRATES INDICUS</i> . A new species, which approaches closely to the <i>Scomber ductor</i> , Lin.	.	129
Fig. 2.— <i>MASTACEMBELUS MACULATUS</i> . It is <i>Rhynchobdella maculata</i> Reinwardt.	.	130
Fig. 3.— <i>SERIOLA RIVOLII</i> §	.	130
Fig. 4.—Head of the <i>ELACATE ATLANTICA</i> ; <i>Centronotus spinosus</i> , Mitchell	.	129
Fig. 5.—Head of the <i>RHYNCHOBDELLA ARAL</i>	.	130
Fig. 6.—Head of the <i>MASTACEMBELUS ARMATUS</i> ; <i>Macrognathus armatus</i> , Lacep.	.	130
Fig. 7.—Head of the <i>NOTACANTHUS NASUS</i> , Bl.	.	130
Plate 40. bis. Fig. 1.— <i>BOVICHTUS DIACANTHUS</i> , Cuv.	.	94

fifth of its length, its thickness one-third of its height. Its head one-fourth less high than long, is a fifth of the total length.

* A new species found near Brazils, and other parts of South America.

† It is to the researches of M. Dussumier, that we owe the knowledge of this singular little fish, which he found in the stomach of a Tropical Bonita in the Indian sea. It belongs to the group of Perches with eight rays to the gills and to the ventrals, consequently it is near the Holocentri; but it constitutes a new genus, known by a prolongation of the carina of the cranium in a point which advances beyond its mouth, almost the same as in the *Lepidoleprus*. The preoperculum has a projecting spine towards its angle; but the angle of the operculum has only very short spines, as in the *Myripristis*. We as yet only know one single species of this genus, which we have named *Rhynchichtys pelamidis*.

‡ There must be added to the division of "Percoides with a single dorsal and six branchial rays, a fish very common in the lakes of Bengal; it is the *Coius nandus* of Buchanan. The description given by this author, however, leaves us too many doubts to be able to fix with correctness the place of this species; but M. Dussumier, to whom science is so much indebted, has recently brought to Europe a *nandus* as fresh as if it were just out of the water; which enables us to give a detailed and exact description of it. This fish we shall now make the type of a genus, to follow immediately after the Doules, p. 94. It is characterised by a very protractile mouth, furnished with fine short soft teeth, like the pile on velvet, in the two jaws, the palate and vomer. The preoperculum and interoperculum have their edges finely indented. The spine of the operculum is so small that it is difficult to perceive it. The protractility of the upper jaw of this fish gives it an appearance quite different from the Doules, and makes it more resemble a Mendole with a shortened body. This affinity is increased, as the Mendoles, like this fish, have some teeth in the palate, but fewer in number, and very few in the vomer. However the *Nandus* cannot be placed in the family of the Menides, on account of the indentations of the operculum. We know but one kind of them—the *Nandus marmoratus*.

§ A species which approaches closely the *Seriola cosmopolita* of Cuvier. It was presented to the Cabinet Du Roi by the Duke of Rivoli.

|| We sought for a long time to know what the jugular fish could be, figured in the

PISCES.		Vol. II. Page
Plate 40. bis. Fig. 2.— <i>APHRITIS URVILLII</i> , Cuv.*	.	94
Fig. 3.— <i>APLODACTYLUS PUNCTATUS</i> , Cuv.†	.	94
Plate 41. bis. Fig. 1.— <i>NOTACANTHUS NASUS</i> , Bl.	.	130
Fig. 2.— <i>MASTACEMBELLUS ARMATUS</i> , Cuv.; <i>Macrognathus armatus</i> , Lacep.	.	130
Fig. 3.— <i>RHYNCHOBDELLA ARAL</i> , Cuv.	.	130

twelfth volume of the Linnaean transactions by Captain Carmichael, under the name of *Callionymus diacanthus*, for it was easy to perceive that it was not a *Callionymus*. We have been fortunate enough to find it among the species collected at Valparaiso, from Chili, by M. d'Orbigny. It is a particular genus, near the *Trachinus*, having like them teeth as the pile on velvet in the jaws, in the palate and on the vomer, but is distinguished not only from the *Trachinus*, but also from all the other jugular *Percoides*, the *Percophis* excepted, by the seven rays of its branchiostegal membrane. The colour appears to have been blackish. The length of the individual is eight inches, the caudal not included. Captain Carmichael says his fish had the body olive colour marbled with greenish spots, and marked with white dots, and the iris of the eye brown. On his figure (pl. 26. vol. 12.) of the Linnaean transactions, twenty two rays are marked to the second dorsal; but in the text the author only enumerates twenty rays. The characters of the penultimate and antepenultimate rays prolonged from the anal, is well expressed in the figure that we quote.

* Cuvier found in the lot of fishes brought to Europe by Messrs. Quoy and Gaimard three small *Percoides* with jugular ventrals, originally from the soft waters of Van Dieman's land, which are all three of the same species, but which must become the type of a new genus, near that of the *Percophis*. (p. 94.) The name of *Aphritis* has been given to the genus, and that of *Aphritis Urvillii* to the species figured by us.

This species has a cylindric body, of which the diameter, in front of the anus, is the sixth of the total length; the colour is reddish, shaded, and marbled with greenish brown on the back. The fins are transparent. On the two dorsals, and on the caudal, we see two or three rows of small red dots.

† This fish forms a new genus of the family of *Percoides* with six branchial rays, and free rays to the pectorals, consequently near the *Cirrhites*; but it is easily distinguished from them, by the form of the teeth, which resemble those of the *Crenidens* Forsk., of the family of *Sparoides*. We give to this genus, the name of *Aploactylus*, to recall the simple form of some of the pectoral rays, and to the species we have figured that of *Aploactylus punctatus*. Its height measures the fourth of its total length. The thickness is contained twice and a half in the height; the snout is obtuse and rounded. The line of the profile of the head, ascends by an arched curve, sustained even to the occiput, from whence it rises, and ascends obliquely, following an almost straight direction, even to the base of the dorsal. The curve of the back is slight; that of the belly is rather more so. The height of the tail is only the third of that of the body, measured perpendicularly from the ventrals. The head is short, and swelled in front. It has its length contained four times and four-fifths in that of the body. Its height from the neck downward, is a fifth less than its length. The eyes, of moderate size, are on the first half of the head, and placed in the upper part of the cheek, without, however, the circle of the orbit encroaching on the line of the profile.

The scales are very small; there are more than a hundred and twenty between the gill and the caudal—seen separately, and through a magnifying glass, each of them presents a square surface, finely striated on the sides, and of which the radical edge is finely indented. The lateral line is traced beneath the third of the height, and forms slight undulations. On a white ground, brownish towards the back, the body is spotted with an infinite number of little blackish dots, equally spread over the fins, which are darker than those on the back.

M. Gay has likewise observed this fish; for we have found a coloured drawing of it among the drawings of this naturalist, which leads us to hope that the species is not very rare, and that we may receive other individuals.

PISCES.

	Vol. II. Page
Plate 42. Fig. 1.— <i>ZEUS FABER</i> , Lin. (<i>The Common Dory</i>)	133
Fig. 2.— <i>LAMPRIS GUTTATUS</i> , Retz. (<i>The Moon Fish of Lacepede</i>)	134
Fig. 3.— <i>STROMATEUS FIATOLA</i> , Lin. (<i>The Mediterranean Pomfret</i>)	135
Plate 42. bis. Fig. 1.— <i>ARGYREYESUS VOMER</i> , Lacep. ; <i>Zeus vomer</i> , Lin.	133
Fig. 2.— <i>CARANX TRACHURUS</i> , Lacep. ; <i>Scomber trachurus</i> , Lin.	131
Fig. 3.— <i>CARANX CHRYSOPHRYNS</i> , Cuv. A new species.	132
Plate 42. ter. Fig. 1.— <i>BLEPHARIS SUTOR</i> , Cuv. (<i>The Cordonnier of Martinique</i>)	133
Fig. 2.— <i>GALLICHTYS MAJOR</i> . A new species of <i>Gallus</i> nearly allied to the <i>Zeus gallus</i> , Lin.	133
Plate 43. Fig. 1.— <i>CORYPHENA AZORICA</i> , Cuv. and Val.	137
Fig. 2.— <i>PTERACLIS TRICHIPTERUS</i> , Cuv. and Val.	139
Plate 44. Fig. 1.— <i>GYMNETRUS FALX</i> , Cuv.	139
Plate 45. Fig. 1.— <i>SIGANUS DOLIATUS</i> , Cuv. and Val.	142
Fig. 2.— <i>ACANTHURUS DELISIANUS</i>	142
Fig. 3.— <i>NASEUS LONGICORNIS</i>	142
Plate 46. Fig. 1.— <i>ANABAS TESTUDINEUS</i> , Bl. (<i>The Tree Climber</i>)	145
Fig. 2.— <i>POLYACANTHUS COLISA</i> , Ham.	144
Fig. 3.— <i>OPHICEPHALUS SERPENTINUS</i> . A new species found by Dr. Finlayson, near the coast of Siam	146
Plate 47. Fig. 1.— <i>MUGIL CEPHALUS</i> , Cuv. (<i>The Common Mullet</i>)	146
Fig. 2.— <i>TETRAGONURUS CUVIERI</i> , Risso.	148
Fig. 3.— <i>ATHERINA PRESBYTER</i> , Cuv.	149
Plate 48. Fig. 1.— <i>BLENNIUS PALMICORNIS</i> , Cuv.	150
Fig. 2.— <i>CLINUS PECTINIFER</i>	151
Fig. 3.— <i>ZOARCES BLENNOIDES</i>	152
Plate 49. Fig. 1.— <i>GOBIUS CAPITO</i> , Cuv. (<i>The Great Goby or Sea Gudgeon</i>)	153
Fig. 2.— <i>GOBIOIDES BROUSSONNETII</i> , Lacep. (<i>The great Gobioides of Broussonnet</i>)	154
Fig. 3.— <i>PERIOPHTHALMUS FREYCINETII</i>	154
Plate 50. Fig. 1.— <i>ELEOTRIS DORMITATRIX</i> , Cuv. (<i>The Sleeper</i>)	155
Fig. 2.— <i>CALLIONYMUS FASCIATUS</i> , Cuv. and Val. (<i>The Fasciated Callionymus</i>). A new species.	156
Fig. 3.— <i>CHIRUS DEAGRAMMUS</i> , Pall. (<i>The Ten-striped Chirus</i>)	157
Plate 51. Fig. 1.— <i>LOPHIUS PISCATORIUS</i> , Lin. (<i>The Common Angler, or Frog-Fish</i>)	158
Fig. 2.— <i>ANTENNARIUS NESOGALLOVICUS</i> . A new species.	159
Fig. 3.— <i>Batrachus SURINAMENSIS</i> , Bl.	160
Plate 52. Fig. 1.— <i>LABRUS VITTATUS</i> , Quoy and Gaym. A new species.	161
Fig. 2.— <i>JULIS TRIMACULATA</i> , Quoy and Gaym. A new species.	162
Fig. 3.— <i>CRENILABRUS CHABROLII</i> , Less. A new species.	163
Plate 53. Fig. 1.— <i>CHROMIS NILOTICA</i> , Cuv. (<i>The Egyptian Coracinus</i>)	165
Fig. 2.— <i>MALACANTHUS FURCATUS</i> . Cuv. A new species.	166
Fig. 3.— <i>SCARUS CREVICUS</i> , Aldrov. and Cuv. (<i>The Parrot-Fish</i>)	167
Plate 54. Fig. 1.— <i>EPIBULUS INSIDIATOR</i> , Pall.	164
Fig. 2.— <i>GOMPHOSUS VIRIDIS</i> , Cuv.	164
Fig. 3.— <i>XIRICHTHYS CYANIROSTRIS</i> , Cuv. A new species	165

PISCES.	Vol. II. Page
Plate 55. Fig. 1.— <i>FISTULARIA TABACARIA</i> , Bl. (<i>The Tobacco-pipe Fish, also called Flute-mouth</i>)	168
Fig. 2.— <i>CENTRISCUS SCOLOPAX</i> , Lin. (<i>The Sea-Snipe</i>)	169
Fig. 3.— <i>AMPHISILE SCUTATUS</i> , Lin. (<i>The Shielded Sea-Snipe</i>)	169
Plate 56. Fig. 1.— <i>CYPRINUS AMARAS</i> , (The <i>Amarous Carp</i>)	171
Fig. 2.— <i>GOBIO VULGARIS</i> , (The <i>Common Gudgeon</i>)	172
Fig. 3.— <i>LABEO NILOTICUS</i> , Geoff. (<i>The Mullet of the Nile</i>)	173
Plate 57. Fig. 1.— <i>LEUCISCUS IDUS</i> , Bloch. (<i>The Albe, commonly called the White Fish</i>)	173
Fig. 2.— <i>COBITIS BARBATULA</i> , Lin. (<i>The Common Bearded Loach</i>)	175
Fig. 3.— <i>PÆCILIA UNIPUNCTATA</i> , Val. (<i>The One-dotted Pæcilia</i>)	176
Plate 58. Fig. 1.— <i>LEBIAS FASCIATA</i> , Val. (<i>The Fasciated Lebias</i>)	176
Fig. 2.— <i>FUNDULUS BRASILIENSIS</i> , Val. (<i>The Fundule of Brazil</i>)	176
Fig. 3.— <i>CYPRINODON FLAVULUS</i> , Val.	177
Fig. 4.—Anatomical details of the <i>FUNDULUS FASCIATUS</i> , Val. (<i>The Fasciated Fundule</i>)	176
Fig. 5.—Dentition of the <i>LEBIAS RHOMBOIDALIS</i> , Val. (<i>The Rhomboid Lebias</i>)	176
Plate 59. Fig. 1.— <i>ESOX LUCIUS</i> , Lin. (<i>The Common Pike</i>)	177
Fig. 2.— <i>MICROSTOMA MEDITERRANEA</i> , Risso (<i>The Mediterranean Microstoma</i>)	178
Fig. 3.— <i>STOMIAS BARBATUS</i> , Risso (<i>The Barbed Stomias</i>)	178
Plate 60. Fig. 1.— <i>SALANX FUSCUS</i> , Cuv. and Val.	179
Fig. 2.— <i>SCOMBRESOX SAURUS</i> , Bl.	180
Fig. 3.— <i>EXOCETUS ACUTUS</i> . (<i>The Sharp-snouted Flying-Fish</i>). A new species	180
Plate 61. Fig. 1.— <i>MORMYRUS OXYRHINCHUS</i> , Geoff. Cuv. (<i>The True Mormyrus oxyrhinchus of the Ancient Egyptians</i>)	182
Fig. 2.— <i>SCHILBE MYSTUS</i> , Geoff. (<i>The Silure of the Nile</i>)	183
Fig. 3.— <i>SYNODONTIS SCHEILAN</i> , Geoff. (<i>The Schal of Egypt</i>)	185
Plate 62. Fig. 1.— <i>BAYARD FITILE</i> , Cuv. and Val. A new species	185
Fig. 2.— <i>DORAS DORSALIS</i> , Cuv. and Val. A new species	186
Fig. 3.— <i>HETEROBRANCHUS PUNCTATA</i> , Cuv. and Val. A new species	186
Plate 63. Fig. 1.— <i>PORGUS BAYAD</i> , Geoff. and Cuv.	184
Fig. 2.—Mouth of the <i>PIMELODES BLOCHII</i> , Cuv. and Val.	185
Fig. 3.— <i>PLOTOSUS ALBILABRIS</i> , Cuv. and Val. A new species	187
Fig. 4.— <i>CALlichthys LONGIFILIS</i> , Cuv. and Val.	187
Plate 64. Fig. 1.— <i>ASPREDO OCTOCIRRUS</i> , Cuv. A new species	188
Fig. 2.— <i>HYPSTOMUS PLEIDOMUS</i> , Cuv.	189
Fig. 3.— <i>LORICARIA CATAPHRACTA</i> , Lin.	189
Plate 65. Fig. 1.— <i>SALMO FARIO</i> , Lin. (<i>The Common Trout</i>)	191
Fig. 2.— <i>MALLOTUS GRAELANDICUS</i> , Bl. Cuv. (<i>The Capelan</i>)	192
Fig. 3.— <i>COREGONUS OXYRHINCHUS</i> , Lin. Cuv. (<i>The Laverett</i>)	193
Plate 66. Fig. 1.— <i>MYLETES HASSELQUISTII</i> , Cuv. (<i>The Salmon of the Nile</i>)	196
Fig. 2.— <i>CURIMATA GILBERTII</i> , Quoy. and Gaym.	194
Fig. 3.— <i>SAURUS MEDITERRANEUS</i> , Cuv. (<i>The Mediterranean Saurus</i>)	197
Plate 67. Fig. 1.— <i>CLUPEA HARENGUS</i> , Lin. (<i>The Common Herring</i>)	200
Fig. 2.— <i>GNATHOLEBOS MUCRONATUS</i> , Lacep.; or <i>Odontognathus aiguillonné</i>	202

TABLE OF THE PLATES.

XXI

PISCES.

Vol. II. Page

Plate 67. Fig. 3.— <i>PRISTIGASTER AMERICANUS</i> , Cuv. and Val. (<i>The American Pristigaster</i>)	203
Plate 68. Fig. 1.— <i>AMIA CALVA</i> , Lin. (<i>The Amia</i>)	206
Fig. 2.— <i>LEPIOSOTEUS OSSEUS</i> , Lin. (<i>The Lepisosteus</i>)	207
Fig. 3.— <i>POLYPTERUS BICHIR</i> , Geoff. (<i>The Birchir</i>)	208
Plate 69. Fig. 1.— <i>MEGALOPS CYPRINOIDES</i> , Bl. (<i>The Great Megalope</i>)	204
Fig. 2.— <i>CHIROCENTRUS DENTEX</i> , Schn. (<i>The Many-toothed Chirocentrus</i>)	204
Fig. 3.— <i>ERYTHRINUS MALABARICUS</i> , Bl. (<i>The Erythrin of Malabar</i>)	206
Plate 70. Fig. 1.— <i>GADUS MERLANGUS</i> , Lin. (<i>The Whiting</i>)	210
Fig. 2.— <i>LOTA VULGARIS</i> (<i>The Common Ling</i>)	210
Fig. 3.— <i>LEPIDOLEPRUS CAELORHYNCHUS</i> , Risso,	212
Plate 71. Fig. 1.— <i>PLATESSA LIMANDA</i> , Lin. (<i>The Dab</i>)	214
Fig. 2.— <i>RHOMBUS</i> ; <i>Platessa rhombus</i> , Lin. (<i>The Barbue or oval Turbot</i>)	215
Fig. 3.— <i>ACHIRUS MARMORATUS</i> , Lacep. (<i>The Marbled Achirus</i>)	217
Plate 72. Fig. 1.— <i>LEPADOGASTER GOUAN</i> , Lacep.	217
Fig. 2.— <i>CYCLOPTERUS LUMPUS</i> , Lin. (<i>The Lump Sucker</i>)	218
Fig. 3.— <i>ECHENEIS REMORA</i> , Lin. (<i>The Remora</i>)	219
Plate 73. Fig. 1.— <i>MURÆNA ANGUILLA</i> , Lin. (<i>The Common Eel</i>)	221
Fig. 2.— <i>GYMNOTUS ELECTRICUS</i> , Lin. (<i>The Gymnotus, also called the Electrical Eel</i>)	225
Plate 74. Fig. 1.— <i>OPHIDIUM BARBATUM</i> , Bl. (<i>The Common Donzelle</i>)	227
Fig. 2.— <i>AMMODYTES TOBIANUS</i> , Bl. (<i>The Lancet</i>)	227
Fig. 3.— <i>AMMODYTES LANCEA</i> , Penn. (<i>The Equille</i>)	228
Plate 75. Fig. 1.— <i>SYNGNATHUS VIRIDIS</i> , Risso (<i>The Green Sea Eel</i>)	229
Fig. 2.— <i>HIPPOCAMPUS BREVIROSTRIS</i> , Cuv. (<i>The Short-snouted Sea Horse</i>)	229
Fig. 3.— <i>PEGASUS DRACO</i> , Lin.	230
Plate 76. Fig. 1.— <i>TRIODON BURSARIUS</i> , Reinw.	234
Fig. 2.— <i>BALISTES RECTANGULUS</i> , Schn. (<i>The File-Fish</i>)	235
Fig. 3.— <i>OSTRACION TRIQUETER</i> , Bl.	237
Plate 77. Fig. 1.— <i>ACIPENSER STURIO</i> , Lin. (<i>The Common Sturgeon</i>)	239
Fig. 2.— <i>SPATULARIA SPATULA</i> , Maud.	240
Fig. 3.— <i>CALLORHYNCHUS ANTARCTICUS</i> , Lin. (<i>The Antarctic Chimaera</i>).	241
Plate 78. Fig. 1.— <i>SCYLIUM MALAIANUM</i> , Lesson.	243
Fig. 2.— <i>SQUALUS CARCHARIAS</i> , Lin. (<i>The White Shark</i>)	244
Fig. 3.— <i>ZYGÆNA MALLEUS</i> , Val.	247
Fig. 4.— <i>PRISTIS ANTIQUORUM</i> , Lath. (<i>The Common Saw-Fish</i>)	248
Plate 79. Fig. 1.— <i>RHINOBATUS LÆVIS</i> (<i>The Smooth Rhinobatos</i>)	249
Fig. 2.— <i>TORPEDO NARKE</i> , Riss. (<i>The Ocellated Torpedo</i>)	250
Fig. 3.— <i>TRYGON HALGANI</i> , Less.*	251
Fig. 4.— <i>CEPHALOPTERA GIORNA</i> , Lacep.	253


* A newly discovered species of Ray, nearly allied to the *Raja Sephen*, Forsk.


PISCES.	Vol. II. Page
Plate 80. Fig. 1.— <i>PETROMYZON MARINUS</i> , Lin. (<i>The Sea Lamprey</i>)	254
Fig. 2.—Anatomical details of the <i>PETROMYZON FLUVIATILIS</i> , Lin. (<i>The River Lamprey</i>)	254
Fig. 3.— <i>AMMOCÆTES BRANCHIALIS</i> , Lin. (<i>The Common Pride</i>)	256
Fig. 4.—Anatomical details of the <i>AMMOCÆTES RUBER</i> , Lacep. (<i>The Red Pride</i>)	256



P I S C E S , O S T E O L O G Y .


Plate I.—Osteology of the <i>Perch</i> . The skeleton	269
Plate 2.—The front and back of the skull, and the bones of the branchiæ	269
Plate 3.—The bones of the face, the branchiæ, the shoulder, cavity of the stomach, and the vertebræ	269
Plate 4.—Myology of the <i>Perch</i> . Exhibiting the external appearance of the nerves, when viewed in a lateral position	310
Plate 5.—Exposing the whole depth of the lateral nerves	312
Plate 6.—Details of the Myology of the <i>Perch</i> . Views of the neck, brain, and Great sympathetic nerve	325
Plate 7.—Details of the Splanchnology and Angiology of the <i>Perch</i>	352
Plate 8.—Details of the Splanchnology and Angiology of the <i>Perch</i>	352





1. *Hemidactylus marginatus*. . . . 2. *Uropeltis ceylanicus*.


3. *Dactylolethra capensis*. /The Thimble or Cape Frog.



1. *Testudo depressa*. 2. Head of the *Testudo clausa*. 3. *Euys concinna*. (Taken from a young Specimen)
4. *Chelonia virgata*. 5. *Chelys fimbria*. (The fringed Tortoise). 6. *Tryonix gangeticus*. (The Tyre of the Ganges).
7. *Tryonix aegyptiacus*, Grotf. (The soft shelled Tortoise of the Nile).




1. *Crocodilus biporcatus*. (The Double-crested Crocodile). 2. *Alligator sclerops*. (The Spectacle Alligator).
3. *Gavialis tenuirostris*.






1. *Testudo serrata*, Bland. 2. *Testudo undosa*, Linnaeus. 3. *Testudo capreolus*, Bland.
 4. *Testudo caretta*, Graa. 5. *Testudo corrucata*, Linnaeus. 6. *Testudo ferox*,
 Pennant. (The Soft-shelled Tortoise of America.)



1. *Testudo fimbriata*, *Wolff.* 2. *Testudo hermanni*, *Lacépède*. 3. *Testudo graeca*, *Linnaeus*.
4. *Testudo stellata*, *Schäpp.* 5. *Testudo geometrica*, *Lacépède*. 6. *Testudo marginata*, *Zinn*.
7. *Testudo radiata*, *Shaw*.



1 the Ameiva (American Lizard)
 2 the scales of the belly of a Sauvegarde.
 3 Centropix calcaratus (the Spurred Cecropix)




1. 2. 3. Head of the Great Python of Java.


4. 5. 6. Head of a Rattlesnake.



2



3



0000
000

1. *Monita crocodilinus*. Murr. /The Great Dragon of Lacede/ 2. *Tupinambis albicularis*. Baud. /The White-throated Monitor/ 3. *Lacerta magnifica*. Seba. /The Beautiful Monitor/




1. *Lacerta ocellata*, Daud. (The Ocellated Lizard) 2. *Lacerta viridis*, Daud. (The Green Lizard)

3. *Lacerta slirpium*, Daud.



1. *Trapelus agypius*, young specimen. 2. *Trapelus agypius*, Geoff. St. Hil. n. s.


3. *Lacerta calotes*, Lin.



1.b




1.a



2

1. *Monitor gemmatus* in the collection of the French Museum.


2. *Dracæna bicarinata*.




1. *Lacerta agilis*. (The Common European Lizard) 2. *Algyra barbarica*, Cuv. *Lacerta Algyra*, Lin.
3. *Tachydromus ocellatus*. (The Ocellated Swift Lizard)



J. S. J. S.




J. S.



2





3



J. S. J. S.

1. *Cordylus microlepidotus.* 2. Head of the *Stellio vulgaris.* (The Common Stellion.) 3. *Dipsosaurus (Dipsoas) vulgaris.* (Short tailed Dipsosaurus of India.) 4. *Uromastyx reticulatus.* (The reticulated Stellion of Bengal.)





Lacerta vittata, Gmel. / The White Banded Gecko / 2. Stellio fimbriatus, Sch. / The Fringed Gecko / 3. Stellio Gecko, Sch. / The White spotted Gecko / 4. Gecko levris D. / The Smooth Gecko /




1



2



3






Anolis velifer / the great crested. *Anolis*.
Echimotes tuberculata / the tuberculated *Echimotes*.
Oplurus torquatus / the Black collared Grey *Oplurus*.

London. G. Henderson 2 Old Bailey.




1 *Anolis velifer* / The Great Crested Anolis; an other view see Pl. L4 / 2 *Anolis equestris* / The Equestrian Anolis
 3 *Gecko marmoratus* / The mottled Gecko / 4 *Gecko ocellatus* / The Ocellated Gecko /
 5 *Gecko cepedianus* / The Cepedian Gecko /




1 *Physignathus cocincinus* (*The Physignath*) 2 *Sitana ponticeriana* (*The Sitana*)
Iguana cychlura (*The Carolina or Arm-tailed Iguana*)


1




1.a




2.a




2




3.a




3




1. *Agama Jacksoniensis*, /The Bachelor/ in the collection of the French Museum. 2. *Leiolepis guttata*, /The striped leiolepis/ 3. *Calotes gutturosa*, /The crested Agama/.





1. *Brachylophus fasciatus*, /The Banded Agua/ 2. *Istiurus cochinensis*, Valencienne, /The
Istiurus or Lophura of Cochin China/ in the collection of the French Museum.


I

 $\frac{1}{2}$ n.s.


1. a




1. b





2. a




2

 $\frac{1}{3}$ n.s.1. *Ophryessa superciliata*.2. *Lyriocephalus marginatus* Mer.

*Anomalous Serpents.*1.2.3. Head of a *Cœcilia*. . . 4.5.6 Head of an *Amphisbana*.7.8.9. Head of an *Ophisaurus*.




1. *Draco fimbriatus*. / *The Dragon* / 2. *Sitana ponticeriana*. / *The Sitan* /



1. *Iguana undicollis*. / *The Naked Iguana* / 2. *Basilicus* / *Lacerta basilicus*. L. / *The Basilisk* /

3. *Polychrus marmoratus*. L. / *The Marbled Lizard* /




Hemidactylus tuberculatus Daud. (The spotted Hemidactylus)






1. *Phyllurus platurus*. White. 2. Foot of the *Platydactylus inunguis*. 3. Foot of the *Hemidactylus trihedrus*. 4. Foot of the *Thecadactylus*. 5. Foot of the *Ptyodactylus lobatus*. 6. Foot of the *Sphaerodactylus*. 7. Foot of the *Stenodactylus guttatus*.




1. *Bipes lepidopodus*, Lacep. / The two footed Lepidopodus / 2. *Chilis tetradactyla*, Lacep.
3. *Chirotes canaliculatus*, Lacep.






1. *Pseudopus Durvillii*. (The false foot of Durville) / 2. Head of the *Anguis fragilis*. Lin. (The common Blind worm of Europe) / 3. *Acontias meleagris*. Lin. (The darting serpent of the Cape of Good Hope)




1. *Amphisbaena fuliginosa*, Lin. /The Double Walker/
 2. *Typhlops cinereus*, Schn. /The Blind Serpent of Schneider/







1 Vipera berus, Lin. /The common Viper/ 2 Crotalus horridus, Lin. /The Diamond Rattle-snake/



1. *Trimeresurus tortor*. Less. /The winding Trimeresurus/


2. *Acanthophis cerastinus*. Band. /The venomous Acanthophis/

1. *Tortrix maculata*. L. / The Variegated Tortrix of America / 2. *Boa canina*. L. / The Canine Boa /3. Tail of the *Uropeltis ceylanicus*. Cav. / The Trepeltis of Ceylon /






1



2




3. a

3. d


3. c

3. b


1. *Chamaeleo jamaicensis*. /The common Chameleon/ 2. *Scincus celestinus*. Valenc. /The blue Scinc/ in the collection of the French Museum. 3. *Seps striata*. /The striped Seps/



Eryx bengalensis, Merr. / The Hairy Serpent of Bengal / 2. Head of the *Scytale ceronata*, Merr.
The Crowned Scytale or False Boa / 3. *Erpeton tentaculatus*, Lapebie / The Tentaculated Serpent



1. *Python Schneiderii*, Merrem. (The Great Python of Schneider) / 2. *Dipsas cyanosticta*, (The fatal thirst causing Serpent of India) / 3. Head of the *Xenopeltis concolor*, Reinwardt. (The many coloured Xenopeltis)

1. *Dendrophis albolabris*. (The Serpent of the Woods.)2. *Drymias nasuta*, Russel. (The Hog-nosed Drymias or Coluber.)



1. *Bungarus annularis*, Daud. 2. *Anguis platurus*, Lin.



1. *Hydrophis eloris*, Russel. 2. *Cœcilia lombricoides*, Daud.

3. *Cœcilia alliventris*, Daud.

London: G. Henderson, 2 Old Bailey.





1. *Hydrophis fasciatus. Russel. (The Bengal Water Serpent.)*

2. *Caecilia bivittata. (The Blind Serpent
or the Caecilia of Ceylon.)*




1. *Rana palmipes*, Leconte. 2. *Ceratophrys granosa*, (The American Horned Frog).

3. *Hyla bicolor*, Daud. (The South American two-coloured Tree Frog).







1. *Bufo vulgaris*. /The Common Toad/ 2. *Oxyrhyynchus bicolor*. Val. /The Angular Muzzle Toad/ in the collection of the French Museum. 3. *Engystoma marmoratum*. /The Marbled Toad/




1 *Salamandra atra*, Laurenti. / The Black Salamander / 2. *Triton vittatus*, Valenci. / The Triton / in the collection of the French. 3. *Dicamptodon giganteus*, Barton. / The great Salamander of North America /






1. *Amphiuma means*, Gard. & Hirlimann of the French Museum. / The two-toed Amphiuma / 2. *Amphiuma trydactylum*. / The three-toed Amphiuma / 3. *Axolotl pisciformis*, Shaw. / The Fish serpent of Mexico / 4. *Monobranchus lateralis*, Harl.








1. *Protens anguinus*. Laar. / The Blind Proteus / 2. *Siren striata*. Lecomte. / The Streaked Syren /
3. *Siren intermedia*. Lecomte / The intermediate Syren of Leconite /






1. *Etelis carbunculus*. /The Etelis/ 2. *Diplopteron bifasciatum*. /The bifasciate Diplopteron/ 3. Head of the *Huro nigricans*. /The Black Basse/ 4. Head of the *Niphon spinosus*. /The Spiny Niphon/ 5. Head of the *Enoplosus armatus*. /The Enoplosus or armed Perch of Japan/




1. *Perca acenta* Cuv et Val. 2. *P. Latr. nobilis* Cuv et Val. 3. *Centropomus undecimalis* n.
Eleven Spined Centropomus




1. *Apogon trimaculatus n./* The Mediterranean Apogon. 2. *Cheilodipterus arabicus n./* Arabian Cheilodipterus. 3. *Ambassis commersonii n./* Commerson's Ambassis.



1. *P. Lucioperc a sandra. n.* *The Common Pike Perch.* 2. *P. Lucioperc a american a.* *The American Pike.*
3. *Huso nigricans. n.* *The Black Bass of the Huron.*




1. *Luciopercus Volgensis* Pallas / The Russian Perch. 2. *Niphon spinosus* / The Spined Niphon /
3. *Enoplosus armatus*, n. / The Armed Enoplosus. /




1. *Serranus anthias*. / The spined Serranus / 2. *Serranus cabrilla*. Nob. / The common Serranus /

3. *Serranus gigas*. Nob. / The Meron of the Mediterranean /


London, G. Henderson. 2. Old Bailey.



1. *Serranus phactus*. Nob. (Long-tailed Serranus.) / 2. *Serranus fasciatus* Nob. (The Banded Serranus.) / 3. *Serranus oculatus* Nob. (The Large-eyed Serranus.)



2






3



1 *Mesopriion unicolor* (One-spotted Mesopriion) 2 *Mesopriion chrysoura* Rob. Golden-tailed Mesopriion

3 *Diacope rivulata* Rob. The Banded Diacope




1. P. *Labax mucronatus*. n. 2. P. *Labax tigris*. n. 3. *Seretulus scriba*. n.

1



2




3



1 *Sternopyx diaphana* 2 *Sternopyx Offerstii*

3 *Gymnarchus senegalensis*




1. *Plectropoma leopardinus*. n. / Leopard Spotted Plectropoma / 2. *Plectropoma puella*. n. / Beautiful Plectropoma / 3. *Serranus altivelis* / Large Finned Serranus. /




1. *Pentaceros capensis*, Rob. 2. *Geistes salmonides*, Rob. (The Salmon-formed trawler)
3. *Ctenopristes nigricans*, Rob. (American Black Bass)






1. *Percus serrato-granulata*. (The Granulated Perch) / 2. *Grammistes orientalis*. (The Oriental Grammist) / 3. *Aspro vulgaris*. (The Common Aspro) / 4. Head of the *Labeo vulgaris*. (The Rasse Perch) / 5. Head of the *Lates nebulosus*. (The American Pike)



1. *Heliaes insolatus* n. 2. *Pomacentrus fasciatus* n. 3. *Glyphisodon calostinus* n.




1/2 n.s.



3 a.s.

1. *Centropristes aurorubens*. The Golden Red Perch / 2. *Gristes marginatus*. The Growler of New South Wales / 3. Head of the *Acerina vulpis* (The Common Grenille) / 4. Head of the *Polyprion cornutus*. The Mediterranean Cernier / 5. Head of the *Rypticus sanguineus* (The Soap Wart) / 6. Fin of the *Cirrhitus fasciatus*.




1. *Dules marginatus*, Rob. ('The Bordered Dule') / 2. *Datnius argenteus*.


3. *Therapon*. *Theraps*.



1. *Apogon rec nullorum.* / *The Red Surnullet.* / 2. *Pomatomus telescopium.* / *The Telescope Pomatomus.* /
3. *Lucioperca sandra.* / *The Common Sandre.* / 4. *Head of the Cheilodipterus arabicus.* / *Tabah of the
Arabs.* / 5. *Head of the Ambassis commersonii.* / *The Dugol.* /




1 *Serranus hexagonatus*, Cuv. s.g. *Meron*. / *The Hexagon Sea Perch*. / 2 *Mesoprion aya*, Cuv. / *The Garra or Red Mesoprion*. / 3. Head of the *Plectropomus puello*, Cuv. / *The White Lady of Martinique*. / 4. Head of the *Serranus anthius*, s.g. *Barbier*, Cuv. / *The Mediterranean Barber*. / 5. Head of the *Diacope rivulata*, Cuv. / *The Coromandel Oruti*.



Centrarchus spariooides. 2. *Dalea rostrata*. (*The Dalea*) / 3. Head of the *Pomotis vulgaris* (*American Pond Perch*) 4. Head of the *Priacanthus japonicus*. (*The Priacanthus of Japan*)

1




2



3



1. *Pelor filamentosum*, n. 2. *Platycephalus asper*. 3. *Dactylopterus orientalis*, nab.




1. *Therapon puta.* (*The Therapon*) 2. *Pelates quadrilineatus.* (*The four-lined Pelates*)
3. *Trichodon stellifer.* (*The Trichodon*) 4. *Head of the Helotes sexlineatus.* (*The six-lined Helotes*)



1 *Holotes sex lineatus*, n. 2 *Myripristes japonicus*, n. 3 *Holocentrum hastatum*, n.

1




4 n.s.




1

5




2






4 n.s.

3






4 n.s.

1. *Trachinus araneus* Rien. 2. *Percis ranceillata*. 3. *Uranoscopns guttatus*.4. Head of the *Pinguipes Brasilianus*. 5. Head of the *Pecophis Brasilianus*.




1. *Sillago erythraea.* 2. *Myripristes seychellensis.*



3. *Beryx decadactylus.*




Percis semilasciata, n. The half banded Percis/2. Percophis brasiliensis, n. The Brasilian Percophis/
3. Uranoscopus unarmatus, n. The unarmed Star Gazer.


Linden G. Henderson, 2. ill. Bailey.






1. *Beryx lineatus*. n. 2. *Trachinus radiatus*. n. 3. *Pinguipes brasiliensis*. n.



1. *Polynemus macrourus*. 2. *Sphyraena vulgaris*. 3. *Paralepis ceregooides* Rissso.4. *Upeneus latetus*. 5. Head of the *Mullus barbatus*.




2






3




1. *Polydactylus quadrifilis* / The four-fingered Paradise Fish/ 2. *Paralepis sphaeroides* Rissso.
3. *Sphyraena* Barracuda Cuv. et Val.

Head of a Cod — *Gadus morrhua* Lin.




1 *Sillago dominica* n. 2 *Mullus barbatus*, Linn. 3 *Upeneus Thaumaturus*.




1. *Agriopus verrucosus*. 2. *Pelor japonicum*, n. 3. *Synanceia venusta*.






3






2




1. *Scorpaena grandicornis*, n. 2. *Bembras japonicus*, n. 3. *Hemilepidotus*. *Tiburon*.



1. *Trigla pacifoptera*. Val. & Cuv. 2. *Cephalacanthus spinarella*. Cuv. 3. *Cottus porosus*. Cuv.4. *Cottus octodecim spinosus*. Cuv.





1. *Priacanthus tribulus nob.*, 2. *Trigla lucerna Brünn.*, 3. *Peristedion cataphractum Lacep.*




1. *Aspidophorus segaliensis*, Gmelin. 2. Head of the *Aspidophorus quadricornis*, Gmelin. 3. *Hemitripterus americanus*, Gmelin. 4. *Hemilepidotus tilesii*, Gmelin. 5. Head of the *Platycephalus osper*, Gmelin.





3




1. *Trigla papilio*, nob. 'The Butterfly Gurnard' / 2. *Cottus bubalis* Faufr. 'The Bull Head' / 3. *Optichthys Lutgsdorffii*.




1. *Sciæna pama* n. 2. *Otolithus ruber* n. 3. *Otolithus terce* n.






1. *Platycephalus grandispinis*, Guv. 1. a. Thickness of the body towards the Head. 1. b. Thickness of the middle. 2. *Scorpaena neogallica*, Guv. 3. *Tarmianotus triacanthus*, Guv.






1. Chelodactylus Antonii. 2. Amphiprion chrysogaster. 3. Heliasys firmatus. 4. Head of the tulopsides tyrogenis. 5. Head of the Premnas semicinctus. 6. Head of the Dacryllus marginatus. 7. Head of the Pomacentrus fasciatus. 8. Head of the Glyptodon celestinus.



1. *Sebastes norvegicus*, n. 2. *Blepsias lilloensis*, n. 3. *Pterois volitans*, n.



1. *Otolithus guatucupa* n. 2. *Aucylodon parvipinnis* n. 3. *Corvina semiluctuosa* n.




1. *Sebastes variabilis*, Gv. 2. *Blepsias trilebus*, inv.

3. *Apistes marmoratus*, Gv.






1. *Pagellus centrodontus*. 2. *Buops salpa*. 3. Head & teeth of the *Sargus Rondeletii*. (The *Sargus* of Rondelet) 4. Head & Dentition of the *Chrysophrys aurata*. (The Golden-Eye bream) 5. Head & Dentition of the *Pagellus erythrinus*. (The Paget)




1. *Eques punctatus* Bl. 2. *Boridio grossidens* n. 3. *Eleginus macrourus* n.


1




2



3




1. *Haemulon quadrilineatum*, n. 2. *Haemulon heterodon*, n. 3. *Pristipoma bilineatum*, n.




2








3



4




1. *Premnas semicinctus*. 2 & 3. The Brain of the *Sciana Pama*. 4. *Dascyllus marginatus* Sh.



$\frac{3}{2}$ n. s.


1. *Agriopuss peruvianus*. Gv. 2. *Pelox maculatum*. Gv.

3. *Synanceya elongata*. Gv.




1. *Lobotes somnolentus*. 2. *Diagramma orientale*, n. 3. *Diagramma caristrons*, n.


1






2



3




1. *Glypthisodon koruschin*. 2. *Amphiprion latidorsus*. n. 3 *Amphiprion tuniceatus*. n.




1. *Gasterosteus trachurus*, Cuv. et Val. 2. *Gasterosteus Gymnurus*, Cuv. et Val. 3. *Gasterosteus noreboracensis*, Cuv. et Val. 4. *Gasterosteus leirus*, Cuv. et Val.




2



3



3. a

1. *Monocentris japonica*, Gv. 2. *Gasterosteus brachycentrus*, Gv.3. *Oreosoma coniferum*, Gv.









Lutjanus breviceps, Nob. (The Larine) / 2. *Leptipterus Francisci*, Nob. (The Leptipter of the Island of St. Francis) / 3. *Corypha ocellata*, Nob. (The Spotted Corvina) and
Anatomical details of the Brain of the Aubas sciaua. 5. Anatomical details of the Brain of the Aplochoplites London. 6. Illustration, see 2. W.H. Bailey.



1. *Pogonias fasciatus*, n. / The Banded Pogonias. / 2. *Umbrina cornicles*, n. 3. *Micropogon liratus*, n.




1. d




1 *Muraena venusta*. Nob. 2 *Boops salpa*. Nob. 3 *Aphareus carolinensis*. Nob. 4 A fragment of the intermaxillary jaw of a *Daurades* with the molars detached. 5 The superior intermaxillary jaw of a *Sargus*. a. The incisors. b. The root of an incisor tooth. c. The molars.


1



2






3




1. *Latilus doliatus*, n. 2. *Cheilodactylus zonatus*, n. 3. *Macquaria australis*, sp. n.










1. *Siganus triacanthus*. 2. *Chaetodon strigatus* Longsd. 3. *Chaetodon reticulatus*.




1. *Centrarchus sparoides*. 2. *Rypticus arenatus*. 3. *Cirrhites fasciatus*.






1 *Taenianotus triacanthus* Lacep. 2 *Apistus trachinooides*. 3 *Apistus monodactylus*, n.

 $\frac{7}{12}$ n. s. $\frac{7}{12}$ n. s. $\frac{7}{15}$ n. s.

1. *Scirna aquila*. 2. *Corvina Eureca*. 3. *Johnius senegalensis*. 4. Head of the *Otolithus ruber*.
5. Head of the *Ancylodon parvipinnis*. 6. Head of the *Umbrina coroides*. 7. Head of the *Pogonias fasciatus*.




1. *Nemipterus Peroni* / The Shepherd / 2. *Caranx hoopo*, / The False Mackerel or carangid
3. *Guttus egyptiacus*, / The Egyptian Guttus / Cuvier's Ichthyology, Vol. 2, page 241





1. *Otolithus toroe*, Nob. 2. *Otolithus guatucupi*, Nob. (*The Striped Otolithe*)


3. *Corvina lobata*, Nob.




1. *Cheilodactylus carponensis*. n. 2. *Scolopsides lycogenis*. n. 3 *Diagramma gatherina*. n.



1. *Chaetodon strigatus*. Sol. 2. *Chaetodon lunula*. n. 3. *Chaetodon ephippium*.




1. *Eques balteatus*. 2. *Lobotes erate*. 3. Head of the *Hermuthion heterodon*. 4. Head of the *Pristipoma bilineatum*. 5. Head of the *Diagramma orientale*.




1




2



3



1. *Hemistomus humeralis*, Rob. or *Johnius humeralis*, var. / 2. *Nebrius micros*, Rob. / The small eyed Nebrius. / 3. *Corvina dentex*, Rob. / The many tooth Corvina. /



1. *Sargus annularis*, Nob. 2. *Sargus rhomboides*, Nob.
3. *Charax punctazzo*, Nob. (*The Common Puntazzo*)




1




2


1. *Zanclus cornutus*, 2. *Heniochus meyerrossi*, n.



2




3



1. *Chrysophrys gibbiceps*, Nob. 2. *Chrysophrys crassirostris*, Nob.

3. *Chrysophrys aurata*, Nob.


Loudon, 1829; Henderson, 2. Old Bailey.



1 *Scatophagus ornatus*, n. 2. *Taurichthys varius*, n. 3. *Holacanthus trimaculatus*, n.




1




3. b



3




4



3. a





2




1. *Simpis audeo*. (The blue-spotted Bream.) 2. *Cesio tile*, var. 3. *Maena venturiana*, var. (Vomer Mudelci)

4. *Gerres Phainieri*, var.


1. *Pagellus lithognathus*, Nob. 2. *Pagellus erythrinus*, Nob. (The Papel)3. *Pagellus calanus*, Nob.



1




2



3

1. *Lethrinus genivittatus*. Nob. 2. *Cantharis vulgaris*. Nob.

3. *Boops vulgaris*. Nob.



2/3 n. s.




2/3 n. s.



3


1. *Chaetodon setifer*. (inv.) 2. *Ephippus orbioides*. (inv. /The Wrasse/)

3. Head of the *Ephippus leagirostris*.



1. *Dentex Peronii*, Nob. (*Peron's Dentex*) 2. *Dentex filamentosus*, Nob.
3. *Dentex vulgaris*, Nob. (*The Common Dentex*)









1. *Chelmon longirostris*, n. The Long-nosed Chelmon. 2. *Aspidophorus monopterygius*, bl.
The Aspidophorus with a single Dorsal. 3. *Apriou virescens*, n.




1. *Spirobranchus capensis*. 2. *Trichopus trichopterus*. 3. Anatomy of the Brain of the Ephippion (*Ephippus gigas*). 4. Shoulder bone. 5 & 6. Interspinal & Anal bones of the Ephippion (*Ephippus gigas*).



1 *Anabas scandens*, n. 2. *Psettus Sebire*, n. 3. The Labyrinthal Organ of the *Ophichthoides Striatus*.
4 Anatomy of the Brain of the *Osfhromenus Gourumi*.




1. *Holocanthus ciliaris*, L. var.



2. *Pomacanthus arcuatus*, L.

1. *Holocanthus ciliaris*, L. var. 2. *Pomacanthus arcuatus*, L.






3/10 n.s.



5




7/10 n.s.

2. ♂.



2



1/10 n.s.

1 Scomber scombrus / the common Mackerel,

2 Thynnus vulgaris / the Tunny;

3 Alosa lele / the Axis;

4. the mouth with the teeth of the Axis.

5. the mouth of Pclanys cardu.

London. G. W. Anderson 2. Old Bailey




1

2. a



2. a




2

2. b




2. d




3

3. b


1. *Pentapodus aurolineatus*. Nob. (The Golden striped Pentapode.) 2. *Pentapodus porosus*. Nob.3. *Lethrinus esculentinus*. Nob.



2



3





1. *Ophicephalus limbatus.* 2. *Ophicephalus striatus.* Bl?



3. *Ophicephalus grandinosus.* Sob.





1. *Colisa vulgaris*. 2. *Macropodus venustus*. n. 3. *Platax guttulatus*. n. 4. *Platax punctulatus*. n.








1. *Platus Gaimardi*, *Thw.* 2. *Psettus rhombus*, *Thw.*



5




1. *Ephippus coreensis*, Nob. 2. *Drepane punctata*, Nob.

3. Head of the Spinobranchie. 4. Head of the Maetropode. 5. Head of the Common Colisa.




1. *Thunnus pelamys*. Nob. / The Tropical Bonito or ray-bellied Tunny / 2. *Thunnus sarda*. Nob. / The common Pelamis / 2. a. Dentition of the *Pelamis sarda*. 3. *Thunnus thynnus*. Nob. / The Thonine / 4. Head of the *Xiphias gladius* L. / The Sword Fish a young subject / 5. The same as Fig. 4. but taken from a full grown Fish.

London, 6. Henderson, 2. Old Bailey.




1. *Pimelepterus Boscii*, Lac. / 2. *Polyacanthus Hasseltii*, n. 3. *Brama Rua* Bl. S. J.

1




3. c



3. a




3



2. a








2






1. *Pimelepterus altipinnis*. 2. *Diptorodon capensis*.

3. Head of the *Pimelepterus Bocci* Lac.






1. *Thysites atun*. Nob. 2. *Cybium tritor*. Nob.

3. *Thysites lepidopoides*.




1. *Gempylus prometheus*. Job. 2. *Lepidorhynchus argyreus*. Job.

3. *Gempylus coluber*. Job.



1. *Helostoma Truncatum* Ritter. 2. *Pempheris taeniensis* n. 3. *Toxotes jaculatrix* n.






1. *Brama Raii. Bl.* 2. *Pompheris mexicana.*

3. *Toxotes jaculator. Sh.*



London: G. Henderson, 2, Old Bailey.






1. *Gerres plumieri*, n. 2. *Caesio erythrogaster*, n. 3. *Suaoris chrysotis*, n.

London: G. Henderson. 2. Old Bailey.




2




3



1. *Flacate atlantica*. n. 2. *Lichia glayces*. n. 3. *Lichia valigo*. n.




1. a




2.



2. a



3.



3. 3 n.s.


1. *Cybium lineolatum*. / The long Tassand. / 2. *Thryssites chilensis*. / The White Thryssites. /
3. *Gempylas prometheus*. / The Gempyle. /




1. *Olistus Malabaricus*. Cuv et Val. 2. *Seyris indica*. Cuv et Val.



1




2




3

1. *Apolectus stromateus*, n. 2. *Chorinomus Sancti Petri*, n. 3. *Trachinotus pampanus*, n.







1. *Tetrapterus Indicus nob.* 2. *Skeleton of the Tetrapterus, Belone Barr.*

London G. Henderson & old Bailey.




1. *Xiphias gladius* A. 2. *Scomber* Fish but of younger age. 3. *Tetrapturus belone*, Raf.
4. *Histiophorus pulchellus*.






Mastacembelus indicus. 2. *Mastacembelus maculatus.* 3. *Seriola Rivolii.* 4. Head of the Elacate *antica.* 5. Head of the *Phynchobdella ural.* 6. Head of the *Mastacembelus armatus.* 7. Head of the *Notacanthus nasus.* 8, 11. 9, 10. 12. 13. n.s.



1. *Bovichtus diacanthus*, n. 2. *Aphritis Urvillii*, n. 3. *Aplodactylus punctatus*, n.


1




2




3




1. *Notacanthus nasus* Bl. 2. *Mastacembelus armatusn.* 3. *Rhynchobdella oralisn.*



$\frac{7}{9}$ n.s.




$\frac{7}{9}$ n.s.



$\frac{7}{9}$ n.s.

1. *Zenus faber*. L. / The common Dory / 2. *Lampris guttatus*. Retz. / The Moon Fish of Lacedepe /


3. *Stromateus fimbriata*. L. / The Mediterranean Pompet /



1. *Argyreyosus vomer*, Lacep. 2. *Caranx trachurus*, Lacep. 3. *Caranx chrysophrys*, n.




1. *Blepharis salomon*. 2. *Callichthys major*.




$\frac{7}{2}$ n.s

Gymnusetus falcatus.


London. C. Henderson 2 Old Bailey.



1 n.s.



2 n.s.




3 n.s.


1. *Siganus dolianus*. 2. *Acanthurus Devisimus*.

3. *Nasens longicornis*.

1


 $\frac{3}{2}$ n.s.


2



n.s.

3

 $\frac{3}{4}$ n.s.1. *Anabas testudineus*, Bl. 2. *Polyacanthus colisa*, Ham.3. *Ophicephalus serpentinus*.



2. a




2. n. s.


2. c



2. b



?




2. n. s.

2. b



?




2. n. s.


1. *Mugil cephalus*. 2. *Tetragonurus Cuvieri*, Riso.3. *Atherina presbyter*.

London & Henderson, 2. Old Bailey.


3










 $\frac{1}{4}$ n. s.

2

 $\frac{1}{4}$ n. s.




1

 $\frac{1}{3}$ n. s.1. *Bleennius palmeicornis*. 2. *Clinus pectinifer*.3. *Zoarces blennoides*.




3.b

1. *Gobius capito*, *an* /The great Goby or Sea Gudgeon/ 2. *Gobiooides Broassenetii* /The great Gobiooides of Broassenet/ 3. *Periophthalmodon Freyinetii*




1. *Eleotris dormitatrice*, Linn. / The Sleeper / 2. *Callionymus fasciatus*, / The fasciated Callionymus /
3. *Chirurus decagrammus*, Ball. / The Ten Striped Chirus /






1. *Lophius piscatorius.* 2. *Antennarius nesogallicus.*

3. *Batrachus surinamensis.*




1




2 n.s.

2



2 n.s.

3





3 n.s.

1. *Labrus vittatus*, Quoy and Gaim. 2. *Julis trimaculata*, Quoy and Gaim.

3. *Crenilabrus chabrolii*, Less.






1. *Chromis nilotica*, Cav. 2. *Malacanthus furcatus*, Cav.3. *Scarus creticus*, Abbrev. Cav.

1. *Epibulus insidiator*. Pall. Cuv. 2. *Gomphosus viridis*. Cuv.3. *Xirichthys cyanirostris*. Cuv.




1. *Fistularia tabacaria* (The Tobacco pipe Fish also called Flute Mouth) / 2. *Centriscus s. cliviger*.
L. (The Sea Snipe) / 3. *Ampelisca dentatus* L. (The Shielded Sea Snipe)

1. *Cyprinus amarae*. (The Amurian Carp) 2. *Gobio vulgaris*. (The common Sea Gudgeon)3. *Labeo niloticus* Geoffr. (The Mullet of the Nile)



1. a

2. n. s.




2.

n. s.

3. b

3. a




H. S.

1. *Lencis sens. illeus*, Block. (The Albe commonly called the White Fish) / 2. *Cobitis barbatula*, Lin. (The common Bande Loach) / 3. *Poecilia unipunctata*, Val. (The One dotted Poecilia)




1. *Lebias fasciata*, Val. / The fasciated Lebias / 2. *Fundulus brasiliensis*, Val. / The Fundule of Brazil / 3. *Cyprinodon flavulus*, Val. 4. Anatomical details of the *Fundulus fasciatus*, Val. / The Fasciated Fundule / 5. Dentition of the *Lebias rhomboidalis*, Val. / The Rhomboid Lebias /

1

 $\frac{1}{3}$ n.s.

1. a


2

 $\frac{1}{2}$ n.s.


2. a

3. a

3

 $\frac{1}{2}$ n.s.

1. *Esox lucius* L. *The Common Pike* 2. *Microstoma mediterranea*. (*The Mediterranean Micrestema*)
3. *Stomias barbatus* Risso. (*The Barbed Stemias*)




I


 $\frac{1}{2}$ n.s.

O

3



 $\frac{1}{2}$ n.s.

2


 $\frac{2}{3}$ n.s.

O

1. *Salpa fuscus*.2. *Scomberesox saurus*. Bloch.3. *Exocoetus acutus*. The sharp snouted Flying Fish /

 $\frac{3}{4}$ n. s. $\frac{1}{2}$ n. s. $\frac{1}{3}$ n. s.


1. *Morayneus oxyrhinchus*, Geoff. Cuv. /The True Morayneus oxyrhinchus of the Ancient Egyptians/
2. *Schilbe mystus*, Geoff. /The Silure of the Nile/ 3. *Synodontis scheilan*, Geoff. /The That of Egypt/




1

1/3 n.s.

2.b




2.a




2

1/3 n.s.

3.b




3.a



3


1. Bayard fitile. Cuv & Val. 2. Doras dorsalis. Cuv & Val.

3. Heterobranchus punctatus. Cuv & Val.






1. *Percus bayad*, Geoff. & Cuv. 2. Mouth of the *Pimelodes Blochii*, Cuv. & Val.

3. *Plotosus albus* Linn. 4. *Callichthys longifilis*, Cuv. & Val.




1. *Aspredo octocirrus*. Cuv. 2. *Hypostomus plecostomus*. Cuv.


3. *Loricaria Cataphracta*. Lin. Cuv.



1. *Salmo fario*, L. Cuv. (The Common Trout) 2. *Mallotus greenlandicus*, Bl. Cuv. (The Capelan)
3. *Coregonus oxyrinchus*, L. Cuv. (The Laverett)




1. *Myletes Hasselquistii*, Gav. (Sometimes called the Salmon of the Isle) / 2. *Cirimata* Gilbert, Gray & Gym.
3. *Saurus mediterraneus*, Gav. (The Mediterranean Saurus)



1

3. H. S.



2

3. a




2

3. H. S.


1. *Clupea harengus*, Lin. ('The common Herring') 2. *Gnathopholus microstomus*, Lc.
3. *Pristigaster divergens*, Gv. ('The divergent Pristigaster')

Lendenius Henderson 2 Old Bailey.







1. *Anutia calva*, Lin. / The Anutie / 2. *Lepisosteus osseus*, L. / The Lepisosteus /

3. *Polypterus birchir*, Geoff. / The Birchir /








1. *Megaleporinus eximius* Bl. / The great Megaleporine / 2. *Chirocentrus dentex*, Sch. / The
sea trout chirocentrus / 3. *Erythrinus malabaricus* Bl. / The Erythrin of Matabar /



1. *Gadus morhua*, L. / The Whiting. / 2. *Lota vulgaris*, / The common Ling)


3. *Lepidolepis colorynchus*, Risso.






1. *Platessa limanda*, L. / The Dab / 2. *Rhambus* / *Platessa rhombus*, L. / The Barbeque or oval Turbot /
3. *Achirus marmoratus*, Lac. / The marbled Achirus /

1



n.s.

2.a




2




2 n.s.

2.a


3.b



3.a



3





3 n.s.

1. *Lepadogaster gowani*. Lac. 2. *Cyclopterus lumpus*. L. (The Lump Sucker)
3. *Echeneis rentora*. L. (The Remora)






1. *Muraena anguilla*. Lin. (The Common Eel)

2. *Gymnotus electricus*. L. (The Gymnotus, also called the Electrical Eel)




1. *Ophidion barbatum*, Bl. (The Common Dentelle) 2. *Ammodytes tobianus*, Bl. (The Lancet)
3. *Ammodytes lancea penn*. (The Equille)



1. *Syngnathus varidus*, Risso. (The green Sea eel). 2. *Hippocampus brevirostris*, Linn. (The short-snouted Sea Horse). 3. *Pegasus draco*, L.





1. *Triodon hirsutus*, Reim. 2. *Balistes rectangularis*, Schm.


3. *Ostracion triqueter*, Ill.

1



420. gr. n.

1 b




1 a

2



410. gr. n.


3 a



3 b



3




430. gr. n.

1. *Acipenser sturio* L. *The Common Sturgeon*, p. 379. 2. *Spatularia spatula* Muraq. 3. *Callorhynchus Antarecticus* I. *Car. t. 2*. p. 382. *The Antarctic Chimaera*.


1




2



3



4





1. *Squalus acanthias* Lesson. 2. *Squalus carcharias*, Lin. (The White Shark) 3. *Zygoprion*
meridionalis Val. 4. *Carcharias taurus*, Lath (The Common Saw Fish)




1. *Rhinobatos levis*. /The smooth Rhinobatid/ 2. *Torpedo narke* Riss. /The ocellated Torpedo/
3. *Trygon Halgani* Tess. 4. *Cephalopterix Giorna* Lac.


London. G. Henderson. 2. Old Bailey.







1. *Petromyzon marinus*, L. / The Sea Lamprey / 2. Anatomical details of the *Petromyzon flavivittatus*, L. / The River Lamprey / 3. *Ammocetes branchialis*, L. / The tommon Pride / 4. Anatomical details of the *Ammocetes ruber*, Lac. / The Red Pride /




OSTEOLOGY OF THE PERCH, PL. I. *The Skeleton.*


London: G. Henderson, 214 Fleet Street

OSTEOLOGY OF THE PERCH. PL. II. *The front & back of the skull & the bones of the Branchia.*







Animal Kingdom




OSTEOLOGY OF THE PERCH. PL. III. The bones of the Face, the Branchia, the Shoulder, cavity of the Stomach, and the Vertebrae.






THE MYOLOGY OF THE PERCH. PL. I. Exhibiting the external appearance of the nerves when viewed in a lateral position.



THE MYOLOGY OF THE PERCH, PL. II Exposing the whole depth of the lateral nerves.

London, J. Henderson, 2 Old Bailey






DETAILS OF THE MYOLOGY OF THE PERCH. Views of the Neck Brain, & great Sympathetic Nerve.


London: W. H. Allen, 2, Old Bailey.





Details of the Splanchnology and Angiology of the Perch.





Details of the Splanchnology and Angiography of the Perch.

London: G. Henderson, 2 Old Bailey

350



