

**Mathematics for natural sciences I****Exercise sheet 25****Warm-up-exercises**

EXERCISE 25.1. Compute the definite integral

$$\int_0^{\sqrt{\pi}} x \sin x^2 dx .$$

In the following exercises, which involve the determination of antiderivative functions, consider an appropriate domain of definition.

EXERCISE 25.2. Determine an antiderivative of the function

$$\tan x .$$

EXERCISE 25.3. Determine an antiderivative of the function

$$x^n \cdot \ln x .$$

EXERCISE 25.4. Determine an antiderivative of the function

$$e^{\sqrt{x}} .$$

EXERCISE 25.5. Determine an antiderivative of the function

$$\frac{x^3}{\sqrt[5]{x^4 + 2}} .$$

EXERCISE 25.6. Determine an antiderivative of the function

$$\frac{\sin^2 x}{\cos^2 x} .$$

EXERCISE 25.7. Let  $I$  be a real interval and let

$$f : I \longrightarrow \mathbb{R}$$

be a continuous function with antiderivative  $F$ . Let  $G$  be an antiderivative of  $F$  and let  $b, c \in \mathbb{R}$ . Determine an antiderivative of the function

$$(bt + c) \cdot f(t) .$$

EXERCISE 25.8. Let  $n \in \mathbb{N}_+$ . Determine an antiderivative of the function

$$\mathbb{R}_+ \longrightarrow \mathbb{R}_+, x \longmapsto x^{1/n},$$

using the antiderivative of  $x^n$  and Theorem 25.4.

EXERCISE 25.9. Determine an antiderivative of the natural logarithm function using the antiderivative of its inverse function.

EXERCISE 25.10. Let

$$f : [a, b] \longrightarrow [c, d]$$

be a bijective, continuous differentiable function. Prove the formula for the antiderivative of the inverse function by the integral

$$\int_a^b f^{-1}(y) dy$$

using the substitution  $y = f(x)$  and then integration by parts.

EXERCISE 25.11. Compute by an appropriate substitution an antiderivative of

$$\sqrt{3x^2 + 5x - 4}.$$

EXERCISE 25.12. Compute the definite integral of the function

$$f : \mathbb{R} \longrightarrow \mathbb{R}, x \longmapsto f(x) = 2x^3 + 3e^x - \sin x,$$

on  $[-1, 0]$ .

EXERCISE 25.13. Compute the definite integral of the function

$$f : \mathbb{R}_+ \longrightarrow \mathbb{R}, x \longmapsto f(x) = \sqrt{x} - \frac{1}{\sqrt{x}} + \frac{1}{2x+3} - e^{-x},$$

on  $[1, 4]$ .

### Hand-in-exercises

EXERCISE 25.14. (4 points)

Compute the definite integral  $\int_0^8 f(t) dt$ , where the function  $f$  is

$$f(t) = \begin{cases} t + 1, & \text{if } 0 \leq t \leq 2, \\ t^2 - 6t + 11, & \text{if } 2 < t \leq 5, \\ 6, & \text{if } 5 < t \leq 6, \\ -2t + 18, & \text{if } 6 < t \leq 8. \end{cases}$$

EXERCISE 25.15. (3 points)

Determine an antiderivative of the function

$$x^3 \cdot \cos x - x^2 \cdot \sin x .$$

EXERCISE 25.16. (2 points)

Determine an antiderivative of the function

$$\arcsin x .$$

EXERCISE 25.17. (4 points)

Determine an antiderivative of the function

$$\sin(\ln x) .$$

EXERCISE 25.18. (5 points)

Determine an antiderivative of the function

$$e^x \cdot \frac{x^2 + 1}{(x + 1)^2} .$$

EXERCISE 25.19. (5 points)

Let  $I$  be a real interval and let

$$f : I \longrightarrow \mathbb{R}$$

be a continuous function with antiderivative  $F$ . Let  $G$  be an antiderivative of  $F$  and  $H$  an antiderivative of  $G$ . Let  $a, b, c \in \mathbb{R}$ . Determine an antiderivative of the function

$$(at^2 + bt + c) \cdot f(t) .$$