

# Characteristics of Multiple Random Variables

Young W Lim

July 4, 2019

Copyright (c) 2018 Young W. Lim. Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.2 or any later version published by the Free Software Foundation; with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts. A copy of the license is included in the section entitled "GNU Free Documentation License".

This work is licensed under a Creative Commons "Attribution-NonCommercial-ShareAlike 3.0 Unported" license.



Based on  
Probability, Random Variables and Random Signal Principles,  
P.Z. Peebles,Jr. and B. Shi

# Outline

## 1 Linear Transformation of Gaussian Random Variables

# Linear Transform

$N$  Gaussian random variables

## Definition

linear transform of Gaussian random variables  $X_1, X_2, \dots, X_N$

$$Y_1 = a_{11}X_1 + a_{12}X_2 + \dots + a_{1N}X_N$$

$$Y_2 = a_{21}X_1 + a_{22}X_2 + \dots + a_{2N}X_N$$

$$\vdots \quad \quad \quad \vdots \quad \quad \quad \vdots \quad \quad \quad \vdots \quad \quad \quad \vdots$$

$$Y_N = a_{N1}X_1 + a_{N2}X_2 + \dots + a_{NN}X_N$$

$$[Y] = \begin{bmatrix} Y_1 \\ Y_2 \\ \vdots \\ Y_N \end{bmatrix}, [T] = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1N} \\ a_{21} & a_{22} & \dots & a_{2N} \\ \vdots & \vdots & \ddots & \vdots \\ a_{N1} & a_{N2} & \dots & a_{NN} \end{bmatrix}, [X] = \begin{bmatrix} X_1 \\ X_2 \\ \vdots \\ X_N \end{bmatrix}$$

# Linear Transform

$N$  Gaussian random variables

## Corollary

*linear transform of Gaussian random variables  $X_1, X_2, \dots, X_N$*

$$[Y] = [T][X], \quad [Y - \bar{Y}] = [T][X - \bar{X}]$$

$$[X] = [T]^{-1}[Y], \quad [X - \bar{X}] = [T]^{-1}[Y - \bar{Y}]$$

$$[Y] = \begin{bmatrix} Y_1 \\ Y_2 \\ \vdots \\ Y_N \end{bmatrix}, \quad [\bar{Y}] = \begin{bmatrix} \bar{Y}_1 \\ \bar{Y}_2 \\ \vdots \\ \bar{Y}_N \end{bmatrix}, \quad [X] = \begin{bmatrix} X_1 \\ X_2 \\ \vdots \\ X_N \end{bmatrix}, \quad [\bar{X}] = \begin{bmatrix} \bar{X}_1 \\ \bar{X}_2 \\ \vdots \\ \bar{X}_N \end{bmatrix}$$

$[T]$  is non-singular

# Linear Transform

$N$  Gaussian random variables

## Corollary

linear transform of Gaussian random variables  $X_1, X_2, \dots, X_N$

$$X_i = T_i^{-1}(Y_1, \dots, Y_N) = a^{i1} Y_1 + \dots + a^{iN} Y_N$$

$$\frac{\partial X_i}{\partial Y_j} = \frac{\partial T_i^{-1}}{\partial Y_j} = a^{ij}$$

$$X_i - \bar{X}_i = a^{i1}(Y_1, \dots, \bar{Y}_1) + \dots + a^{iN}(Y_N, \dots, \bar{Y}_N)$$

$a^{ij}$  denotes the  $ij$ -th element of  $[T]^{-1}$

## Using Jacobian

 $N$  Gaussian random variables

## Corollary

linear transform of Gaussian random variables  $X_1, X_2, \dots, X_N$

$$|J| = \|[T]^{-1}\| = \frac{1}{\|[T]\|}$$

$$C_{X_i X_j} = E[(X_i - \bar{X}_i)(X_j - \bar{X}_j)]$$

$$= \sum_{k=1}^N a^{ik} \sum_{m=1}^N a^{im} E[(Y_k - \bar{Y}_k)(Y_m - \bar{Y}_m)]$$

$$= \sum_{k=1}^N a^{ik} \sum_{m=1}^N a^{im} C_{Y_k Y_m}$$

# Using Jacobian

$N$  Gaussian random variables

## Corollary

$C_{X_i X_j}$  is the  $ij$ -th element of the covariance matrix  $[C_X]$   
and  $C_{Y_k Y_m}$  is the  $km$ -th element of the covariance matrix  $[C_Y]$

$$[C_X] = [T]^{-1} [C_Y] ([T]^t)^{-1}$$

$$[C_X]^{-1} = [T]^t [C_Y]^{-1} [T]$$

$$\|[C_X]^{-1}\| = \|[C_Y]^{-1}\| [T]^2$$





