


# First Order ODEs (H.1)

20151222

Copyright (c) 2015 Young W. Lim.

Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.2 or any later version published by the Free Software Foundation; with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts. A copy of the license is included in the section entitled "GNU Free Documentation License".

# Another kind of triangles and their slopes



ODE Background (0P)


8

Young Won Lim  
12/21/15

## Differentials and Derivatives (1)

$$\begin{aligned} dy &= f'(x) dx \\ dy &= \frac{df}{dx} dx \\ \text{differentials} && \text{derivative} \\ \frac{dy}{dx} &= f'(x) \end{aligned}$$

ratio                      not a ratio



ODE Background (0P)

12


Young Won Lim  
12/21/15

## Differentials and Derivatives (2)

$$\begin{aligned} f(x_1 + dx) &\approx f(x_1) + dy \\ &= f(x_1) + f'(x_1)dx \end{aligned}$$

for small enough  $dx$

$$\begin{aligned} f(x_1 + dx) &\stackrel{\textcircled{=}}{=} f(x_1) + dy \\ &= f(x_1) + f'(x_1)dx \end{aligned}$$



$$\lim_{dx \rightarrow 0} \frac{f(x_1 + dx) - f(x_1)}{dx} \stackrel{\textcircled{=}}{=} f'(x_1)$$

## Differentials and Derivatives (3)

$$dy = f'(x) dx \quad \rightarrow \quad \int dy = \int f'(x) dx$$

$$dy = \frac{df}{dx} dx \quad \rightarrow \quad \int dy = \int \frac{df}{dx} dx$$

$$dy = \dot{f} dx \quad y = f(x)$$

$$dy = D_x f dx \quad \int dy = \int 1 dy = y$$

# Integration Constant C

place a constant

place another constant

$$\int dy$$

$$= \int f'(x) dx$$

$$\int dy$$

$$= \int \frac{df}{dx} dx$$

$$y + C_1 = f(x) + C_2$$

$$y = f(x) + C$$

differs by a constant

place only one constant from the beginning

$$\int dy$$

$$= \int f'(x) dx$$

$$+ C$$

$$\int dy$$

$$= \int \frac{df}{dx} dx$$

$$+ C$$

$$y = f(x) + C$$

# Types of First Order ODEs

---

## A General Form of First Order Differential Equations

$$\frac{dy}{dx} = g(x, y)$$

$$y' = g(x, y)$$

## Separable Equations

$$\frac{dy}{dx} = g_1(x)g_2(y)$$

$$y' = g_1(x)g_2(y)$$

$$y = f(x)$$

## Linear Equations

$$a_1(x) \frac{dy}{dx} + a_0(x)y = g(x)$$

$$a_1(x)y' + a_0(x)y = g(x)$$

$$y = f(x)$$

## Exact Equations

$$M(x, y)dx + N(x, y)dy = 0$$

$$\frac{\partial z}{\partial x} dx + \frac{\partial z}{\partial y} dy = 0$$

$$z = f(x, y)$$

# 1차 미분 방정식


ch 2.

- { ① Separable eq.      편미분식      2.2
- ② Linear eq.      선형방정식      2.3
- ③ Exact eq.      완전방정식      2.4

↑

1st order differential eq

Solution:  $y(x)$ :  $y$ 는  $x$ 의 함수



ODE Ordinary Differential Equation 一階微分方程

PDE Partial Differential Equation 二階微分方程

$$\frac{df}{dx} \quad f(x) \quad \dots \quad \text{2階微分}$$

$$\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \quad f(x, y) \quad \dots \quad \text{3階微分}$$

~~~~~  
partial derivative.

# Separable Equation

1차 미분 방정식  $y' = \underbrace{g(x, y)}$   $y(x) = ?$

모든  $y$ 에 대해  $y'$ 를 미분

$x, y$ 의 수식

$$y' = [g_1(x)] \times [g_2(y)]$$

(x식)      (y식)      Separable

$$y' \left( \frac{1}{g_2(y)} \right) = (g_1(x))$$

$$\underline{y'} \left( y^{\frac{1}{g_2}} \right) = (x^{\frac{1}{g_1}})$$

$$\underline{y'} \left( y^{\frac{1}{g_2}} \right) dx = (x^{\frac{1}{g_1}}) dx$$

$$(y^{\frac{1}{g_2}}) \left[ \frac{dy}{dx} \right] dx = (x^{\frac{1}{g_1}}) dx$$

$$(y^{\frac{1}{g_2}}) \boxed{\frac{dy}{dx}} = (x^{\frac{1}{g_1}}) dx$$

$$\int (y^{\frac{1}{g_2}}) dy = \int (x^{\frac{1}{g_1}}) dx$$

$(y^{\frac{1}{g_2}})$  은

$y$ 에 대하여 적을

$(x^{\frac{1}{g_1}})$  은

$x$ 에 대하여 적을

$$\frac{dy}{dx} = 6y^2x \sim (6x)(y^2)$$

$$\textcircled{1} \quad y' = \frac{3x^2 + 4x - 4}{2y - 4} = (3x^2 + 4x - 4) \left( \frac{1}{2y - 4} \right)$$

$$\textcircled{2} \quad y' = \frac{xy^3}{\sqrt{1+x^2}} = \left( \frac{x}{\sqrt{1+x^2}} \right) (y^3)$$

$$\textcircled{3} \quad y' = e^{-y} (2x - 4) = (2x - 4) (e^{-y})$$


$$\textcircled{4} \quad \frac{dr}{d\theta} = \frac{r^2}{\theta} \quad r(\theta) \text{ を } \frac{r^2}{\theta} \text{ の極値を取る。}$$

$$\frac{d}{d\theta} r(\theta) = \frac{r^2(\theta)}{\theta} \Rightarrow r(\theta) ?$$

$$\frac{dr}{d\theta} = \frac{r^2}{\theta} = \left( \frac{1}{\theta} \right) (r^2)$$


$$\textcircled{5} \quad \frac{dy}{dt} = e^{y-t} \sec(y)(1+t^2) = (e^{-t}(1+t^2)) (e^y \sec(y))$$

$$\frac{dy}{dx} = 6y^2x \approx (6x)(y^2)$$



$$\frac{dy}{dx} = \frac{1}{y^2} = 6x$$

$$\frac{1}{y^2} \frac{dy}{dx} dx = 6x dx$$



$$\int \frac{1}{y^2} \left[ \frac{dy}{dx} dx \right] = \int 6x dx$$


$$\int \frac{1}{y^2} dy = \int 6x dx$$

$$-\frac{1}{3} y^{-3} = 3x^2 + C$$

$$y^{-3} = -9x^2 - 3C$$

$$y^3 = \frac{1}{-9x^2 - 3C}$$

$$y = \sqrt[3]{\frac{1}{-9x^2 - 3C}}$$



# Linear Equation

1차 미분 방정식  $y' = \underbrace{g(x, y)}$  y(x) ?  
x, y의 수식  
비율  
함수  $y_1, y_2$

Linear Eq

$$\frac{1}{a_1(x)} \left[ y' \right] + \frac{a_2(x)}{a_1(x)} \left[ y \right] = g(x)$$

$\frac{1}{a_1(x)}$   $a_1(x)$ 의 수식  
 $a_2(x)$   $x$ 의 수식  
 $a_1(x)$   $a_1(x)$ 의 수식

$$1 \cdot \left[ y' \right] + p(x) \left[ y \right] = g(x)$$

$p(x)$   $x$ 의 수식  
 $y'$   $x$ 의 수식

$$1. \boxed{y'} + p(x) \boxed{y} = g(x)$$

$$\textcircled{1} \quad \boxed{\frac{dv}{dt}} = 9.8 - 0.196 \boxed{v} \quad v(t) = ?$$

$$1. \boxed{\frac{dv}{dt}} + 0.196 \boxed{v} = 9.8$$

$$\textcircled{2} \quad \cos(x) \boxed{y'} + \sin(x) \boxed{y} = 2 \cos^3(x) \sin(x) - 1$$

$$1. \boxed{y'} + \frac{\sin(x)}{\cos(x)} \boxed{y} = 2 \cos^2(x) \sin(x) - \frac{1}{\cos(x)}$$

$$\textcircled{3} \quad t \boxed{y'} + 2 \boxed{y} = t^2 - t + 1$$

$$\boxed{y'} + \frac{2}{t} \boxed{y} = t - 1 + \frac{1}{t}$$

$$\textcircled{4} \quad t \boxed{y'} - 2 \boxed{y} = t^5 \sin(2t) - t^3 + 4t^4$$

$$1. \boxed{y'} - \frac{2}{t} \boxed{y} = t^4 \sin(2t) - t^2 + 4t^3$$

$$\textcircled{5} \quad 2 \boxed{y'} - \boxed{y} = 4 \sin(3t) \quad y(t) ?$$

$$1. \boxed{y'} - \frac{1}{2} \boxed{y} = 2 \sin(3t)$$

homogeneous eq

$$y' + p(x) y = 0 \quad \text{Sol } \underline{y_h}$$

$$\Leftrightarrow y'_h + p(x) y_h = 0$$

non-homogeneous eq

$$y' + p(x) y = Q(x) \quad \text{Sol } \underline{y_p}$$

$$\Leftrightarrow y'_p + p(x) y_p = Q(x)$$

$$y'_h + p(x) y_h = 0$$

$$y'_p + p(x) y_p = Q(x)$$

$$(y_p + y_h)' + p(x)(y_p + y_h) = Q(x)$$

$$\Leftrightarrow \boxed{y' + p(x) y = Q(x)} \quad \text{Sol } \underline{(y_p + y_h)}$$

homogeneous eq

$$y' + p(x) y = 0 \quad \text{Sol} \quad \underline{y_h} = C \cdot y_1 \\ = C \cdot e^{-\int p(x) dx}$$

non-homogeneous eq

$$y' + p(x) y = Q(x) \quad \text{Sol} \quad \underline{y_p} = u \cdot y_1 \\ = u(x) e^{-\int p(x) dx}$$

$$\textcircled{1} \quad \frac{dy}{dx} + y = 1$$

$$y = 1 + ce^{-x}$$

$$\textcircled{2} \quad \frac{dy}{dx} + y = x$$

$$y = (x-1) + ce^{-x}$$

$$\textcircled{3} \quad \frac{dy}{dx} + y = x^2$$

$$y = x^2 - 2x + 2 + ce^{-x}$$

$$y = 1$$

$$\frac{d}{dx}(1) + 1 = 0 + 1 = 1$$

$$y = x-1$$

$$\frac{d}{dx}(x-1) + (x-1) = 1 + x-1-x$$

$$y = x^2 - 2x + 2$$

$$\begin{aligned} & (x^2 - 2x + 2)' + (x^2 - 2x + 2) \\ &= 2x - 2 + x^2 - 2x + 2 \\ &= x^2 \end{aligned}$$

$$y = ce^{-x}$$

$$\begin{aligned} & (ce^{-x})' + (ce^{-x}) \\ &= -ce^{-x} + ce^{-x} = 0 \end{aligned}$$

$$y = ce^{-x}$$

$$\begin{aligned} & (ce^{-x})' + (ce^{-x}) \\ &= -ce^{-x} + ce^{-x} = 0 \end{aligned}$$

$$y = ce^{-x}$$

$$\begin{aligned} & (ce^{-x})' + (ce^{-x}) \\ &= -ce^{-x} + ce^{-x} = 0 \end{aligned}$$

$$y = 1 + ce^{-x}$$

$$(1 + ce^{-x})' + (1 + ce^{-x})$$

$$\begin{aligned} &= -ce^{-x} + 1 + ce^{-x} \\ &= 1 \end{aligned}$$

$$y = x-1 + ce^{-x}$$

$$(x-1 + ce^{-x})' + (x-1 + ce^{-x})$$

$$\begin{aligned} &= 1 - ce^{-x} + x - 1 + ce^{-x} \\ &= x \end{aligned}$$

$$y = x^2 - 2x + 2 + ce^{-x}$$

$$= x^2$$

$$a_1(x) \boxed{y'} + a_2(x) \boxed{y} = g(x)$$

General Solution

...  
infinitely  
many  
solutions

$y_p + y_h$

$$y_p$$

$$y_p + C \cdot \boxed{x^{\lambda_j}}$$

$$y_h$$

$$a_1(x) \boxed{y'} + a_2(x) \boxed{y} = 0$$

$$1 \boxed{y} + p(x) \boxed{y}$$

$$C \cdot e^{-\int p(x) dx}$$

## Solving the Homogeneous DE : finding $y_h$

$$\left( \frac{dy}{dx} + P(x)y = 0 \right)$$

$$y' + P(x)y = 0$$

homogeneous solution

$$y_h = f_h(x)$$

$$\frac{dy}{dx} = -P(x)y$$

$$y' = -P(x)y$$

$$\frac{1}{y} \frac{dy}{dx} = -P(x)$$

$$\frac{1}{y} y' = -P(x)$$

$$\int \frac{1}{y} \frac{dy}{dx} dx = - \int P(x) dx$$

$$\int \frac{1}{y} y' dx = - \int P(x) dx$$

$$dy = \frac{df}{dx} dx$$

$$\int \frac{1}{y} dy = - \int P(x) dx$$

$$\int \frac{1}{y} dy = - \int P(x) dx$$

$$\ln|y| = - \int P(x) dx + C$$

$$\ln|y| = - \int P(x) dx + C$$

$$|y| = e^{- \int P(x) dx + C}$$



$$|y| = e^{- \int P(x) dx + C}$$

$$\begin{aligned} y &= \pm e^{- \int P(x) dx} \cdot e^C \\ &= \pm e^C \cdot e^{- \int P(x) dx} \end{aligned}$$

$$y = c e^{- \int P(x) dx}$$

$$y = c e^{- \int P(x) dx}$$

## Derivatives of Inverse Functions



$$m_1 m_2 = f'(\text{a}) g'(\text{b}) = 1 \Rightarrow g'(\text{b}) = \frac{1}{f'(\text{a})} \Rightarrow g'(\text{b}) = \frac{1}{f'(g(\text{b}))}$$

$$g(\text{b}) = \text{a}$$

ODE Background: Integral (2A)

42

Young Won Lim  
12/23/15

## Derivatives of Inverse Functions

$$m_1 m_2 = f'(\text{a}) g'(\text{b}) = 1 \Rightarrow g'(\text{b}) = \frac{1}{f'(\text{a})} \Rightarrow g'(\text{b}) = \frac{1}{f'(g(\text{b}))}$$

$$g(\text{b}) = \text{a} \qquad \qquad \qquad g'(x) = \frac{1}{f'(g(x))}$$

To find  $g'(x)$   
 (1) find  $f'(x)$   
 (2) find  $1/f'(x)$   
 (3) substitute  $x$  with  $g(x)$

|                                                                                             |                                                                                                            |                                                                                                                        |
|---------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------|
| $\frac{d}{dx} \ln x \Rightarrow \frac{1}{x}$<br><br>$f'(x)$<br>$\frac{1}{f'(x)}$<br>$g'(x)$ | $(e^x)' \Rightarrow e^x$<br>$\Rightarrow \frac{1}{e^x}$<br>$\Rightarrow \frac{1}{e^{\ln x}} = \frac{1}{x}$ | $\frac{d}{dx} e^x \Rightarrow e^x$<br>$(\ln x)' = \frac{1}{x}$<br>$\Rightarrow \frac{1}{1/x} = x$<br>$\Rightarrow e^x$ |
|---------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------|

ODE Background: Integral (2A)

43

Young Won Lim  
12/23/15

## Derivative of $\ln |x|$

$$\frac{d}{dx} \ln x = \frac{1}{x} \quad x > 0$$

$$\frac{d}{dx} \ln(-x) = \frac{1}{(-x)} \cdot (-1) = \frac{1}{x} \quad x < 0$$

$$\begin{array}{ll} \ln x & x > 0 \\ \ln(-x) & x < 0 \end{array} \quad \begin{array}{l} \frac{d}{dx} \rightarrow \\ \frac{d}{dx} \end{array} \quad \begin{array}{l} \frac{1}{x} \\ \frac{1}{x} \end{array}$$

$$\frac{d}{dx} \ln|x| = \frac{1}{x} \quad x \neq 0$$

## Method of Solving First Order ODEs

$$\frac{dy}{dx} + P(x)y = Q(x)$$

$y_p?$

$$y' + P(x)y = Q(x)$$

$$y = f(x)$$

$$\left[ e^{\int P(x)dx} \right] \cdot \frac{dy}{dx} + P(x)y = \left[ e^{\int P(x)dx} \right] Q(x)$$

$$y_1 = e^{-\int P(x)dx} \quad \frac{1}{y_1} = e^{\int P(x)dx}$$

$$\left[ e^{\int P(x)dx} \right] \frac{dy}{dx} + \left[ \left[ e^{\int P(x)dx} \right] P(x) \right] y = \left[ e^{\int P(x)dx} \right] Q(x)$$

$$\left[ e^{\int P(x)dx} \right] P(x) = \frac{d}{dx} \left[ e^{\int P(x)dx} \right] \quad f'(g(x))g'(x)$$

$$\left[ e^{\int P(x)dx} \right] \frac{dy}{dx} + \frac{d}{dx} \left[ e^{\int P(x)dx} \right] y = \left[ e^{\int P(x)dx} \right] Q(x)$$

$$\left[ e^{\int P(x)dx} \right] \frac{dy}{dx} + \frac{d}{dx} \left[ e^{\int P(x)dx} \right] y = \frac{d}{dx} \left[ e^{\int P(x)dx} \cdot y \right]$$

$$\frac{d}{dx} \left[ \left[ e^{\int P(x)dx} \right] \cdot y \right] = \left[ e^{\int P(x)dx} \right] Q(x)$$

$$f'(x)g(x) + f(x)g'(x)$$

$$\int \frac{d}{dx} \left[ \left[ e^{\int P(x)dx} \right] \cdot y \right] dx = \int \left[ e^{\int P(x)dx} \right] Q(x) dx + c$$

$$\left[ \left[ e^{\int P(x)dx} \right] \cdot y \right] = \int \left[ e^{\int P(x)dx} \right] Q(x) dx + c \quad \Rightarrow$$

$$y(x) = ce^{-\int P(x)dx} + e^{-\int P(x)dx} \cdot \left[ \int \left[ Q(x) \cdot e^{\int P(x)dx} \right] dx \right]$$

# 1st Order Linear Equations

$y_p$  part)

$$1 \cdot y' + \boxed{P(x)} y = Q(x)$$

$$y_h = C \cdot e^{-\int P(x) dx}$$

$$y_p = u(x) e^{-\int P(x) dx}$$

$$e^{+\int P(x) dx}$$

Integrating Factor

{ • const. c X  
 • sign X

$$e^{+\int P(x) dx}$$

$$(1 \cdot y' + \boxed{P(x)} y) = (Q(x))$$

$$e^{+\int P(x) dx}$$

$$(e^{+\int P(x) dx} \cdot \boxed{y_p})' = Q(x) e^{+\int P(x) dx}$$

$$e^{+\int P(x) dx} \cdot y_p = \int Q(x) e^{+\int P(x) dx} dx$$

$$y_p = e^{-\int P(x) dx} \cdot \int Q(x) e^{+\int P(x) dx} dx$$

$$y = y_h + y_p$$

$$y = \underline{\underline{C \cdot e^{-\int P(x) dx}}} + \underline{\underline{e^{-\int P(x) dx} \cdot \int Q(x) e^{+\int P(x) dx} dx}}$$

$y_h$                      $y_p$

Calculus 1

Review : Exponential

Review : Logarithmic

Derivatives : Trig Derivatives

Differential Equation

First Order : Linear Equation

First Order Differential Equations (1P.pdf) .....  
Linear Equation (2A.pdf)

cf) Partial Derivatives (9.4 Zill & Wright)

## Exact Equation

$$M(x, y) dx + N(x, y) dy = 0$$

$y(x) = ?$  ... 1 variable  
 $f(x, y)$  ... 2 variables

Exact  $\rightarrow f(x, y)$  exists

total differential df =  $\frac{\partial f}{\partial x}$  dx +  $\frac{\partial f}{\partial y}$  dy  
=  $M(x, y) dx + N(x, y) dy = 0$

$$\frac{\partial}{\partial y} \left( \frac{\partial f}{\partial x} \right) = \frac{\partial}{\partial x} \left( \frac{\partial f}{\partial y} \right)$$

$$\frac{\partial}{\partial y}(M) = \frac{\partial}{\partial x}(N)$$

$f(x, y)$   $df = 0$   $f(x, y) = C$   
  $y(x)$  ← Curve ← Surface  $f(x, y) = C$

$$(2xy - 9x^2) + (2y + x^2 + 1) \frac{dy}{dx} = 0$$

$$(2xy - 9x^2) dx + (2y + x^2 + 1) dy = 0$$

$$\begin{array}{c} \boxed{\begin{array}{c} M \\ \frac{\partial F}{\partial x} \end{array}} \\ || \\ dx \end{array} + \begin{array}{c} \boxed{\begin{array}{c} N \\ \frac{\partial F}{\partial y} \end{array}} \\ || \\ dy \end{array} = 0$$

$$dx + \begin{array}{c} \boxed{\begin{array}{c} N \\ \frac{\partial F}{\partial y} \end{array}} \\ || \\ dy \end{array} = 0$$

\* Exact?

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

$$2x = 2x \quad \therefore \text{exact}$$

$$f(x, y) = \left\{ \begin{array}{l} \frac{\partial f}{\partial x} dx + C \\ \frac{\partial f}{\partial x} dx + g(y) \end{array} \right.$$

variable  $y$  may be included

$$= \int (2xy - 9x^2) dx + g(y)$$

$$f(x, y) = x^2y - 3x^3 + g(y) ?$$

$$\frac{\partial f}{\partial y} = x^2 + g'(y) = (2y + x^2 + 1)$$

$$g'(y) = 2y + 1$$

$$g(y) = y^2 + y$$

$$f(x, y) = x^2y - 3x^3 + y^2 + y$$

$$df = 0$$

$$f(x, y) = x^2y - 3x^3 + y^2 + y = C$$

$$\underline{y(x)}$$

$$M(x, y) dx + N(x, y) dy = 0 \implies df = 0$$

$\downarrow$        $y(x) = ?$  ... 1 variable  $f(x, y) = c$

$f(x, y)$  ... 2 variables  $f(x, y(x))$

$$M(x, y) dx + N(x, y) dy = 0$$

$$\frac{\partial f}{\partial x} \quad \text{"} \frac{\partial f}{\partial y} \text{"}$$

① Verify "exactness"

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

②

$$M(x, y)$$

$$\frac{\partial f}{\partial x}$$

$$f(x, y) = \int M(x, y) dx + g(y)$$

③

$$N(x, y)$$

$$\frac{\partial f}{\partial y}$$

$$\frac{\partial f}{\partial y} = \frac{\partial}{\partial y} \left[ \int M(x, y) dx \right] + g'(y) \\ = N(x, y)$$

④

$$f(x, y) = C$$

flat surface  $\cap$   $y = y(x)$

## Non-Exact Case

Zill Ch 4)

$$xy \, dx + (2x^2 + 3y^2 - 20) \, dy = 0$$

$$\frac{\partial}{\partial y} (xy) = x \neq \frac{\partial}{\partial x} (2x^2 + 3y^2 - 20) = 4x$$

non-exact eq

$y^3$

$\leftarrow$

Integrating Factor  $\mu(x)$   $\mu(y)$

$$x y^4 \, dx + (2x^2 + 3y^2 - 20) y^3 \, dy = 0$$

$$x y^4 \, dx + (2x^2 y^3 + 3y^5 - 20y^3) \, dy = 0$$

$$\frac{\partial}{\partial y} (x y^4) = 4x y^3 = \frac{\partial}{\partial x} (2x^2 y^3 + 3y^5 - 20y^3) = 4x y^3$$

Now exact!

## Non-Exact Case : Finding Integrating Factor

$$\text{Exact} \rightarrow \frac{\partial M}{\partial y} = \frac{\partial N}{\partial x} \quad M_y = N_x$$

$$M_y - N_x = 0$$

$$\text{Non-Exact} \rightarrow M_y - N_x \neq 0$$

$$\frac{M_y - N_x}{N} : x \text{의 } \frac{d}{dx} \checkmark \quad \text{IF.} = e^{\int \frac{M_y - N_x}{N} dx}$$

$$- \frac{M_y - N_x}{M} : y \text{의 } \frac{d}{dy} \checkmark \quad \text{IF.} = e^{-\int \frac{M_y - N_x}{M} dy}$$

## Integrating Factor

$\left\{ \begin{array}{l} \cdot \text{ sign } x \\ \cdot \text{ const } \propto x \end{array} \right.$

$$\mu(x) = e^{\int \frac{M_y - N_x}{N} dx} \Leftrightarrow \frac{M_y - N_x}{N} \dots x \text{의 수식}$$

$$\mu(y) = e^{\int -\frac{M_y + N_x}{M} dy} \Leftrightarrow -\frac{M_y + N_x}{M} \dots y \text{의 수식}$$

$$M(x, y) dx + N(x, y) dy = 0$$

$$\frac{\partial M}{\partial y} = M_y \quad \frac{\partial N}{\partial x} = N_x$$

difference

$x$  또는  
 $y$  수식

$$xy \, dx + (2x^2 + 3y^2 - 20) \, dy = 0$$

$$\frac{\partial}{\partial y} (xy) = x \neq \frac{\partial}{\partial x} (2x^2 + 3y^2 - 20) = 4x$$

non-exact eq

$$\frac{\partial M}{\partial y} = M_y = x$$

$$\frac{\partial N}{\partial x} = N_x = 4x$$

x only expression

$$\mu(x) = e^{\int \frac{M_y - N_x}{N} dx}$$

$$\mu(x) = C \int \frac{-3x}{(2x^2 + 3y^2 - 20)} dx$$

$$xy \, dx + (2x^2 + 3y^2 - 20) \, dy = 0$$

$$\frac{\partial}{\partial y} (xy) = x \neq \frac{\partial}{\partial x} (2x^2 + 3y^2 - 20) = 4x$$

non-exact eq

$$\frac{\partial M}{\partial y} = M_y = x \quad \frac{\partial N}{\partial x} = N_x = 4x$$

x only expression

$$\mu(y) = e^{\int \frac{-M_y + N_x}{M} dy} \Leftarrow -M + N \dots y \text{ only}$$

$$\mu(y) = e^{\int \frac{3x}{xy} dy} = e^{3 \int \frac{1}{y} dy} = e^{\ln(y)^3} \\ = [y]^3$$

$$\mu(y) = y^3, -y^{-3} \Rightarrow y^3$$

$$xy^3 \, dx + (2x^2 + 3y^2 - 20) y^3 \, dy = 0$$

# integrating factor 통계우도 과정

$$(f \cdot g)' = f'g + fg'$$

①  $\frac{\partial(\mu M)}{\partial y} = \frac{\partial(\mu N)}{\partial x}$

$\mu(x)$   
이상적입니다.

②  $\frac{\partial M}{\partial y} \neq \frac{\partial N}{\partial x}$

$$\frac{\partial \mu}{\partial y} M + \mu \frac{\partial M}{\partial y} = \frac{\partial \mu}{\partial x} N + \mu \frac{\partial N}{\partial x}$$

$$\mu \frac{\partial M}{\partial y} = \frac{d\mu}{dx} N + \mu \frac{\partial N}{\partial x}$$

$$\frac{d\mu}{dx} N = \mu \frac{\partial M}{\partial y} - \mu \frac{\partial N}{\partial x}$$

$$\frac{d\mu}{dx} N = \mu M_y - \mu N_x$$

$$\frac{d\mu}{dx} = \left( \frac{M_y - N_x}{N} \right) \mu$$

$$\frac{d\mu}{dx} = P(x)\mu$$

이는  $d\mu/dx = P(x)\mu$ 입니다.

이

$$\frac{d\mu}{dx} - P(x)\mu = 0$$

$$\mu(x) = e^{\int P(x) dx}$$

$$\mu(x) = e^{\int \left( \frac{M_y - N_x}{N} \right) dx}$$

# Substitution Method

$$\frac{dy}{dx} = g(x, y) \quad \text{...} \quad \left(\frac{y}{x}\right) \text{ 항은 } u \text{로 대체하는 경우입니다.}$$

$\Leftrightarrow u \quad y = ux$

Substitute

$$\text{ex)} \quad y = ux$$

$$y = h(x, u)$$

$$u = \Phi(x)$$

$$\frac{dy}{dx} = \frac{\partial h}{\partial x} + \frac{\partial h}{\partial u} \frac{du}{dx}$$

$$\frac{d}{dx} y = \frac{d}{dx} h(x, u)$$

$$\frac{\partial h}{\partial x} = u \quad \frac{\partial h}{\partial u} = x \quad \frac{du}{dx} = u' \quad \Rightarrow \quad u + xu'$$

$$g(x, h(x, u)) = \frac{\partial h}{\partial x} + \frac{\partial h}{\partial u} \frac{du}{dx}$$

$$g(x, h(x, u)) = h_x(x, y) + h_u(x, u) \frac{du}{dx}$$

$$\frac{dy}{dx} = g(x, y)$$

\* Substitute

$$(y)$$

and

$$(y')$$

$$\left(\frac{y}{x}\right) \rightarrow u$$

$$(y) \rightarrow ux$$

$$(y') \rightarrow u'x + u$$

both  $y$  &  $y'$

$$x y y' + 4x^2 + y^2 = 0$$

↑

divide by  $x^2$

$$\left(\frac{y}{x}\right)y' + 4 + \left(\frac{y}{x}\right)^2 = 0$$

↑                      ↑                       $u = \frac{y}{x} = \frac{y(x)}{x}$

Substitute both  $y$  and  $y'$

$$\frac{x \text{의 } u \text{ } \cancel{\text{을}}}{x \text{의 } u}$$

$$\textcircled{y} = xu$$

$$\textcircled{y'} = u + xu'$$

$$\left(\frac{y}{x}\right)y' + 4 + \left(\frac{y}{x}\right)^2 = 0$$

$$\left(\frac{xu}{x}\right) \cdot (u + xu') + 4 + \left(\frac{xu}{x}\right)^2 = 0$$

$$u(u + xu') + 4 + u^2 = 0$$

$$2u^2 + xuu' + 4 = 0$$

$$2u^2 + xu' + 4 = 0$$

$$\frac{xu'u}{u} = -\frac{2u^2 + 4}{u}$$

$$xu' = -\frac{4+2u^2}{u}$$

$$\left(\frac{u}{4+2u^2}\right) u' = -\frac{1}{x}$$

$$u' = \frac{du}{dx}$$

$$\frac{u}{4+2u^2} \cdot \frac{du}{dx} = -\frac{1}{x} dx$$

$$\frac{u}{4+2u^2} du = -\frac{1}{x} dx$$

$$\frac{1}{4} \int \frac{4u}{4+2u^2} du = \int -\frac{1}{x} dx$$

$$\ln(4+2v^2)^{\frac{1}{4}} = \ln(x)^{-1} + c$$

sides and do a little rewriting

$$(4+2v^2)^{\frac{1}{4}} = e^{\ln(x)^{-1} + c} = e^c e^{\ln(x)^{-1}} = \frac{c}{x}$$

$$\frac{1}{4} \ln |4+2u^2| = -\ln |x| + C$$

$$\ln |4+2u^2|^{\frac{1}{4}} = \ln |x|^{-1} + C$$

$$|4+2u^2|^{\frac{1}{4}} = e^{\ln|x|^{-1} + C} = e^{\ln|x|^{-1}} e^C$$

$$(4+2u^2)^{\frac{1}{4}} = \frac{1}{x} C$$

$$(4+2u^2)^{\frac{1}{4}} = \frac{1}{x}(\textcircled{C})$$

$$(4+2u^2) = \left(\frac{c}{x}\right)^4 \quad u = \frac{y}{x}$$

$$4+2\left(\frac{y}{x}\right)^2 = \left(\frac{c}{x}\right)^4$$

$$2\left(\frac{y}{x}\right)^2 = \left(\frac{c_1}{x^4} - 4\right)$$

$$\frac{y^2}{x^2} = \frac{1}{2} \left( \frac{c_1 - 4x^4}{x^4} \right)$$

$$y^2 = \frac{1}{2} x^2 \left( \frac{c_1 - 4x^4}{x^4} \right)$$

$$= \frac{c_1 - 4x^4}{2x^2}$$

$$xy' = y(\ln x - \ln y)$$

$$x > 0$$

$$xy' = y \ln\left(\frac{x}{y}\right)$$

$$y' = \left(\frac{y}{x}\right) \ln\left(\frac{x}{y}\right)$$

$$\frac{y}{x} = u \quad y = xu$$

$$(u + xu') = u \ln\left(\frac{1}{u}\right)$$

$$y' = u + xu'$$

$$xu' = u \ln\left(\frac{1}{u}\right) - u$$

$$\int \frac{1}{u \left( \ln\left(\frac{1}{u}\right) - 1 \right)} \frac{du}{dx} dx = \int \frac{1}{x} dx$$

$$\int \frac{1}{u \left( \ln\left(\frac{1}{u}\right) - 1 \right)} du = \int \frac{1}{x} dx$$

$$\int \frac{-1}{\left( \ln\left(\frac{1}{u}\right) - 1 \right)} \left( \frac{-1}{u} du \right)$$

$$\boxed{\begin{aligned} t &= \ln\left(\frac{1}{u}\right) - 1 \\ dt &= \left(\frac{1}{u}\right)\left(\frac{1}{u}\right)' du \\ dt &= u \cdot (-u^{-2}) du \\ dt &= -\frac{1}{u} du \end{aligned}}$$

$$-\int \frac{1}{\left( \ln\left(\frac{1}{u}\right) - 1 \right)} dt = \int \frac{1}{x} dx$$

$$dt = u du$$

$$\ln\left(\ln\left(\frac{1}{u}\right) - 1\right) = -(\ln x + c)$$

$$\ln\left(\ln\left(\frac{1}{u}\right) - 1\right) \therefore = C - \ln x$$

$$\ln(\ln(\frac{t}{n}) - i) \cdot \cdot \cdot = C - \ln x$$

$$\begin{aligned}\ln(\frac{t}{n}) - i &= e^{C - \ln x} \\ &= c_1 e^{\ln \frac{1}{x}} \\ &= \frac{c_1}{x}\end{aligned}$$

$$\ln(\frac{t}{n}) - i = \frac{c_1}{x}$$

$$\ln(\frac{t}{n}) = \frac{c_1}{x} + 1$$

$$\frac{1}{n} = e^{\frac{c_1}{x} + 1}$$

$$u = e^{-\frac{c_1}{x} - 1}$$

$$\frac{y}{x} = e^{-\frac{c_1}{x} - 1}$$

$$y = x \cdot e^{-\frac{c_1}{x} - 1}$$

$$\star \int \tan x \, dx \rightarrow \text{substitution } u = \cos x$$

$$\int t^2 \sin(2t) dt \rightarrow$$

$$\int \tan x \, dx = \int \frac{\sin x}{\cos x} \, dx \quad u = \cos x \\ du = -\sin x \, dx$$

$$= \int \frac{-1}{u} \, du = -\ln|u| = \ln|u|^{-1}$$

$$= \ln\left|\frac{1}{\cos x}\right| = \ln|\sec x|$$

integrating factor  $\mu(x)$

$\rightarrow$  multiplied to the original differential eq

$$y' + p(x) y = Q(x)$$

$$\pm \mu(x) (y' + p(x) y) = (Q(x)) \pm \mu(x)$$

(Op Ex 6)

2F  
3g

$$\int f \ g' dt = f \ g - \int f' \ g dt$$

$$\int t^2 \sin(2t) dt$$

$$= -\frac{1}{2} t^2 \cos(2t) - \int (t^2)' \left[ \frac{1}{2} \cos(2t) \right] dt$$

$$= -\frac{1}{2} t^2 \cos(2t) + \left[ \int t \cos(2t) dt \right]$$

$$\left[ t \left[ \frac{1}{2} \sin(2t) \right] - \int t' \left[ \frac{1}{2} \sin(2t) \right] dt \right]$$

$$\frac{1}{2} t \sin(2t) - \frac{1}{2} \int \sin(2t) dt$$

$$-\frac{1}{2} \left[ -\frac{1}{2} \cos(2t) \right]$$

$$-\frac{1}{2} t^2 \cos(2t) + \frac{1}{2} t \sin(2t) + \frac{1}{4} \cos(2t)$$

$$-\frac{1}{2} \left( t^2 - \frac{1}{2} \right) \cos(2t) + \frac{1}{2} t \sin(2t)$$

(%i2) `integrate(% , t);`

(%o2)  $\frac{4t \sin(2t) + (2 - 4t^2) \cos(2t)}{8}$

$$\int f(x)g'(x) dx = f(x)g(x) - \int f'(x)g(x) dx$$

$$\int xe^x dx = xe^x - \int e^x dx = xe^x - e^x + c_1 = (x-1)e^x + c_1$$

$$\int x^2e^x dx = x^2e^x - 2\int xe^x dx = x^2e^x - 2xe^x + 2e^x + c_2 = (x^2 - 2x + 2)e^x + c_2$$

$$\int x^3e^x dx = x^3e^x - 3\int x^2e^x dx = x^3e^x - 3x^2e^x + 6xe^x - 6e^x + c_3 = (x^3 - 3x^2 + 6x - 6)e^x + c_3$$

$$\int xe^x dx = (x-1)e^x + c_1$$

$$\int x^2e^x dx = (x^2 - 2x + 2)e^x + c_2$$

$$\int x^3e^x dx = (x^3 - 3x^2 + 6x - 6)e^x + c_3$$

$$\int e^x dx = \int \left[ \frac{d}{dx} e^x \right] dx = e^x + c$$

$$\int xe^{x^2/2} dx = \int \left[ \frac{d}{dx} e^{x^2/2} \right] dx = e^{x^2/2} + c$$

$$\int x^2 e^{x^3/3} dx = \int \left[ \frac{d}{dx} e^{x^3/3} \right] dx = e^{x^3/3} + c$$



Chain Rule

$$f(g(x)) \xrightarrow{\frac{d}{dx}} f'(g(x)) g'(x)$$

$\therefore \int \cdot dx$

Substitution rule

Product Rule

$$(fg)' = f'g + \cancel{fg'}$$

$$fg' = (f \cdot g)' - f'g$$

$$\int fg' dx = \int (f \cdot g)' dx - \int f'g dx$$

$$\int f g' dx = fg - \int f' g dx$$

Pauls online math note  
Calculus 1

Integral :

Computing Indefinite Integrals  
Substitution Rule  
More about substitution rule

a b  
c d  
e f  
g h

Differential Equation  
1st Order

Exact Equation  
Substitution

CR&P

range

Graph X

$x_0 \neq 3m$   
initial cond

IVP

Substitution Rule

$$\int f(g(x)) g'(x) dx = \int f(u) du$$

||  
u

Chain rule

$$\frac{d}{dx} f(g(x)) = \frac{d}{dg} f(g) \frac{d}{dx} g(x)$$

