Laurent Series and z-Transform

Geometric Series Double Pole Examples A

7	\cap 1	L8	\cap	1	\mathbf{T}	1
/		ı×	()	≺		
_ \	$oldsymbol{\cup}$	${f L}$	V	_	_	_

Copyright (c) 2016 - 2018 Young W. Lim.

Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.2 or any later version published by the Free Software Foundation; with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts. A copy of the license is included in the section entitled "GNU Free Documentation License".

2 formulas of z

$$\frac{-1}{(2-1)(2-2)} = \left(\frac{2-1}{2-1} - \frac{2-2}{2-2}\right)$$

$$\frac{-05\xi^{2}}{(2-1)(2-0.5)} = \left(-\frac{\xi}{(\xi-1)} + \frac{0.5 \xi}{(\xi-0.5)}\right)$$

$$\frac{-1}{(2^{-1})(2^{-2})} = \left(\frac{1}{\xi - 1} - \frac{1}{\xi - 2}\right)$$

$$= \left(\frac{1}{\xi - 0.5} - \frac{1}{\xi - 2}\right)$$

$$= \left(\frac{1}{\xi^{-1} - 1} - \frac{1}{\xi^{-2}}\right)$$

$$= \left(\frac{\xi}{1 - \xi} - \frac{\xi}{1 - 2\xi}\right)$$

$$= \left(\frac{-\xi}{2 - 1} + \frac{0.5\xi}{2 - 0.5}\right)$$

$$= \xi \left(\frac{-0.5\xi}{(\xi - 1)(\xi - 0.5)}\right)$$

$$= \frac{-0.5\xi^{2}}{(\xi - 1)(\xi - 0.5)}$$

$$\frac{-0.52^{2}}{(2-1)(2-0.5)} = \left(-\frac{2}{(2-1)} + \frac{0.52}{(2-0.5)}\right)$$

$$\frac{(3-1)(3-2)}{-1} = \left(\frac{\xi-1}{1} - \frac{\xi-2}{1}\right)$$

$$\frac{f(z)}{|z| > 2} \qquad \frac{f(z)}{|z|} = \frac{z^{-1}}{|z|^{-2}} - \frac{z^{-1}}{|z|^{-2}} + |z|^{n+1} - (\frac{1}{2})^{n+1} \qquad (n < 0)$$

$$\frac{\chi(\xi)}{|\xi| > 2} \qquad \chi(\xi) = \frac{\xi^{-1}}{|-\xi^{-1}|} - \frac{\xi^{-1}}{|-2\xi^{-1}|} \qquad + \frac{|-1|}{|-2|^{n-1}} - \frac{|-1|}{|-2|}$$

$$\frac{-05z^2}{(z-1)(z-0.5)} = \left(-\frac{z}{(z-1)} + \frac{0.5z}{(z-0.5)}\right)$$

(2)-(A)
$$|\xi| < 05$$
 $f(\xi) = + \frac{\xi}{|-\xi|} - \frac{\xi}{|-\chi\xi|} |n-1| - 2^{n-1} (n > 1)$

$$\frac{\chi(5)}{|\xi| > 1} \qquad \chi(5) = -\frac{1}{1 - \epsilon_{-1}} + \frac{0.5}{1 - 0.5 \epsilon_{-1}} \qquad - \frac{1}{n+1} + \left(\frac{1}{7}\right)_{u+1} \qquad (u > 0)$$

 $(p,q) = (1,2) \qquad (p,q) = (0.5, 1)$ $-\frac{1}{(2-1)(2-2)} \qquad 2 \qquad \frac{-0.5 z^2}{(2-1)(2-0.5)}$ $A) \quad |z| 0) \qquad +\frac{n-1}{1+2^{n-1}} (n>1)$ $f(z) \quad |z| > p \qquad +\frac{n+1}{1+2^{n-1}} (n<1) \qquad +\frac{n+1}{1+2^{n-1}} (n<1)$ $B) \quad |z|
<math display="block">X(z) \quad |z| > q \qquad +\frac{1}{1+2^{n-1}} (n>1) \qquad -\frac{1}{1+2^{n+1}} (n>0)$

	_	(P, 4) = (1, 2)	(7,4)=(0.5,1)
		(1) (2-1) (2-2)	$2 \frac{-052^{2}}{(2-1)(2-0.5)}$
121 < 10	f(2)	$\frac{-1}{n+1} + \left(\frac{7}{17}\right)_{U+1} (U > 0)$	$+ ^{n-1} - 2^{n-1} (n \ge)$
z < P	Χ(₹)		$+ _{u+1} - (\frac{7}{1})_{u+1} (u < 0)$
	f(2)	$+ \frac{1}{n+1} - (\frac{1}{2})^{n+1} $ (n<0)	$-1^{n-1}+2^{n-1}(n<1)$
& > 	X(2)	+ n-1 - 2 n-1 (n>1)	$\frac{- \left(\frac{2}{1}\right)^{n+1} + \left(\frac{2}{1}\right)^{n+1} (n \ge 0)}{n+1}$

(P, 4) = (0.5, 1)(7,4)=(1,2) $2 \frac{-052^2}{(2-1)(2-0.5)}$ (-| (2-1)(2-2) 121 < P f(2) causal (n>1) causal (n>0) anticausal (n<1) |2| > B f(2) anticausal (n<0) anticausal (n<1) X(z)121 < P anticausal (N<0) causal (170) causal (n>1) X(z)121 > 8

(P, 4) = (0.5, 1)(P, 4)=(1, 2) -0522 (2-1)(2-0.5) causal (n>1) 121 < p | f(2) causal (n>0) X(Z) |2| < |2 anticausal (n<1) anticausal (n<0) anticausal (N<0) f(2) anticausal (M<1) 121 > B X(z)causal (931) causal (n>0) 121 > 8

$$\frac{-1}{(2-1)(2-2)} = 2 \frac{-05z^2}{(2-1)(2-0.5)}$$

$$\left(\frac{1}{\xi-1}-\frac{1}{\xi-2}\right)$$

$$\left(\frac{2}{(2-1)} + \frac{0.5 z}{(2-0.5)}\right)$$

$$\frac{2}{|-\xi|} + \frac{0.5}{|-0.5\xi|}$$

$$+\frac{z}{|-z|}$$

15/<1 |5/<1 |0.58)<1

[2]<0.5 |2]<1 |22]<1

$$\frac{\xi^{1}}{|-\xi^{1}|} - \frac{\xi^{1}}{|-2\xi^{1}|}$$

$$\frac{2^{-1}}{1-\epsilon^{-1}}+\frac{0.5}{1-0.5\epsilon^{-1}}$$

|と| 72 | ほりく| | 1287 | く |

|&|7 | |&1|<| |0.521|<|

$$\frac{-1}{(2-1)(2-2)} = 2 \frac{-05\xi^2}{(2-1)(2-0.5)}$$

$$\left(\frac{1}{\xi-1}-\frac{1}{\xi-2}\right)$$

$$\left(-\frac{2}{(2-1)}+\frac{0.5 z}{(2-0.5)}\right)$$

$$\frac{2}{1-\xi} + \frac{0.5}{1-0.5\xi} + \frac{\xi}{1-\xi} - \frac{\xi}{1-2\xi}$$

$$(1 \leq n)$$

$$\frac{z^{1}}{|-z^{1}} - \frac{z^{1}}{|-2z|}$$

$$\frac{\xi^{-1}}{|-\xi^{-1}|} - \frac{\xi^{-1}}{|-2\xi^{-1}|} = \frac{|-\xi^{-1}|}{|-\xi^{-1}|} + \frac{0.5}{|-0.5\xi^{-1}|}$$

$$(n \ge 1)$$

$$X(2)$$
 causal $(n \ge 0)$

$$\frac{-1}{(2-1)(2-2)} = 2 \frac{-052^2}{(2-1)(2-0.5)}$$

$$\left(\frac{1}{\xi-1}-\frac{1}{\xi-2}\right) \qquad \left(-\frac{\xi}{(\xi-1)}+\frac{0.5 z}{(\xi-0.5)}\right)$$

$$\frac{2}{1-z} + \frac{0.5}{1-0.5z} + \frac{z}{1-z} - \frac{z}{1-z}$$

$$|Z| < 1$$
 f(z) causal $(n > 0)$ $|Z| < 0.5$ f(z) causal $(n > 1)$


$$X(z)$$
 anticausal $(n \le 0)$ $X(z)$ anticausal $(n \le -1)$

$$\frac{z^{-1}}{|-z^{-1}|} - \frac{z^{-1}}{|-2z^{-1}|} - \frac{|-z^{-1}|}{|-z^{-1}|} + \frac{0.5}{|-0.5z^{-1}|}$$

|
$$|\xi|$$
72 | f(z) anticausal (n \leq -1) | $|\xi|$ 7| | f(z) anticausal (n \leq 0) | X(z) causal (n \geq 0)

$$(n > 0) \xrightarrow{-n} (n < 0) \equiv (n < 1)$$

$$(n > 1) \xrightarrow{-n} (n < -1) \equiv (n < 0)$$



$$\frac{-1}{(2-1)(2-2)} = 2 \frac{-052^2}{(2-1)(2-0.5)}$$

151<1

$$-\frac{1}{1-\xi}+\frac{0.5}{1-0.5\xi}$$

151<0.5

$$+\frac{z}{1-z}-\frac{z}{1-2z}$$

$$\frac{f(z)}{f(z)} = -\left[\frac{1}{1} + \frac{1}{2}z^{1} + \frac{1}{3}z^{2} + \cdots\right] - \frac{f(z)}{f(z)}$$

$$+\left[\left(\frac{1}{2}\right) + \left(\frac{1}{2}\right)^{2}z^{2} + \left(\frac{1}{2}\right)^{3}z^{2} + \cdots\right] + \left(\frac{1}{2}\right)^{n+1}$$

$$\frac{f(z) = +\left[1^{0}z^{1} + 1^{1}z^{2} + 1^{2}z^{3} + \cdots\right] + |^{n+1}}{-\left[2^{0}z^{1} + 2^{1}z^{2} + 2^{2}z^{3} + \cdots\right] - 2^{n+1}}$$

18/72

|そ| 7 |

$$\frac{1}{1-858^{-1}} + \frac{0.5}{1-0.58^{-1}}$$

$$\frac{(2) = -\left[\left[\left(\frac{1}{2} \right)^{1} z^{9} + \left[\left(\frac{1}{2} \right)^{3} z^{-1} + \left(\frac{1}{2} \right)^{3} z^{-2} + \cdots \right] - \left[\frac{n+1}{2} \right]^{n+1}}{+ \left[\left(\frac{1}{2} \right)^{1} z^{9} + \left(\frac{1}{2} \right)^{3} z^{-1} + \left(\frac{1}{2} \right)^{3} z^{-2} + \cdots \right] + \left[\frac{1}{2} \right]^{n+1}}$$

$$\frac{1}{2}$$

$$\frac{-1}{(2-1)(2-2)} \longrightarrow 2 \frac{-05\xi^2}{(2-1)(2-0.5)}$$

$$-\frac{1}{1-z}+\frac{0.5}{1-0.5z}$$

$$f(z) = -\left[1' + 1^{2}z' + 1^{3}z^{2} + \cdots\right] \qquad -|f|$$

$$+\left[\left(\frac{1}{2}\right) + \left(\frac{1}{2}\right)^{2}z' + \left(\frac{1}{2}\right)^{3}z^{2} + \cdots\right] \qquad +\left(\frac{1}{2}\right)^{n+1}$$

$$X (Z) = -\left[\left(\frac{1}{1}\right)^{-1} + \left(\frac{1}{1}\right)^{-2} z^{1} + \left(\frac{1}{1}\right)^{-3} z^{2} + \cdots \right] + 2^{n-1} + 2^{n-1}$$

$$\frac{\xi^{-1}}{1-\xi^{-1}} - \frac{\xi^{-1}}{1-2\xi^{-1}}$$

$\lambda = \left(\frac{1}{2}\right)^{-1}$ $\left(\frac{1}{2}\right) = \lambda^{-1}$

$$f(z) = + \left[\left(\frac{1}{1} \right)_{0}^{2} z_{1} + \left(\frac{1}{1} \right)_{-1}^{2} z_{-2} + \left(\frac{1}{1} \right)_{-2}^{2} z_{-3} + \cdots \right] - \left(\frac{2}{1} \right)_{4l+1}$$

$$- \left[\left(\frac{1}{1} \right)_{0}^{2} z_{1} + \left(\frac{1}{1} \right)_{-1}^{2} z_{-2} + \left(\frac{1}{1} \right)_{-2}^{2} z_{-3} + \cdots \right] - \left(\frac{1}{1} \right)_{4l+1}$$

151<0.5

$$\frac{f(z)}{-[2^{n}z^{1}+1^{1}z^{2}+1^{2}z^{3}+\cdots]} + |^{n+1}$$

$$-[2^{n}z^{1}+2^{1}z^{2}+2^{2}z^{3}+\cdots] - 2^{n+1}$$

$$N = -1 \qquad -5 \qquad -3$$

$$-\left[\left(\frac{1}{1}\right)_{0} \bar{s}_{1} + \left(\frac{1}{1}\right)_{-1} \bar{s}_{2} + \left(\frac{1}{1}\right)_{-2} \bar{s}_{3} + \cdots \right] \qquad + \left[\frac{1}{4+1} \right]_{0} \bar{s}_{1} + \left(\frac{1}{1}\right)_{-2} \bar{s}_{2} + \left(\frac{1}{1}\right)_{-3} \bar{s}_{3} + \cdots \right] \qquad + \left[\frac{1}{4+1} \right]_{0} \bar{s}_{1} + \left[\frac{1}{4} \right]_{0} \bar{s}_{2} + \left[\frac{1}{4} \right]_{0} \bar{s}_{3} + \cdots$$

$$\frac{1}{|-\epsilon^{-1}|} + \frac{0.5}{|-0.5\epsilon^{-1}|}$$

$$f(z) = -\left[\left(\frac{1}{1}\right)^{3} z^{0} + \left(\frac{1}{1}\right)^{2} z^{-1} + \left(\frac{1}{1}\right)^{-3} z^{-2} + \cdots \right] - |^{n-1} + \left[2^{-1} z^{0} + 2^{-2} z^{-1} + 2^{-3} z^{-2} + \cdots \right] + 2^{n-1}$$

151<

$$-\frac{1}{1-\xi}+\frac{0.5}{1-0.5\xi}$$

$$\frac{f(z) = -\left[|+|\frac{1}{2}|^2 z' + |\frac{1}{2}|^2 z^2 + \cdots\right]}{+\left[\left(\frac{1}{2}\right) + \left(\frac{1}{2}\right)^2 z' + \left(\frac{1}{2}\right)^3 z^2 + \cdots\right]}$$

$$(n \ge 0)$$

$$\frac{f(z)}{f(z)} = + \left[1^{o}z^{1} + 1^{1}z^{2} + 1^{2}z^{3} + \cdots \right]$$

$$\Delta_n = \pm |^{n-1} - 2^{n-1} \quad (n \geqslant 1)$$

$$(n = -1^{n-1} + 2^{n-1})$$

$$\frac{(2) = + \left[\left(\frac{1}{1}\right)^{-1} z^{1} + \left(\frac{1}{1}\right)^{-2} z^{2} + \left(\frac{1}{1}\right)^{-2} z^{3} + \cdots \right]}{- \left[\left(\frac{1}{2}\right)^{0} z^{1} + \left(\frac{1}{2}\right)^{-2} z^{3} + \left(\frac{1}{2}\right)^{-2} z^{3} + \cdots \right]}$$

$$Q_n = \uparrow \left(\frac{1}{2}\right)^{n+1} - \left(\frac{1}{2}\right)^{n+1} \quad (n < 0)$$

18172

$$\frac{\xi^{-1}}{1-\xi^{-1}}-\frac{\xi^{-1}}{1-2\xi^{-1}}$$

1817 1

$$\frac{1}{1-8^{-1}} + \frac{0.5}{1-0.5^{-1}}$$

$$f(z) = + \left[\left(\frac{1}{1} \right)^{0} z^{1} + \left(\frac{1}{1} \right)^{-1} z^{-2} + \left(\frac{1}{1} \right)^{-2} z^{-3} + \cdots \right]$$

$$- \left[\left(\frac{1}{1} \right)^{0} z^{1} + \left(\frac{1}{1} \right)^{-1} z^{-2} + \left(\frac{1}{1} \right)^{-2} z^{-3} + \cdots \right]$$

$$a_n = \frac{1}{2} \left(\frac{n+1}{2} - \left(\frac{1}{2} \right)^{n+1} \right) \quad (n < 0)$$

$$f(z) = -\left[\left(\frac{1}{1}\right)^{-1} z^{0} + \left(\frac{1}{1}\right)^{-2} \overline{z}^{-1} + \left(\frac{1}{1}\right)^{-3} \overline{z}^{-2} + \cdots \right] + \left[2^{-1} z^{0} + 2^{-2} z^{-1} + 2^{-3} \overline{z}^{-2} + \cdots \right]$$


$$\frac{-\left[2_{0}\xi_{1}+1_{1}\xi_{2}+1_{2}\xi_{3}+...\right]}{-\left[2_{0}\xi_{1}+1_{2}\xi_{2}+1_{2}\xi_{3}+...\right]}$$

$$\alpha_n = + |^{n-1} - 2^{n-1} \qquad (n \geqslant 1)$$

$$\frac{1}{\left(\frac{1}{2}\right)^{1}} \overline{z}^{0} + \left(\frac{1}{2}\right)^{2} \overline{z}^{-1} + \left(\frac{1}{2}\right)^{3} \overline{z}^{-2} + \cdots \right]$$

$$\Delta_n = -|^{n+1} \quad t(\frac{1}{2})^{n+1} \quad (n > 0)$$


	R0(€)	f(E)	\iff	a n	RNG(n)	
	2 < p				n≥ o	
I	R0((₹ ¹)	f(E)	←	- a n	RNG (n)	
_	1 2 1 > 1				n < 0	
	R0(€)	ƒ(そ)		Q n	RNG(n)	
	2 < p				n≥ o	
	P 0 6 (7-1)	f (7-1)		0	71166)	
Œ	RO((₹¹)	7(6)		OL-n	RNG(-n)	
	1717 P				n < 1	
	R0(€)	f(Z)	\iff	Оn	RNG(n)	
	131 < p				n> o	
$\qquad \qquad \blacksquare \qquad \qquad$	R0(₹)	f(z-')	\longleftrightarrow	<u>-</u> α-n	« RNG(n) »	I+I
_	} < p				n≯I	
	R0(€)	f(Z)	\longleftrightarrow	Дn	RNG(n)	
	2 < p	, ,			n> o	
$\overline{\mathbb{W}}$	R0((Z)	X (E)	\longleftrightarrow	a-n	RNG (-n)	
	} < p	, , ,			n < 1	
	161 - 1					





(\mathbb{I})	R0(€)	f(z')	$\langle \rightarrow \rangle$	<u>- Д-</u> п	« RNG(n) »	I+I
	} < p				n≯I	
	ROC(Z)	f(E)	\longleftrightarrow	Qп	RNG(n)	
	131 < p				n≥ o	
I						
						<u></u>
- 3	RO((z')	f(E)	\iff	— Q n	RNG (n)	
	171 > 				n < 0	
					n ≤ -1	
I						
	R0(€)	f(z')	\longleftrightarrow	- a-n	RNG (-n)	
	1 2 1 < p				n≽l	





	RO((₹)	f(E)	\iff	Q n	RNG(n)	
	1 3 1 < p				n≥ o	
(I)	ROC(E)	f(Z)	←	— a n	RNG (n)	
	1717P				n < 0	
			•			
				-	complement	
$\overline{\mathbb{Z}}$	R0(€)	X(E)	\iff	a-n	RNG(-n)	
	2 < p				n < 1	
				<u>-N</u>	-n	
					Symmetrical	











(W)	ROC(Z)	X (そ)	\longleftrightarrow	a-n	RNG(n)	
	2 < p				n < 1	

(P, 4) = (0.5, 1)(P,4)=(1,2) 2 (2-1)(2-2) (2-1)(2-0.5) f(2) (n > 0) (F) |z| < p X(Z) (n<1)f({}) <u>(-10)</u> 121 > 8 (n>1)X(Z)

	_	(P,4)=(1,2)	(7,4)=(0.5,1)
		(2-1)(2-2)	$2 \frac{(3-1)(2-0.5)}{(3-1)(2-0.5)}$
z < P	f(2)	-	$+ ^{n-1} - 2^{n-1} (n \ge 1)$
	Χ(₹)	•	$+ \frac{1}{n+1} - (\frac{7}{7})_{U+1} (U < 0)$
	f(2)		$-1^{n-1}+2^{n-1}(n<1)$
z > 	X(£)		$-\mid_{U+1}+\left(\frac{\tau}{T}\right)_{U+1}\left(U>0\right)$

$$f(z)$$
 $|z| < 0.5$ $|z| > 2$

Causal anticausal

$$-\left(|_{1} + |_{2} + |_{2} + |_{2} + |_{2} + \dots \right) + \left(\left(\frac{7}{1} \right) + \left(\frac{7}{1} \right)_{3} + \left(\frac{7}{1} \right)_{3} + \left(\frac{7}{1} \right)_{3} + \dots \right)$$

$$|\mathbf{z}| > 2 \qquad \frac{\xi^{-1}}{|-\xi^{-1}|} - \frac{\xi^{-1}}{|-2\xi^{-1}|} + |\frac{n+1}{n+1} - (\frac{1}{2})^{n+1}$$
 (n < 0)
$$(|^{0}\xi^{-1} + |^{1}\xi^{-2} + |^{2}\xi^{-3} + \cdots) - ((\frac{1}{2})^{0}\xi^{-1} + (\frac{1}{2})^{-1}\xi^{-2} + (\frac{1}{2})^{2}\xi^{-3} + \cdots)$$

$$(|^{0}\xi^{-1} + |^{-1}\xi^{-2} + |^{-2}\xi^{-3} + \cdots) - ((\frac{1}{2})^{0}\xi^{-1} + (\frac{1}{2})^{-1}\xi^{-2} + (\frac{1}{2})^{2}\xi^{-3} + \cdots)$$

$$|^{n-1} \qquad |^{n-2} \qquad |^{n-2} \qquad |^{n-2} \qquad |^{n-2} \qquad |^{n-2}$$

$$2-A \frac{-05\xi^{2}}{(2-1)(2-0.5)} = \left(-\frac{\xi}{(\xi-1)} + \frac{0.5\xi}{(\xi-0.5)}\right)$$

$$|\xi| < 0.5$$
 $f(a) = + \frac{\xi}{|-\xi|} - \frac{\xi}{|-\chi|\xi|} ||\eta^{-1}| - 2^{n-1}| (\eta > 1)$

$$+ (|_{0}\xi_{1} + |_{1}\xi_{2} + |_{3}\xi_{3} + \cdots) - (5_{0}\xi_{1} + 5_{1}\xi_{3} + 5_{2}\xi_{2} + \cdots)$$

$$\frac{1}{|\xi|} > \frac{1}{|\xi|} = \frac{1}{|\xi|^{2}} + \frac{1$$

$$(z)$$
 $|z| < 0.5$ $|z| > 2$

anticausal causal

$$\mathbb{D} - \mathbb{B} \quad \frac{-1}{(2-1)(2-2)} = \left(\frac{1}{\xi-1} - \frac{1}{\xi-2}\right)$$


$$|\xi| < |\chi| = -\frac{1}{1-\xi} + \frac{0.5}{1-0.5\xi} - |\chi| + 2^{n-1}$$

$$-(|\chi|^2 + |\chi|^2 + |\chi|^3 + |\chi|^3 + |\chi|^3 + |\chi|^4 + |\chi|^5 +$$

$$|\mathbf{z}| > 2 \qquad \times (2) = \frac{\mathbf{z}^{-1}}{|-\mathbf{z}^{-1}|} - \frac{\mathbf{z}^{-1}}{|-2\mathbf{z}^{-1}|} + \frac{\mathbf{z}^{-1}}{|$$

$$-0.52^{2} - (2-1)(2-0.5) - \left(-\frac{2}{(2-1)} + \frac{0.52}{(2-0.5)}\right)$$

$$|\xi| < 0.5 \qquad \times (\epsilon) \qquad = \qquad + \frac{\xi}{|-\xi|} - \frac{\xi}{|-\chi|^2} \qquad + |^{n+1} - (\frac{1}{2})^{n+1} \qquad (n < 0)$$





$$\frac{z}{|-\alpha z|} \Rightarrow \sum_{n=1}^{\infty} \alpha^{n-1} z^{n}$$


$$= -\sum_{k=0}^{\infty} \alpha^{k-1} z^{k}$$


$$\alpha + \alpha^{2} \xi^{1} + \alpha^{3} \xi^{2} + \alpha^{4} \xi^{3} + \cdots$$
 $\xi^{-1} + \alpha^{-1} \xi^{2} + \alpha^{2} \xi^{-3} + \alpha^{3} \xi^{-4} + \cdots$

$$z + \alpha z^{2} + \alpha^{2} z^{3} + \alpha^{3} z^{4} + \cdots$$
 $z^{4} + \alpha^{4} z^{5} + \alpha^{4} z^{5} + \cdots$











$$\frac{3}{2} \frac{-1}{(2-0.5)(2-2)} = \left(\frac{1}{2-0.5} - \frac{1}{2-2}\right)$$

$$|z| < 0.5 \qquad f(z) = -\frac{2}{1-2z} + \frac{0.5}{1-0.5z} - 2^{n+1} + (\frac{1}{2})^{n+1} + (\frac{1}{2})^{n+1}$$





$$X(z) = \frac{1}{1 - z^{-1}} - \frac{1}{1 - 2z^{-1}} \quad (|z| > 2)$$

$$a_n = 1^n + 2^n \quad (n \ge 0)$$

$$a_{n-1} = \frac{z^{-1}}{|-z^{-1}|} - \frac{z^{-1}}{|-zz^{-1}|} - (|z| > 2)$$

$$f(z) = (+1) \frac{1}{1-z} - \frac{1}{1-2z} \quad (|z| < 0.5)$$

$$a_n = |n| - 2^n \quad (n > 0)$$

$$a_{n-1} = \frac{\xi}{|-\xi|} - \frac{\xi}{|-\xi|} - \frac{\xi}{|-\xi|} \left(|\xi| < 0.5 \right)$$

$$f(z) = \frac{1}{1-z} - \frac{1}{1-2z} \qquad (|z| < 0.5)$$

$$a_n = |^n - 2^n \qquad (n \ge 0)$$

$$a_{n-1} = \frac{\xi}{|-\xi|} - \frac{\xi}{|-\lambda\xi|} \left(|z| < 0.5\right)$$

$$a_{n-1} = ||-\lambda\xi|| - 2^{n-1} \left(|n\rangle|\right)$$

$$\mathbf{Z}^{-1}\mathbf{f}(\mathbf{z}^{-1}) = \frac{\mathbf{z}^{-1}}{|-\mathbf{z}^{-1}|} - \frac{\mathbf{z}^{-1}}{|-\mathbf{z}^{-1}|} \left(|\mathbf{z}| > 2\right)$$

$$\mathbf{a}_{-\mathbf{n}-\mathbf{1}} = |\mathbf{n}+\mathbf{1}| - \left(\frac{1}{2}\right)^{\mathbf{n}+\mathbf{1}} \quad (\mathbf{n} < 0)$$

$$\mathbf{a}_{-(\mathbf{n}+\mathbf{1})}$$

$$X(\xi) = \frac{1}{1-\xi^{-1}} - \frac{1}{1-2\xi^{-1}} \quad (|z| > 2)$$

$$a_n = ||^n - 2^n \quad (n \ge 0)$$

$$a_{n-1} = \frac{z^{-1}}{|-z^{-1}|} - \frac{z^{-1}}{|-zz^{-1}|} (|z| > 2)$$

$$\mathbf{Z} \times (\mathbf{Z}^{-1}) = \frac{\mathbf{Z}}{|-\mathbf{Z}|} - \frac{\mathbf{Z}}{|-\mathbf{Z}|} \left(|\mathbf{Z}| < 0.5\right)$$

$$\mathbf{A}_{-\mathbf{n}-1} = |\mathbf{n}+1| - \left(\frac{1}{2}\right)^{\mathbf{n}+1} \qquad (\mathbf{n} < 0)$$

$$\mathbf{A}_{-(\mathbf{n}+1)}$$

Causal
$$f(z)$$
 $X(z)$ $|z| < |z| > 2$

$$\begin{array}{ccc}
\boxed{ -1} & = & \left(\frac{1}{\xi - 1} - \frac{1}{\xi - 2} \right)
\end{array}$$


$$f(z) = (-1)\frac{1}{1-z} + \frac{0.5}{1-0.5z}$$
 $(|z|<1)$

|2|< 2

|2|<|

$$a_n = -1^{n+1} + \left(\frac{1}{2}\right)^{n+1} \quad (n \ge 0)$$

$$\frac{-1}{(2-1)(2-2)} = \left(\frac{\xi-1}{\xi-1} - \frac{\xi-2}{\xi}\right)$$



$$X (z) = \frac{z^{-1}}{1-z^{-1}} - \frac{z^{-1}}{1-zz^{-1}} \quad (|z| > 2)$$

|2| > | |2| > 2

$$a_n = |^{n-1} - 2^{n-1} \quad (n \ge 1)$$

(n > 0)

Causal
$$f(z)$$
 $X(z)$ $|z| < 0.5$ $|z| > 1$

$$2-A \frac{-052^2}{(2-1)(2-0.5)} = \left(-\frac{2}{(2-1)} + \frac{0.52}{(2-0.5)}\right)$$

$$f(z) = (+1)\frac{z}{|-z|} - \frac{z}{|-2z|} (|z| < 0.5)$$

$$|^{n} - 2^{n} (n \ge 0)$$

$$a_{n} = |^{n-1} - 2^{n-1} (n \ge 1)$$

$$2-B\frac{-05z^2}{(z-1)(z-0.5)} = \left(-\frac{z}{(z-1)} + \frac{0.5z}{(z-0.5)}\right)$$

$$X(z) = -\frac{1}{1-z^{-1}} + \frac{o.s}{1-o.sz^{-1}} (|z| > 1)$$

$$a_n = -|^{n+1} + (\frac{1}{2})^{n+1} \qquad (n \ge 0)$$

Anti-causal
$$f(z)$$
 $X(z)$ $|z| > 2$ $|z| < |z|$

 a_n

$$f(z) = \frac{z^{-1}}{|-z^{-1}|} - \frac{z^{-1}}{|-zz^{-1}|} \quad (|z|>2)$$

$$|^{n-1} - 2^{n-1} \quad (n \ge 1)$$

(n > 0)


|2| > 2

$$a_n = |^{n+1} - (\frac{1}{2})^{n+1} \quad (n < 0)$$

$$\frac{-1}{(2-1)(2-2)} = \left(\frac{1}{\xi-1} - \frac{1}{\xi-2}\right)$$

=

|2|>|



$$X = \frac{1}{1-\xi} + \frac{0.5}{1-0.5 z} (|z| < 1)$$

$$-|^{n+1} + (\frac{1}{2})^{n+1} (n \ge 0)$$

$$-|^{n-1} + 2^{n-1} (n < 1)$$

Anti-Causal
$$f(z)$$
 $\chi(z)$ $|z| > 1$

$$2-A \frac{-05\xi^2}{(2-1)(2-0.5)} = \left(-\frac{\xi}{(\xi-1)} + \frac{0.5\xi}{(\xi-0.5)}\right)$$

 a_n

$$f(z) = (-1)\frac{1}{1-z^{-1}} + \frac{0.5}{1-0.5z^{-1}} (|z| > 1)$$

$$- |^{n+1} + (\frac{1}{2})^{n+1}$$

$$= -|^{n-1} + 2^{n-1}$$

$$|-B| \frac{-0.5 z^2}{(2-1)(2-0.5)} = \left(-\frac{z}{(z-1)} + \frac{0.5 z}{(z-0.5)}\right)$$

$$\frac{\xi}{|-\xi|} - \frac{\xi}{|-2\xi|} \left(|\xi| < 1\right)$$

(n > 0)

(n<1)

$$a_n$$

$$\frac{(3-1)(3-2)}{-1} = f(3)$$

anticausal

$$\frac{-1}{(2-1)(2-2)} = \left(\frac{2-1}{2-1} - \frac{1}{2-2}\right) = -\frac{1}{1-2} + \frac{0.5}{1-0.52}$$

$$f(\overline{z}) = -\frac{(1)}{1 - (\overline{z})} + \frac{(\frac{1}{a})}{1 - (\frac{\overline{z}}{2})}$$

$$= -\sum_{n=0}^{\infty} (1)^{n+i} (\overline{z})^n + \sum_{n=0}^{\infty} (\frac{1}{2})^{n+i} (\overline{z})^n$$

$$= -\sum_{n=0}^{\infty} (1)^{n+i} \overline{z}^n + \sum_{n=0}^{\infty} (\frac{1}{2})^{n+i} \overline{z}^n$$

$$\alpha_n =$$

$$a_n = -1^{n+i} + \left(\frac{1}{2}\right)^{n+i}$$

$$\frac{-1}{(2-1)(2-2)} = \left(\frac{1}{\xi-1} - \frac{1}{\xi-2}\right) = -\frac{1}{1-\xi} + \frac{0.5}{1-0.5\xi}$$

$$\frac{1 - (\frac{\pi}{5})}{1 - (\frac{\pi}{5})} = \frac{\frac{1 - (\frac{\pi}{5})}{1 - (\frac{\pi}{5})}}{\frac{\pi}{5}} = \frac{\frac{1 - (\frac{\pi}{5})}{1 - (\frac{\pi}{5}$$

$$=\sum_{n=-1}^{-\infty} \left(1\right)^{n+1} \, \xi^n - \sum_{n=-1}^{-\infty} \left(\frac{1}{2}\right)^{n+1} \, \xi^n$$

$$a_n$$

$$\frac{-1}{(2-1)(2-2)} = \begin{bmatrix} \chi(2) \\ \text{anticausal} \end{bmatrix} = \begin{bmatrix} \xi & 0.5 \\ \text{anticausal} \end{bmatrix}$$

$$|z| < 0.5$$
 $|z| > 2$ anticausal causal

$$\frac{-1}{(2-1)(2-2)} = \left(\frac{1}{\xi-1} - \frac{1}{\xi-2}\right) = -\frac{1}{1-\xi} + \frac{0.5}{1-0.5\xi}$$

$$= -\sum_{n=0}^{-\infty} (1)^{n-1} \xi^{-n} + \sum_{n=0}^{-\infty} (2)^{n-1} \xi^{-n}$$

$$(n \le 0)$$
 $a_n = -(1)^{n-1} + 2^{n-1}$

$$\frac{-1}{(2-1)(2-2)} = \left(\frac{1}{\xi-1} - \frac{1}{\xi-2}\right) = -\frac{1}{1-\xi} + \frac{0.5}{1-0.5\xi}$$

$$(n > 0)$$
 $a_n = (1)^{n-1} - (2)^{n-1}$

$$2-\triangle \frac{-0.5 z^2}{(2-1)(2-0.5)} = f(2) \qquad |z| < 0.5 \qquad |z| > 2$$

$$causal \qquad anticausal$$

$$\frac{-0.5 \, z^2}{(2-1)(2-0.5)} = \left(-\frac{z}{(2-1)} + \frac{0.5z}{(z-0.5)}\right) = \left(\frac{z}{(1-z)} - \frac{z}{(1-2z)}\right)$$


$$\frac{f(z)}{f(z)} = \frac{\frac{(z)}{1-(z)}}{1-(z)} = \frac{(z)}{1-(2z)} \neq$$

$$= \sum_{n=0}^{\infty} (1)^n (z)^{n+1} - \sum_{n=0}^{\infty} (2)^n (z)^{n+1}$$

$$= \sum_{n=1}^{\infty} (1)^{n-1} z^n - \sum_{n=1}^{\infty} (2)^{n-1} z^n$$

$$a_n = 1^{n-1} - (2)^{n-1}$$

$$\frac{-0.5 \, \xi^2}{(2-1)(2-0.5)} = \left(-\frac{\xi}{(2-1)} + \frac{0.5\xi}{(2-0.5)} \right) = \left(-\frac{\xi}{(1-\xi)} - \frac{\xi}{(1-2\xi)} \right)$$



$$\frac{1}{1-\left(\frac{1}{2}\right)} = -\frac{\left(\frac{1}{2}\right)}{1-\left(\frac{1}{2}\right)} + \frac{\left(\frac{1}{2}\right)}{1-\left(\frac{1}{2}\right)}$$

$$= -\sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^{n+1} \left(\frac{1}{2}\right)^n + \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^{n+1} \left(\frac{1}{2}\right)^n$$

$$= -\sum_{n=0}^{\infty} (1)^{n+1} \xi^{-n} + \sum_{n=0}^{\infty} (\frac{1}{2})^{n+1} \xi^{-n}$$

$$= -\sum_{n=0}^{\infty} (1)^{n-1} \xi^{n} + \sum_{n=0}^{\infty} (2)^{n-1} \xi^{n}$$



$$(n \leq o)$$

$$a_n = -|^{n-1} + (2)^{n-1}$$

$$2 - \beta \qquad \frac{-0.5 \, \xi^2}{(2-1)(2-0.5)} = \chi(2)$$

$$|z| < 0.5$$
 $|z| > 2$

anticausal causal

$$\frac{-0.5 \, \xi^2}{(2-1)(2-0.5)} = \left(-\frac{\xi}{(2-1)} + \frac{0.5\xi}{(2-0.5)}\right) = \left(\frac{\xi}{(1-\xi)} - \frac{\xi}{(1-2\xi)}\right)$$

$$\frac{-0.5 \, z^2}{(2-1)(2-0.5)} = \left(-\frac{z}{(2-1)} + \frac{0.5z}{(z-0.5)}\right) = \left(\frac{z}{(1-z)} - \frac{z}{(1-2z)}\right)$$

$$X(\xi) = -\frac{\left(\frac{1}{2}\right)}{\left|-\left(\frac{1}{2}\right)\right|} + \frac{\left(\frac{1}{2}\right)}{\left|-\left(\frac{1}{2}\right)\right|}$$

$$= -\sum_{n=0}^{\infty} \left(1\right)^{n+1} \left(\frac{1}{2}\right)^n + \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^{n+1} \left(\frac{1}{2}\right)^n$$

$$= -\sum_{n=0}^{\infty} \left(1\right)^{n+1} \xi^{-n} + \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^{n+1} \xi^{-n}$$

$$(n \geqslant 0) \qquad a_n = -(1)^{n+1} + (\frac{1}{2})^{n+1}$$



