Laurent Series and z-Transform

Geometric Series Double Pole Examples B

20180222

Copyright (c) 2016 - 2018 Young W. Lim.

Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.2 or any later version published by the Free Software Foundation; with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts. A copy of the license is included in the section entitled "GNU Free Documentation License".

2 formulas of z

$$\frac{-1}{(2-1)(2-2)} = \left(\frac{\xi-1}{\xi-1} - \frac{\xi-2}{\xi-2}\right)$$



$$\frac{-052^2}{(2-1)(2-0.5)} = \left(-\frac{2}{(2-1)} + \frac{0.52}{(2-0.5)}\right)$$

$$\frac{-1}{(2^{-1})(2^{-2})} = \left(\frac{1}{\xi - 1} - \frac{1}{\xi - 2}\right)$$

$$= \left(\frac{1}{\xi - 0.5} - \frac{1}{\xi - 2}\right)$$

$$= \left(\frac{1}{\xi^{-1} - 1} - \frac{1}{\xi^{-2}}\right)$$

$$= \left(\frac{\xi}{1 - \xi} - \frac{\xi}{1 - 2\xi}\right)$$

$$= \left(\frac{-\xi}{2 - 1} + \frac{0.5\xi}{2 - 0.5}\right)$$

$$= \xi \left(\frac{-0.5\xi}{(\xi - 1)(\xi - 0.5)}\right)$$

$$= \frac{-0.5\xi^{2}}{(\xi - 1)(\xi - 0.5)}$$

$$\frac{-0.52^{2}}{(2-1)(2-0.5)} = \left(-\frac{2}{(2-1)} + \frac{0.52}{(2-0.5)}\right)$$

$$\frac{(5-1)(5-5)}{-1} = \left(\frac{\xi-1}{1} - \frac{\xi-2}{1}\right)$$

$$\frac{f(z)}{|z| > 2} \qquad f(z) = \frac{z^{-1}}{|-z^{-1}|} - \frac{z^{-1}}{|-z|^{2}} \qquad + |^{n+1} - (\frac{1}{z})^{n+1}| \qquad (n < 0)$$

$$\frac{\chi(\xi)}{|\xi| > 2} \qquad \chi(\xi) = \frac{\xi^{-1}}{|-\xi^{-1}|} - \frac{\xi^{-1}}{|-2\xi^{-1}|} \qquad + \frac{|-1|}{|-2|^{n-1}} - \frac{|-1|}{|-2|}$$

$$\frac{-05z^2}{(z-1)(z-0.5)} = \left(-\frac{z}{(z-1)} + \frac{0.5z}{(z-0.5)}\right)$$

(2)-(A)
$$|\xi| < 05$$
 $f(\xi) = + \frac{\xi}{|-\xi|} - \frac{\xi}{|-\chi\xi|} |n-1| - 2^{n-1} (n > 1)$

$$\frac{f(z)}{|z|} |z| > | f(z) = -\frac{1}{1-z^{-1}} + \frac{0.5}{1-0.5z^{-1}} - | \frac{1}{n-1} + 2^{n-1} | (n < 1)$$

$$\frac{\chi(\xi)}{|\xi| > 1} \qquad \chi(\xi) = -\frac{1}{|-\xi^{-1}|} + \frac{0.5}{|-0.5\xi^{-1}|} - \frac{1}{|-1|} + \frac{1}{2} \frac{1}{|-1|} (1) > 0$$

(P,4)=(0.5, 1) (P, G)=(1, 2) -0522 (2-1)(2-2) (2-1)(2-0.5) A 121 < p f(2) $+ | \frac{1}{n+1} - (\frac{2}{1})^{n+1} (n < 0)$ 121 > 8 (n<|) B |z| < p X(Z) $+ | ^{n-1} - 2^{n-1} (n > 1)$ 121 78

	_	(P, G) = (1, L)	(P,4)=(0.5, 1)
		(1) (2-1) (2-2)	2 <u>-052</u> (2-1)(2-0.5)
121 < 10	f(2)	$\frac{-1}{n+1} + \left(\frac{7}{17}\right)_{U+1} (U > 0)$	$+ ^{n-1} - 2^{n-1} (n \ge 1)$
2 < P	Χ(₹)		$+ _{u+1} - (\frac{7}{17})_{u+1} (u < 0)$
z > \f	f(2)	$+ \frac{1}{n+1} - (\frac{1}{2})^{n+1} $ (n<0)	$-1^{n-1}+2^{n-1}(n<1)$
	X(2)	$+ _{u-1} - 5_{u-1} (u)$	$\frac{- \left(\frac{1}{2}\right)^{n+1} + \left(\frac{1}{2}\right)^{n+1} (n \geqslant 0)}{n+1}$

(P, 4) = (0.5, 1)(7,4)=(1,2) $2 \frac{-052^2}{(2-1)(2-0.5)}$ (-| (2-1)(2-2) 121 < P f(2) causal (n>1) causal (n>0) anticausal (n<1) |2| > B f(2) anticausal (n<0) anticausal (n<1) X(z)121 < P anticausal (N<0) causal (170) causal (n>1) X(z)121 > 8

(P, 4) = (0.5, 1)(P, 4)=(1, 2) -0522 (2-1)(2-0.5) causal (n>1) 121 < p | f(2) causal (n>0) X(Z) |2| < |2 anticausal (n<1) anticausal (n<0) anticausal (N<0) f(2) anticausal (M<1) 121 > B X(z)causal (931) causal (n>0) 121 > 8

$$\frac{-1}{(2-1)(2-2)} = 2 \frac{-05z^2}{(2-1)(2-0.5)}$$

$$\left(\frac{1}{\xi-1}-\frac{1}{\xi-2}\right)$$

$$\left(\frac{2}{(2-1)} + \frac{0.5 z}{(2-0.5)}\right)$$

$$\frac{2}{|-\xi|} + \frac{0.5}{|-0.5\xi|}$$

$$+\frac{z}{|-z|}$$

15/<1 |5/<1 |0.58)<1

[2]<0.5 |2]<1 |22]<1

$$\frac{\xi^{1}}{|-\xi^{1}|} - \frac{\xi^{1}}{|-2\xi^{1}|}$$

$$\frac{2^{-1}}{1-\epsilon^{-1}}+\frac{0.5}{1-0.5\epsilon^{-1}}$$

|と| 72 | ほりく| | 1287 | く |

|&|7 | |&1|<| |0.521|<|

$$\frac{-1}{(2-1)(2-2)} = 2 \frac{-05\xi^2}{(2-1)(2-0.5)}$$

$$\left(\frac{1}{\xi-1}-\frac{1}{\xi-2}\right)$$

$$\left(-\frac{2}{(2-1)}+\frac{0.5 z}{(2-0.5)}\right)$$

$$\frac{2}{1-\xi} + \frac{0.5}{1-0.5\xi} + \frac{\xi}{1-\xi} - \frac{\xi}{1-2\xi}$$

$$\frac{z^{1}}{|-z^{1}} - \frac{z^{1}}{|-2z|}$$

$$\frac{\xi^{-1}}{|-\xi^{-1}|} - \frac{\xi^{-1}}{|-2\xi^{-1}|} = \frac{|-\xi^{-1}|}{|-\xi^{-1}|} + \frac{0.5}{|-0.5\xi^{-1}|}$$

$$(n \ge 1)$$

$$X(2)$$
 causal $(n \ge 0)$

$$\left(\frac{1}{\xi-1}-\frac{1}{\xi-2}\right)$$

$$\left(-\frac{\xi}{(\xi-1)} + \frac{0.5 z}{(\xi-0.5)}\right)$$

[21<0.5] f(2) causal (n>1)

$$\frac{\xi^{-1}}{|-\xi^{-1}|} - \frac{\xi^{-1}}{|-2\xi^{-1}|}$$

$$\frac{z}{|-z^{-1}|} + \frac{0.5}{|-0.5|^{-1}}$$

$$X(2)$$
 causal $(n \ge 1)$

$$X(2)$$
 causal $(n \ge 0)$

$$(n \ge 0)$$
 $(n \ge 1)$ $(n \le 0)$ $(n \ge 1)$ $(n \le 0)$ $(n \ge 1)$ $(n \ge 0)$ $(n \ge 1)$ $(n \ge 0)$ $(n \ge 1)$ $(n \ge 0)$ $(n \ge 0)$

$$\frac{-1}{(2-1)(2-2)} = 2 \frac{-052^2}{(2-1)(2-0.5)}$$

151<1

$$-\frac{1}{1-\frac{2}{5}}+\frac{0.5}{1-0.5^{2}}$$

151<0.5

$$+\frac{z}{1-z}-\frac{z}{1-2z}$$

$$\frac{f(z) = -\left(\left|\frac{1}{2}\right| + \left|\frac{1}{2}\right|^2 z' + \left|\frac{1}{2}\right|^3 z^2 + \cdots\right) - \left|\frac{1}{2}\right|^{n+1}}{+\left(\left|\frac{1}{2}\right|\right)^2 z' + \left(\left|\frac{1}{2}\right|\right)^3 z^2 + \cdots\right) + \left(\left|\frac{1}{2}\right|\right)^{n+1}}$$

$$\frac{f(z) = +\left[1^{0}z^{1} + 1^{1}z^{2} + 1^{2}z^{3} + \cdots\right] + |^{n+1}}{-\left[2^{0}z^{1} + 2^{1}z^{2} + 2^{2}z^{3} + \cdots\right] - 2^{n+1}}$$

18/72

$$\frac{1}{1-858^{-1}} + \frac{0.5}{1-0.58^{-1}}$$

$$\frac{(2) = -\left[\left[\left(\frac{1}{2} \right)^{1} z^{9} + \left[\left(\frac{1}{2} \right)^{3} z^{-1} + \left(\frac{1}{2} \right)^{3} z^{-2} + \cdots \right] - \left[\frac{n+1}{2} \right]^{n+1}}{+ \left[\left(\frac{1}{2} \right)^{1} z^{9} + \left(\frac{1}{2} \right)^{3} z^{-1} + \left(\frac{1}{2} \right)^{3} z^{-2} + \cdots \right] + \left[\frac{1}{2} \right]^{n+1}}$$

$$\frac{1}{2}$$

$$\frac{-1}{(2-1)(2-2)} \longrightarrow 2 \frac{-05\xi^2}{(2-1)(2-0.5)}$$

$$-\frac{1}{1-\xi}+\frac{0.5}{1-0.5\xi}$$

121<0.5

$$\frac{f(z)}{f(z)} = -\left[|+|^2 \overline{z}^1 + |^3 \overline{z}^2 + \cdots\right] \qquad -|^{\frac{1}{100}}$$

$$+\left[\left(\frac{1}{2}\right) + \left(\frac{1}{2}\right)^2 \overline{z}^1 + \left(\frac{1}{2}\right)^3 \overline{z}^2 + \cdots\right] \qquad +\left(\frac{1}{2}\right)^{n+1}$$

$$\frac{f(z)}{-[2^{0}z^{1}+1^{1}z^{2}+1^{2}z^{3}+\cdots]} + |^{n+1}$$

$$-[2^{0}z^{1}+2^{1}z^{2}+2^{2}z^{3}+\cdots]-2^{n+1}$$

$$\begin{array}{c|c}
2 = \left(\frac{1}{2}\right)^{-1} & \times (2) = -\left[\left(\frac{1}{1}\right)^{-1} + \left(\frac{1}{1}\right)^{-2} z^{1} + \left(\frac{1}{1}\right)^{-3} z^{2} + \cdots\right] & -1 \\
+ \left[2^{-1} + 2^{-2} z^{1} + 2^{-3} z^{2} + \cdots\right] & +2^{n-1}
\end{array}$$

$$N = -1 - 5 - 3$$

$$-\left[\left(\frac{7}{1}\right)_{0}\xi_{1} + \left(\frac{7}{1}\right)_{2}\xi_{2} + \left(\frac{7}{1}\right)_{2}\xi_{3} + \cdots\right] - \left(\frac{7}{1}\right)_{M+1}$$

$$\times (5) = +\left[\left(\frac{1}{1}\right)_{1}\xi_{1} + \left(\frac{1}{1}\right)_{2}\xi_{2} + \left(\frac{1}{1}\right)_{2}\xi_{3} + \cdots\right] - \left(\frac{7}{1}\right)_{M+1}$$

18172

$$\frac{\xi^{-1}}{1-\xi^{-1}}-\frac{\xi^{-1}}{1-2\xi^{-1}}$$

$$2 = \left(\frac{1}{2}\right)^{-1} \quad f(z)$$

$$\left(\frac{1}{2}\right) = 2^{-1}$$

$$f(z) = + \left[\left(\frac{1}{1} \right)^{0} z^{-1} + \left(\frac{1}{1} \right)^{-1} z^{-2} + \left(\frac{1}{1} \right)^{-2} z^{-3} + \cdots \right] + \left[\frac{1}{1} \right]^{-1} z^{-2} + \left(\frac{1}{2} \right)^{-2} z^{-3} + \cdots \right] - \left(\frac{1}{2} \right)^{-1} z^{-1} + \left(\frac{1}{2} \right)^{-1} z^{-2} z^{-3} + \cdots \right] - \left(\frac{1}{2} \right)^{-1} z^{-1} + \cdots + \left[\frac{1}{2} \right]^{-1} z^{-2} z^{-3} + \cdots$$

$$\frac{-\frac{1}{1-\epsilon^{-1}}+\frac{0.5}{1-0.5\epsilon^{-1}}}{-\frac{1}{1-\epsilon^{-1}}+\frac{1}{1-0.5\epsilon^{-1}}}$$

+ [2 2 2 + 2 2 1 + 2 2 2 1 + ...] + 2 1 -1

151<

$$-\frac{1}{1-\xi}+\frac{0.5}{1-0.5\xi}$$

$$+\frac{\xi}{|-\xi|}-\frac{\xi}{|-2\xi|}$$

$$\frac{f(z) = -\left[|1+1|^2 z' + |3|^2 z' + \cdots\right]}{+\left[\left(\frac{1}{2}\right) + \left(\frac{1}{2}\right)^2 z' + \left(\frac{1}{2}\right)^3 z' + \cdots\right]}$$

$$(n \ge 0)^{n+1} + (\frac{1}{2})^{n+1} \qquad (n \ge 0)$$

$$\frac{f(z)}{f(z)} = + \left[1_0 f_1 + 1_1 f_2 + 1_2 f_3 + \cdots \right]$$

$$\Delta_n = \pm |^{n-1} - 2^{n-1} \quad (n \geqslant 1)$$

$$Q_n = -1^{n-1} + 2^{n-1} \quad (n < [)$$

$$\frac{(2) = + \left[\left(\frac{1}{1}\right)^{-1} z^{1} + \left(\frac{1}{1}\right)^{-2} z^{2} + \left(\frac{1}{1}\right)^{-2} z^{3} + \cdots \right]}{- \left[\left(\frac{1}{2}\right)^{0} z^{1} + \left(\frac{1}{2}\right)^{\frac{1}{2}} z^{2} + \left(\frac{1}{2}\right)^{\frac{2}{2}} z^{3} + \cdots \right]}$$

$$a_n = \pm 1^{n+1} - \left(\frac{1}{2}\right)^{n+1} \quad (n < 0)$$

18172

$$\frac{\xi^{-1}}{1-\xi^{-1}}-\frac{\xi^{-1}}{1-2\xi^{-1}}$$

1817 1

$$\frac{1}{1-2^{-1}}+\frac{0.5}{1-0.5^{-1}}$$

$$f(z) = + \left[\left(\frac{1}{1} \right)^{0} z^{-1} + \left(\frac{1}{1} \right)^{-1} z^{-2} + \left(\frac{1}{1} \right)^{-2} z^{-3} + \cdots \right] - \left[\left(\frac{1}{2} \right)^{0} z^{-4} + \left(\frac{1}{2} \right)^{-1} z^{-2} + \left(\frac{1}{2} \right)^{-2} z^{-3} + \cdots \right]$$

$$a_n = + \left| \frac{n+1}{2} - \left(\frac{1}{2} \right)^{n+1} \right| \quad (n < 0)$$

$$\frac{f(z)}{+\left[2^{-1}z^{0}+2^{2}z^{-1}+2^{-3}z^{-2}+\cdots\right]} + \left[2^{-1}z^{0}+2^{2}z^{-1}+2^{-3}z^{-2}+\cdots\right]$$

$$-\left[J_{0}\xi_{1} + J_{1}\xi_{2} + J_{2}\xi_{3} + \cdots \right]$$

$$\times (\xi) = +\left[\left[\int_{0}^{\xi_{1}} \xi_{1} + \left[\int_{1}^{\xi_{2}} \xi_{2} + \left[\int_{1}^{\xi_{2}} \xi_{3} + \cdots \right] \right] \right]$$

$$\alpha_n = + |^{n-1} - 2^{n-1} \qquad (n \geqslant 1)$$

$$\frac{(5)}{(5)} = -\left[\left[\left(\frac{1}{2} \right)_{1}^{3} \xi^{0} + \left(\frac{1}{2} \right)_{2}^{3} \xi^{-1} + \left(\frac{1}{2} \right)_{3}^{3} \xi^{-2} + \cdots \right] + \left[\left(\frac{1}{2} \right)_{1}^{3} \xi^{0} + \left(\frac{1}{2} \right)_{2}^{3} \xi^{-1} + \left(\frac{1}{2} \right)_{3}^{3} \xi^{-2} + \cdots \right]$$

$$\Delta_n = -|^{n+1} \quad t(\frac{1}{2})^{n+1} \quad (n > 0)$$

 $(p,q) = (1,2) \qquad (p,q) = (0.5, 1)$ $-\frac{-1}{(2-1)(2-2)} \qquad 2 \qquad \frac{-0.5 \, 2^2}{(2-1)(2-0.5)}$ |z| < p $f(z) \qquad + \frac{n+1}{n+1} - \left(\frac{1}{2}\right)^{n+1} (n < 0) \qquad - \frac{n-1}{n+1} + 2^{n-1} (n < 1)$ |z| > 9

$$(p,q) = (1,2) \qquad (p,q) = (0.5, 1)$$

$$-\frac{-1}{(2-1)(2-2)} \qquad 2 \qquad \frac{-0.5 \, 2}{(2-1)(2-0.5)}$$

$$|z|
$$|z| > q \qquad \qquad \times (z) \qquad +|^{n-1} - 2^{n-1} \quad (n\geqslant|) \qquad -|^{n+1} + \left(\frac{1}{2}\right)^{n+1} \quad (n\geqslant0)$$$$

$$2^{-n+1} = \left(\frac{1}{2}\right)^{n} \cdot 2 = \left(\frac{1}{2}\right)^{n-1} \qquad \left(\frac{1}{2}\right)^{-n-1} = 2^{n} \cdot 2 = 2^{n+1}$$

$$\left(\frac{1}{2}\right)^{-n+1} = 2^{n} \cdot \frac{1}{2} = 2^{n-1} \qquad 2^{-n-1} = \left(\frac{1}{2}\right)^{n} \cdot \frac{1}{2} = \left(\frac{1}{2}\right)^{n+1}$$

	_		
		(2-1)(2-2)	$2 \frac{-0.57}{(2-1)(2-0.5)}$
z < P	f(2)	$\frac{-1}{n+1} + \left(\frac{7}{17}\right)_{U+1} (U > 0)$	
iei × r		-n	4
121 > 6	f(2)	$+ _{u+1} - (\frac{7}{1})_{u+1} $ (u<0)	$-1^{n-1}+2^{n-1}(n<1)$
 			
		(P, G) = (1, L)	(P, 4) = (0.5, 1)

$$|z| < P$$

$$|z| < P$$

$$|z| > \theta$$


$$(P,G)=(1,2)$$
 $(P,G)=(0.5,1)$

		_		
			(2-1)(2-2)	$2 \frac{-052}{(2-1)(2-0.5)}$
	z < P	f(2)	$\frac{-1}{u+1} + \left(\frac{z}{T}\right)_{U+I} (U > 0)$	$+ ^{n-1} - 2^{n-1} (n \ge 1)$
'	161 , 1		-n,	-1
		f(2)	$+ \mid_{u+1} - \left(\frac{\tau}{T}\right)_{u+1} (u < 0)$	$-1^{n-1}+2^{n-1}(n<1)$
	ट । > फ		-n,	1

$$(P, G) = (1, 2)$$

₹-<u>]</u>

		(2-1)(2-2)	2 -052 (2-1)(2-0.5)
z < P		- K	-1
lei · r	Χ(₹)	$- ^{n-1}+2^{n-1}$ (n<)	$+ \frac{1}{n+1} - (\frac{7}{1})_{U+1} (U < 0)$
		-n,	-1
E > %	X(2)	$+ \frac{1}{n-1} - 2^{n-1} - (n > 1)$	$\frac{- \mathbf{u}+ +(\frac{2}{1})_{\mathbf{U}+1}}{(\mathbf{U}>0)}$



$$f(z)$$
 $|z| < 0.5$ $|z| > 2$

Causal anticausal

$$|\xi| < 0.5$$
 $f(\xi) = -\frac{2}{1-2\xi} + \frac{0.5}{1-0.5\xi} -2^{n+1} + (\frac{1}{2})^{n+1}$ $(N > 0)$

$$\frac{1-\alpha\xi}{\alpha} \frac{\xi^{-1}}{\alpha^{4}\xi^{4}-1} \frac{-\left(2+2^{3}\xi+2^{3}\xi^{2}+\cdots\right)+\left(\frac{1}{2}+\frac{1}{2}\xi+\frac{1}{2}\xi+\frac{1}{2}\xi+\frac{1}{2}\xi^{2}\xi^{2}+\cdots\right)}{n=0 \quad n=1 \quad n=2}$$

$$|\xi| > 2 \qquad f(\xi) = \frac{\xi^{-1}}{|-0.5\xi^{-1}|} - \frac{\xi^{-1}}{|-2\xi^{-1}|} + 2^{n+1} - (\frac{1}{2})^{n+1} \qquad (n < 0)$$

$$(\xi^{-1} + (\frac{1}{2})^{1} \xi^{-2} + (\frac{1}{2})^{2} \xi^{-3} + \cdots) - (\xi^{-1} + 2 \xi^{-2} + 2^{2} \xi^{-3} + \cdots)$$

$$\left(2^{5}\xi^{-1} + 2^{-1}\xi^{-2} + 2^{-2}\xi^{-5} + \cdots\right) - \left(\left(\frac{1}{2}\right)^{6}\xi^{-1} + \left(\frac{1}{2}\right)^{-1}\xi^{-1} + \left(\frac{1}{2}\right)^{2}\xi^{-3} + \cdots\right)$$

$$N = -1 \qquad N = -2 \qquad N = -3$$

$$N = -1 \qquad N = -2 \qquad N = -3$$

$$-A = \frac{3}{2} \frac{-2^2}{(2-2)(2-0.5)} = \left(\frac{0.52}{(2-0.5)} - \frac{22}{(2-2)}\right)$$

$$|\xi| < 0.5$$
 $f(\xi) = -\frac{\xi}{1-2\xi} + \frac{\xi}{1-0.5\xi} -2^{n-1} + (\frac{1}{2})^{n-1} \quad (n > 1)$

$$|\xi| > 2$$
 $f(\xi) = \frac{0.5}{|-a.5\xi^{-1}|} - \frac{2}{|-2\xi^{-1}|} + 2^{n_1} - (\frac{1}{2})^{n_{-1}}$ $(n < 1)$

$$\frac{\left(\left(\frac{1}{2}\right) + \left(\frac{1}{2}\right)^{2} \xi^{-1} + \left(\frac{1}{2}\right)^{3} \xi^{-2} + \cdots\right) + \left(\left(\frac{1}{2}\right)^{-1} + \left(\frac{1}{2}\right)^{-2} \xi^{-1} + \left(\frac{1}{2}\right)^{-3} \xi^{-2} + \cdots\right)}{\left(2 + 2^{2} \xi^{-1} + 2^{-3} \xi^{-2} + \cdots\right) + \left(2 + 2^{2} \xi^{-1} + 2^{3} \xi^{-2} + \cdots\right)}$$

$$(z)$$
 $|z| < 0.5$ $|z| > 2$ anticausal causal

$$|\xi| < 0.5$$
 $\chi(\xi) = -\frac{2}{1-2\xi} + \frac{0.5}{1-0.5\xi} -(\frac{1}{2})^{n-1} + 2^{n-1}$ $(n < 1)$

$$-\left(2^{i}\xi^{0}+2^{2}\xi^{1}+2^{3}\xi^{2}+\cdots\right)+\left(\frac{1}{2}\xi^{0}+\left(\frac{1}{2}\xi^{0}+\left(\frac{1}{2}\xi^{0}+2^{3}\xi^{1}+\cdots\right)\right)\right)$$
$$-\left(\left(\frac{1}{2}\xi^{0}+\left(\frac{1}{2}\xi^{0}+\left(\frac{1}{2}\xi^{0}+2^{3}\xi^{1}+\cdots\right)+\left(2^{-1}\xi^{0}+2^{-2}\xi^{1}+2^{3}\xi^{0}+\cdots\right)\right)\right)$$

n=0 n=-1 n=-2 n=0 n=-1 n=-2

$$|\xi| > 2$$
 $\chi(\xi) = \frac{\xi^{-1}}{|-\rho.5\xi^{-1}|} - \frac{\xi^{-1}}{|-\rho.5\xi^{-1}|} + (\frac{1}{2})^{n-1} - 2^{n-1}$ $(n \ge 1)$

$$\frac{U=1}{\left(\frac{7}{17} \right)_{0}^{2} \xi_{-1} + \left(\frac{7}{17} \right)_{1}^{2} \xi_{-2} + \left(\frac{7}{17} \right)_{2}^{2} \xi_{-2} + \cdots } - \left(\frac{7}{17} \right)_{0}^{2} \xi_{-1} + \frac{1}{17} \left(\frac{7}{17} \right)_{1}^{2} \xi_{-2} + \frac{1}{17} \left(\frac{7}{17} \right)_{1}^{2} \xi_{-2} + \cdots }$$

$$-\mathbf{B}^{\frac{3}{2}}\frac{-\mathbf{z}^{2}}{(2-2)(2-0.5)} = \frac{0.52}{(2-0.5)} - \frac{22}{(2-1)}$$

$$|\xi| < 0.5$$
 $\chi(\xi) = -\frac{\xi}{|-2\xi|} + \frac{\xi}{|-0.5\xi|} -(\frac{1}{2})^{\eta+1} + 2^{\eta+1}$ $(\eta < 0)$

$$- \left(\frac{1}{4} \right)_{0} \xi + \left(\frac{1}{4} \right)_{-1} \xi_{2} + \left(\frac{1}{4} \right)_{-2} \xi_{3} + \cdots \right) + \left(\frac{1}{4} \right) \xi_{3} + \left(\frac{1}{4} \right)_{5} \xi_{3} + \cdots \right)$$

$$- \left(\frac{1}{4} \right)_{0} \xi + \left(\frac{1}{4} \right)_{-1} \xi_{3} + \left(\frac{1}{4} \right)_{-2} \xi_{3} + \cdots \right) + \left(\frac{1}{4} \right) \xi_{3} + \cdots \right)$$

$$|\xi| > 2$$
 $|\xi| > 2$ $|-as \epsilon^{-1}| - \frac{2}{|-as \epsilon^{-1}|} + \frac{1}{2} |-as \epsilon^{-1}| + \frac{1}{2} |-as \epsilon^{-1}|$

$$\frac{\left(\frac{1}{2}\right) + \left(\frac{1}{2}\right)^{2} \xi^{-1} + \left(\frac{1}{2}\right)^{3} \xi^{-2} + \cdots + \left(2 + 2^{2} \xi^{-1} + 2^{3} \xi^{-2} + \cdots \right)}{n = 0 \quad n = 1 \quad n = 2}$$

$$\frac{3}{2} \frac{-1}{(2-05)(2-2)} = \begin{pmatrix} \frac{1}{2-0.5} - \frac{1}{2-2} \end{pmatrix}$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| >$$

$$\frac{3}{2} \frac{-1}{(2-0.5)(2-2)} = \left(\frac{1}{\xi-0.5} - \frac{1}{\xi-2}\right)$$

$$|\xi| < 0.5 \qquad f(z) = -\frac{2}{1-2\xi} + \frac{0.5}{1-0.5\xi} -2^{n+1} + (\frac{1}{2})^{n+1} \qquad (n > 0)$$

$$-\left(2z^{0} + 2^{1} \xi^{1} + 2^{3} \xi^{2} + \cdots\right) + \left((\frac{1}{2})z^{0} + (\frac{1}{2})^{3} \xi^{1} + (\frac{1}{2})^{3} \xi^{1} + \cdots\right)$$

$$|\xi| < 0.5 \qquad \chi(z) = -\frac{2}{1-2\xi} + \frac{0.5}{1-0.5\xi} -(\frac{1}{2})^{n-1} + 2^{n-1} \qquad (n < 0)$$


$$|\xi| < 0.5 \qquad \chi(\xi) = -\frac{2}{1 - 2\xi} + \frac{0.5}{1 - 0.5\xi} - (\frac{1}{2})^{n-1} + 2^{n-1} \qquad (n < 0)$$


$$- (2'\xi'' + 2^2\xi' + 2^3\xi^2 + \cdots) + ((\frac{1}{2})\xi'' + (\frac{1}{2})^3\xi^1 + (\frac{1}{2})^3\xi^1 + \cdots)$$

$$- ((\frac{1}{2})^1 \underline{z}^0 + (\frac{1}{2})^2 \xi^1 + (\frac{1}{2})^3 \xi^1 + \cdots) + (2^{-1}\underline{z}^0 + 2^{-2}\xi^1 + 2^3\xi^2 + \cdots)$$


$$= 0 \quad n = -1 \quad n = -2$$

$$= 0.5 \quad (n < 0)$$





$$\frac{3}{3} \frac{-1}{(3-0.5)(3-2)} = \left(\frac{\xi-0.5}{1} - \frac{1}{\xi-2}\right)$$




$$\int (\xi) = (-2) \frac{0.5}{0.5 - \xi} + (0.5) \frac{2}{2 - \xi} \qquad \left(|\xi| < 0.5 \right)$$

$$a_n = (-2) \ 2^n + (0.5) \ (\frac{1}{2})^n \ (n \ge 0)$$

$$-2^{n+1} + (\frac{1}{2})^{n+1}$$

$$\frac{3}{2} \frac{-2^{2}}{(2-2)(2-0.5)} = \left(\frac{0.52}{(2-0.5)} - \frac{22}{(2-1)}\right) \qquad |2| > 2$$



$$X(z) = 0.5 \frac{z}{z-0.5} - 2 \frac{z}{z-1} \qquad (|z| > 2)$$




$$Q_n = (0.5) \left(\frac{1}{2}\right)^n - 2 \cdot 2^n \qquad (n \geqslant 0)$$

$$\left(\frac{1}{2}\right)^{n+1} - 2^{n+1}$$

Anti-Causal
$$f(z)$$
 $X(z)$ $|z| > 2$ $|z| < 0.5$

$$\frac{3}{2} \frac{-1}{(2-0.5)(2-2)} = \left(\frac{2-0.5}{2-0.5} - \frac{1}{2-2}\right)$$


|2| >2 |2| >0.5



$$\int (\xi) = (-2) \frac{-0.5}{0.5 - \xi} + (0.5) \frac{-2}{2 - \xi} \qquad (|\xi| > 0.5)$$

$$a_n = (+2) \ 2^n - (0.5) \ (\frac{1}{2})^n \ (n < 0)$$
 $+2^{n+1} - (\frac{1}{2})^{n+1}$

$$\frac{3}{2} \frac{-2^{2}}{(2-2)(2-0.5)} = \left(\frac{0.52}{(2-0.5)} - \frac{22}{(2-1)}\right) \qquad |2| < 2$$



$$X(z) = 0.5 \frac{-\xi}{\xi - 0.5} - 2 \frac{-\xi}{\xi - \lambda} \qquad (|z| < 2)$$



$$\alpha_n = -(0.5)(\frac{1}{2})^n + 2 \cdot 2^n \qquad (n < 0)$$

$$-(\frac{1}{2})^{n+1} + 2^{n+1}$$

$$\bigcirc -\bigcirc = \frac{3}{2} \frac{(3-0.5)(3-2)}{(3-2)} = \boxed{f(3)}$$


$$\frac{3}{2} \frac{-1}{(2-0.5)(2-2)} = \left(\frac{1}{\xi-0.5} - \frac{1}{\xi-2}\right) = \frac{-2}{1-2\xi} + \frac{0.5}{1-0.5\xi}$$

$$f(\bar{z}) = \frac{\frac{(-2)}{1 - (2\bar{z})} + \frac{(\frac{1}{2})}{1 - (\frac{2}{2})}}{= -\sum_{n=0}^{\infty} (2)^{n+i} (\bar{z})^n + \sum_{n=0}^{\infty} (\frac{1}{2})^{n+i} (\bar{z})^n}$$
$$= -\sum_{n=0}^{\infty} (2)^{n+i} \bar{z}^n + \sum_{n=0}^{\infty} (\frac{1}{2})^{n+i} \bar{z}^n$$

$$a_n =$$

$$a_n = -2^{n+i} + \left(\frac{1}{2}\right)^{n+i}$$

$$\frac{3}{2} \frac{-1}{(2-0.5)(2-2)} = \left(\frac{1}{\xi-0.5} - \frac{1}{\xi-2}\right) = \frac{\xi^{-1}}{1-0.5\xi^{-1}} - \frac{\xi^{-1}}{1-2\xi^{-1}}$$



$$\frac{f(\xi)}{f(\xi)} = \frac{\left(\frac{1}{\xi}\right)}{1 - \left(\frac{1}{2\xi}\right)} - \frac{\left(\frac{1}{\xi}\right)}{1 - \left(\frac{3}{\xi}\right)}$$

$$= \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n \left(\frac{1}{\xi}\right)^{n+1} - \sum_{n=0}^{\infty} \left(2\right)^n \left(\frac{1}{\xi}\right)^{n+1}$$

$$= \sum_{n=1}^{\infty} \left(\frac{1}{2}\right)^{n-1} \xi^{-n} - \sum_{n=1}^{\infty} \left(2\right)^{n-1} \xi^{-n}$$

$$= \sum_{n=-1}^{-\infty} (2)^{n+1} \xi^n - \sum_{n=-1}^{-\infty} (\frac{1}{2})^{n+1} \xi^n$$

$$a_n$$

$$\frac{3}{2} \frac{-1}{(2-0.5)(2-2)} = \boxed{\chi(3)} \quad |z| < 0.5 \quad |z| > 2$$
anticausal causal

$$|z| < 0.5$$
 $|z| > 2$

anticausal Causal


$$\frac{3}{2} \frac{-1}{(2-0.5)(2-2)} = \left(\frac{1}{\xi-0.5} - \frac{1}{\xi-2}\right) = \frac{-2}{1-2\xi} + \frac{0.5}{1-0.5\xi}$$

$$\begin{array}{c} \times (\overline{z}) = \frac{\left(-2\right)}{1 - \left(2\overline{z}\right)} + \frac{\left(\frac{1}{a}\right)}{1 - \left(\frac{2}{a}\right)} \\ = -\sum\limits_{n=0}^{\infty} \left(2\right)^{n+i} (\overline{z})^n + \sum\limits_{n=0}^{\infty} \left(\frac{1}{2}\right)^{n+i} (\overline{z})^n \\ = -\sum\limits_{n=0}^{\infty} \left(2\right)^{n+i} \overline{z}^n + \sum\limits_{n=0}^{\infty} \left(\frac{1}{2}\right)^{n+i} \overline{z}^n \end{array}$$

$$= -\sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^{n-1} \xi^{-n} + \sum_{n=0}^{\infty} (2)^{n-1} \xi^{-n}$$

$$(n \le 0)$$
 $a_n = -(\frac{1}{2})^{n-1} + 2^{n-1}$

$$\frac{3}{2} \frac{-1}{(2-05)(2-2)} = \left(\frac{1}{\xi-0.5} - \frac{1}{\xi-2}\right) = \frac{\xi^{-1}}{1-0.5\xi^{-1}} - \frac{\xi^{-1}}{1-2\xi^{-1}}$$



$$(n > 0)$$
 $a_n = (\frac{1}{2})^{n-1} - (2)^{n-1}$

$$(2)-\triangle \frac{3}{2}\frac{-\xi^{2}}{(2-2)(2-0.5)} = \int (3) \frac{|\xi| < 0.5}{\text{causal}} \frac{|\xi| > 2}{\text{anticausal}}$$


$$\frac{3}{2} \frac{-2^{2}}{(2-2)(2-0.5)} = \left(\frac{0.52}{(2-0.5)} - \frac{22}{(2-1)}\right) = -\frac{2}{1-22} + \frac{2}{1-0.52}$$

$$\frac{1}{1 - (2\xi)} = -\frac{(\xi)}{1 - (2\xi)} + \frac{(\xi)}{1 - (\frac{\xi}{2})} \neq \frac{1}{1 - (\frac{\xi}{2})} = -\sum_{n=0}^{\infty} (2)^n (\xi)^{n+1} + \sum_{n=0}^{\infty} (\frac{1}{2})^n (\xi)^{n+1} = -\sum_{n=0}^{\infty} (2)^{n-1} \xi^n + \sum_{n=0}^{\infty} (\frac{1}{2})^{n-1} \xi^n$$

$$= -\sum_{n=0}^{\infty} (2)^{n-1} \xi^n + \sum_{n=0}^{\infty} (\frac{1}{2})^{n-1} \xi^n$$

$$(n > 0)$$
 $a_n = -2^{n-1} + (\frac{1}{2})^{n-1}$

$$\frac{3}{2} \frac{-2}{(2-2)(2-0.5)} = \left(\frac{0.52}{(2-0.5)} - \frac{22}{(2-1)}\right) = \frac{0.5}{1-0.52^{-1}} - \frac{2}{1-22^{-1}}$$



$$(n \leq 0) \qquad a_n = 2^{n-1} - \left(\frac{1}{2}\right)^{n-1}$$

$$-\left(\frac{1}{2}\right)^{n-1}$$

(2) - (B)
$$\frac{3}{2} \frac{-\xi^2}{(2-2)(2-0.5)} = [\chi(\xi)]$$

$$|z| < 0.5$$
 $|z| > 2$

anticausal causal

$$\frac{3}{2} \frac{-2^{2}}{(2-2)(2-0.5)} = \left(\frac{0.52}{(2-0.5)} - \frac{22}{(2-1)}\right) = -\frac{2}{1-22} + \frac{2}{1-0.52}$$

(n < 0) $a_n = -(\frac{1}{2})^{n+1} + 2^{n+1}$

$$\frac{3}{2} \frac{-2^{2}}{(2-2)(2-0.5)} = \left(\frac{0.52}{(2-0.5)} - \frac{22}{(2-1)}\right) = \frac{0.5}{1-0.52^{-1}} - \frac{2}{1-22^{-1}}$$

$$X(\xi) = \frac{\left(\frac{1}{2}\right)}{1 - \left(\frac{1}{2\xi}\right)} - \frac{\left(\frac{2}{2}\right)}{1 - \left(\frac{2}{\xi}\right)}$$

$$= \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^{n+1} \left(\frac{1}{\xi}\right)^n - \sum_{n=0}^{\infty} \left(2\right)^{n+1} \left(\frac{1}{\xi}\right)^n$$

$$= \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^{n+1} \xi^{-n} - \sum_{n=0}^{\infty} \left(2\right)^{n+1} \xi^{-n}$$

$$(n \geqslant 0) \qquad \alpha_n = \left(\frac{1}{2}\right)^{n+1} - 2^{n+1}$$

