

This is a very clear and well described overview of an important parasite. It needs in my opinion some more general and applied information to cover the potential audience of this article (e.g. clinical signs, diagnostics and some more details on the option for treatment – specifically anthelmintic).

Formatted: Font: Not Italic

Teladorsagia circumcincta is a nematode that parasitises sheep and goats. It was also previously known as *Ostertagia circumcincta* and colloquially refer to as the brown stomach worm and is common in cool temperate areas, such as south-eastern and south-western Australia and the United Kingdom. *Teladorsagia davtiani* and *Teladorsagia trifurcata* are probably phenotypic variants (morphotypes) of *T. circumcincta*.^[1] This parasite is responsible for huge considerable economic losses in sheep.^{[2][3][4]}

Comment [VB1]: What about goats?

Taxonomy[edit]

Teladorsagia is a member of the subfamily Ostertagiinae in the family Trichostrongylidae and the superfamily Trichostrongyoidea^{[5][6]}. Other taxa that are members of the subfamily are *Ostertagia*, *Haemonchus* and *Graphidium* are closely related taxa that are usually assigned to separate subfamilies.

Comment [VB2]: Should start from phylum, class etc.

Morphology[edit]

Adults are slender with a short buccal cavity and are ruddy brown in colour^[7]. The average worm size varies considerably among sheep. Females range in size from 0.6 to 1.2 cm^[8] with males typically about 20% smaller^[7].

Comment [VB3]: I don't think this is the right reference to this information

Life cycle[edit]

The life cycle is relatively simple. Male and female adults breed on the abomasal surface. Eggs are passed in the faeces and develop through first (L1), second (L2) and third stage larvae (L3). Third stage larvae are unable to feed and are the infective stage. Following ingestion, third-stage larvae rapidly moult within 48 hours into fourth stage larvae (L4) which develop in the gastric glands of the abomasum. About 8 days after infection, the young adults emerge from the gastric glands, mature and breed. The prepatent period can be as short as 12 days and up to 21 days^[9]. The number of eggs in the uterus of mature females can vary from less than 10 to more than 60^[9] and is heavily dependent on the host immune response^[10]. The number of eggs produced per day by an adult female worm has been estimated as ranging from 0 to approximately 350^[10]. Longer females lay more eggs.

Comment [VB4]: Is it proven? Can you add a citation?

Epidemiology[edit]

In most countries, infection resumes in the spring and is triggered by the production of large numbers of eggs by periparturient females^[11]. The key trigger appears to be a relative protein deficiency in the ewe^[12] and the periparturient rise can be prevented by feeding supplementary protein^{[8][13]}. In the stereotypical pattern, egg production (assessed by faecal egg counts in the lambs) rises till midsummer then declines^{[14][15]}. Nematode development is influenced by temperature and moisture and there is considerable variation among years in the pattern of egg production during the year^[15]. The infection gradually dies down at the end of the season. The end of the season is likely to vary among countries depending upon their climate and is also likely to vary from year to year within countries depending upon local weather but is poorly documented. In Scotland, the season usually ends about late October.

Comment [VB5]: This needs clarification (e.g. hypobiosis) or to be taken away as otherwise it is not clear

There is considerable variation among animals in faecal egg counts^[16] and much of this variation is genetic in origin^[17]. The sources of variation are dynamic and their relative importance changes over the course of the year. Other sources of variation, apart from inherited effects include maternal and common environmental effects and effects specific to each individual,

Comment [VB6]: Reduced? I don't think you can totally prevent it

Comment [VB7]: Rather than focusing on a specific area, can you give indication of temperature below which development is arrested (e.g. below 4°C)?

including variation in intake and non-additive genetic effects. The gender of the lamb, type of birth (single or twin), date of birth and intensity of early exposure to nematode infection have relatively minor but still important effects [16]. The inevitable variation between the observed count and the true faecal egg count also contributes to the observed variation [16].

Genetics^[edit]

The heritability of a trait in the narrow sense is the proportion of inherited variation divided by the total variation [18]. It determines the response to selection [19]. Faecal egg count is widely used to identify and select animals that are relatively resistant to nematode infection [20][21][22]. Selection is most advanced in Australia and New Zealand where the dominant nematodes are *Haemonchus contortus* or a mixture of *T. circumcincta* and *Trichostrongylus colubriformis*.

For naturally infected Scottish Blackface lambs grazing fields contaminated with predominantly *T. circumcincta* eggs, the heritability of faecal egg count was not significantly different from zero for the first two months after birth then rose to approximately 0.3 at the end of the grazing season [17][23]. Similar heritability estimates were obtained for a flock of Texel sheep [24]. This heritability is similar to the heritability of growth rate in beef cattle and milk production in dairy cattle and indicates the feasibility of selective breeding for resistance to natural predominantly *T. circumcincta* infection [24].

Genetic variation in faecal egg count in lambs following natural infection is predominantly due to genetic variation influencing worm size and fecundity rather than worm establishment and survival or larval inhibition [8]. Indeed, the heritability of adult female worm size, which is closely related to fecundity, has been estimated at 0.6 [8] whereas the heritability of worm number was only 0.14 [8]. The high heritability of worm size is one of the highest heritabilities known for a fitness related trait.

There has been a lot of effort devoted to identifying the causal polymorphisms that underlie resistance to infection [25][26]. Two genomic regions in particular have shown associations in a number of different studies: the *DRB1* locus of the major histocompatibility complex [27][28][29][30][31] and the interferon gamma locus [32]. Linkage disequilibrium in sheep is quite high compared to humans and the causative loci need not be in these regions but are probably in or around these regions. There are a number of regions which have shown suggestive (nearly but not quite statistically significant) associations in more than one study [26] and some of these are probably harbouring causal polymorphisms for nematode resistance.

Many of the putative candidate genes influence the immune response and genetic variation seems to predominantly influence immune responsiveness to infection. Although there are tantalising clues that other mechanisms may also contribute to genetic variation [33].

Immune responsiveness^[edit]

There are two major manifestations of resistance to *T. circumcincta*: the regulation of worm growth and the control of worm establishment and survival. The regulation of worm size and fecundity is strongly and consistently associated with IgA activity especially against fourth-stage larvae [9][34][35]. This effect is so strong and consistent that it is likely causal because there is no other immunological mechanism that is so strongly associated with IgA activity. Eosinophil number is also associated with worm length and fecundity [36] and it is likely that eosinophils and IgA interact to influence worm growth and fecundity. Eosinophils have been implicated in resistance to infection to *Haemonchus contortus* [37][38]. Eosinophils play a much weaker role in resistance of mice to parasitic infections [39] but this may reflect the fact that mice eosinophils lack the IgA receptor [40].

The other major mechanism of resistance to *T. circumcincta* is the IgE and mast cell mediated control of worm establishment and survival [9]. Again, this association is so strong and consistent that it is likely to be causal rather than acting as a marker for some other response. IgE activity against third-stage larvae seems to be most important [41][42]. The number of molecules recognised by IgE seems to be relatively small; two-dimensional Western blotting identified only 9 molecules

Comment [VB8]: Before this section, another one focusing on the diagnostic methods (explaining faecal egg count and other methods) should be added. Also, if the article is intended for general public (e.g. farmers, students), this section should be simplified and condensed

although there is evidence for at least two other allergens [43]. For comparison, 155 molecules were recognised by IgA on third-stage larvae [44].

Pathology [edit]

The immune response appears to be responsible for much of the pathology following nematode infection [45]. Immune suppression abolishes many of the clinical signs [46].

Infection causes a relative protein deficiency that leads to reduced weight gain or even weight loss. This due in part to a reduction in appetite [47][48]. There is also a loss in digestive efficiency. Lesions in the epithelial barrier allow a loss of protein and in addition protein is diverted to tissue repair and immune and inflammatory processes. Protein supplementation of the diet can prevent the appearance of clinical signs [47][49] which argues strongly that pathogenesis is a consequence of the relative protein deficiency.

Following infection, there is a breakdown of the mucosal barrier which is a consequence of mast cell degranulation releasing a serine protease mast cell protease II that digests occludin and claudin in the tight junctions between epithelial cells [45]. Many of the **clinical-pathological lesions signs** such as epithelial hyperplasia, increased pH in the abomasum **and** pepsinogenaemia **and** **anorexia** are a consequence of repair processes triggered by the binding of epithelial growth factor in saliva to receptors on the internal surface of epithelial cells [45]. Prolonged and sustained infection subverts the repair process into a pathogenic mechanism.

Mathematical modelling [edit]

A number of groups have modelled the infection process ranging from descriptive to semi-mechanistic to almost entirely mechanistic models [50][51][52][53]. These models can test the hypotheses arising from natural and deliberate infections [54] and suggest potential strategies for the control of infections [11][53][54][55].

The control of infection [edit]

The methods to control nematode infection in livestock can be grouped into 6 categories [56][57]: Anthelmintic treatment, grazing management, biological control, nutritional supplementation, vaccination and genetic resistance. Each method has its advantages and disadvantages. Anthelmintic treatment is cheap, simple and effective but is threatened by the development of drug resistance in parasite populations [58]. Grazing management is **cheap, simple and widely used** but is seldom very effective in isolation and can be impractical [59]. Biological control with the fungus *Duddingtonia flagrans* has recently been licensed in Australia and can be very effective [60] but requires regular, possibly daily, feeding. Nutritional supplementation can eliminate clinical signs but **can be very expensive** [61]. There is currently no completely reliable vaccine but great progress is being made [62]. Genetic resistance includes the use of relatively resistant breeds such as the Texel [29], cross-breeding and selective breeding. Selective breeding can be very effective [22] and almost completely eliminate the need for anthelmintic treatment but requires specialist expertise. Most experts feel that a combination of control methods will be more effective and sustainable. However, the most effective combination of control methods may vary from farm to farm and require specialist expertise to design and implement.

Comment [VB9]: This section would be better straight after epidemiology and should be called Pathology and clinical signs

Comment [VB10]: Here an expansion on the clinical signs (e.g. reduced appetite, poor growth performances, weight loss, diarrhoea etc.) should be added. A paragraph/sentence on the economics and the effect on production would also be beneficial

Comment [VB11]: After this section, one briefly outlining anthelmintic resistance should be added, with mention on the difference between whole flock methaphylactic treatments and targeted selective treatments. Also a brief overview of the 5 classes of anthelmintics available should be outlined

Comment [VB12]: It's not. Farmers argue all the time it takes too much of their time and most (in Scotland at least) won't do it.

Comment [VB13]: And also impractical

Additional information [edit]

Acknowledgements [edit]

Any people, organisations, or funding sources that you would like to thank.

Competing interests[edit]

Any conflicts of interest that you would like to declare. Otherwise, a statement that the authors have no competing interest.

Ethics statement[edit]

An ethics statement, if appropriate, on any animal or human research performed should be included here or in the methods section.

References[edit]

1. ↑ Grillo, Victoria; Craig, Barbara H.; Wimmer, Barbara; Gillear, John Stuart (2008-05). "[Microsatellite genotyping supports the hypothesis that *Teladorsagia davtiani* and *Teladorsagia trifurcata* are morphotypes of *Teladorsagia circumcincta*](#)". *Molecular and Biochemical Parasitology* **159** (1): 59–63. doi:[10.1016/j.molbiopara.2008.01.001](https://doi.org/10.1016/j.molbiopara.2008.01.001). ISSN [0166-6851](#).
2. ↑ Nieuwhof, G. J.; Bishop, S. C. (2005-08). "[Costs of the major endemic diseases of sheep in Great Britain and the potential benefits of reduction in disease impact](#)". *Animal Science* **81** (01). doi:[10.1079/asc41010023](https://doi.org/10.1079/asc41010023). ISSN [1357-7298](#).
3. ↑ Morgan, Eric; Charlier, Johannes; Hendrickx, Guy; Biggeri, Annibale; Catalan, Dolores; von Samson-Himmelstjerna, Georg; Demeler, Janina; Müller, Elizabeth *et al.* (2013-08-26). "[Global Change and Helminth Infections in Grazing Ruminants in Europe: Impacts, Trends and Sustainable Solutions](#)". *Agriculture* **3** (3): 484–502. doi:[10.3390/agriculture3030484](https://doi.org/10.3390/agriculture3030484). ISSN [2077-0472](#).
4. ↑ Geoffrey, Lane, Joe Jubb, Tristan Shephard, Richard Webb-Ware, John Fordyce, (2015). [Priority list of endemic diseases for the red meat industries](#). Meat & Livestock Australia Limited. [OCLC 1002179693](#).
5. ↑ Hoberg, E. P.; Lichtenfels, J. R. (1994-12). "[Phylogenetic Systematic Analysis of the Trichostrongylidae \(Nematoda\), with an Initial Assessment of Coevolution and Biogeography](#)". *The Journal of Parasitology* **80** (6): 976. doi:[10.2307/3283448](https://doi.org/10.2307/3283448). ISSN [0022-3395](#).
6. ↑ Lichtenfels, J.R.; Hoberg, E.P. (1993-02). "[The systematics of nematodes that cause ostertagiiasis in domestic and wild ruminants in North America: an update and a key to species](#)". *Veterinary Parasitology* **46** (1-4): 33–53. doi:[10.1016/0304-4017\(93\)90046-p](https://doi.org/10.1016/0304-4017(93)90046-p). ISSN [0304-4017](#).
7. ↑ [Jump up to:7.0 7.1](#)
8. ↑ [Jump up to:8.0 8.1 8.2 8.3 8.4](#) Stear, M. J.; Bairden, K.; Duncan, J. L.; Holmes, P. H.; McKellar, Q. A.; Park, M.; Strain, S.; Murray, M. *et al.* (1997-09). "[How hosts control worms](#)". *Nature* **389**(6646): 27–27. doi:[10.1038/37895](https://doi.org/10.1038/37895). ISSN [0028-0836](#).
9. ↑ [Jump up to:9.0 9.1 9.2 9.3](#) STEAR, M. J.; BISHOP, S. C.; DOLIGALSKA, M.; DUNCAN, J. L.; HOLMES, P. H.; IRVINE, J.; McCIRRIE, L.; McKELLAR, Q. A. *et al.* (1995-12). "[Regulation of egg production, worm burden, worm length and worm fecundity by host responses in sheep infected with *Ostertagia circumcincta*](#)". *Parasite Immunology* **17** (12): 643–652. doi:[10.1111/j.1365-3024.1995.tb01010.x](https://doi.org/10.1111/j.1365-3024.1995.tb01010.x). ISSN [0141-9838](#).
10. ↑ [Jump up to:10.0 10.1](#) Stear, M.J.; Bishop, S.C. (1999-05). "[The curvilinear relationship between worm length and fecundity of *Teladorsagia circumcincta*](#)". *International Journal for Parasitology* **29** (5): 777–780. doi:[10.1016/s0020-7519\(99\)00019-3](https://doi.org/10.1016/s0020-7519(99)00019-3). ISSN [0020-7519](#).
11. ↑ [Jump up to:11.0 11.1](#) SINGLETON, D. R.; STEAR, M. J.; MATTHEWS, L. (2010-10-15). "[A mechanistic model of developing immunity to *Teladorsagia circumcincta* infection in lambs](#)". *Parasitology* **138** (03): 322–332. doi:[10.1017/s0031182010001289](https://doi.org/10.1017/s0031182010001289). ISSN [0031-1820](#).
12. ↑ Donaldson, J.; van Houtert, M. F. J.; Sykes, A. R. (1998-12). "[The effect of nutrition on the periparturient parasite status of mature ewes](#)". *Animal Science* **67** (03): 523–533. doi:[10.1017/s1357729800032951](https://doi.org/10.1017/s1357729800032951). ISSN [1357-7298](#).
13. ↑ Houdijk, J.G.M.; Kyriazakis, I.; Jackson, F.; Huntley, J.F.; Coop, R.L (2000-07). "[Can an increased intake of metabolizable protein affect the periparturient relaxation in immunity against *Teladorsagia circumcincta* in sheep?](#)". *Veterinary Parasitology* **91** (1-2): 43–62. doi:[10.1016/s0304-4017\(00\)00255-7](https://doi.org/10.1016/s0304-4017(00)00255-7). ISSN [0304-4017](#).
14. ↑ STEAR, M. J.; BAIRDEN, K.; BISHOP, S. C.; GETTINBY, G.; McKELLAR, Q. A.; PARK, M.; STRAIN, S.; WALLACE, D. S. (1998-08). "[The processes influencing the distribution of parasitic](#)

Comment [VB14]: Reference missing

- nematodes among naturally infected lambs". *Parasitology* **117** (2): 165–171. doi:[10.1017/s0031182098002868](https://doi.org/10.1017/s0031182098002868). ISSN [0031-1820](#).
- 15. ↑ Jump up to:[15.0](#) [15.1](#) STEAR, M. J.; ABUAGOB, O.; BENOTHMAN, M.; BISHOP, S. C.; INNOCENT, G.; KERR, A.; MITCHELL, S. (2005-10-06). "Variation among faecal egg counts following natural nematode infection in Scottish Blackface lambs". *Parasitology* **132** (02): 275. doi:[10.1017/s0031182005009029](https://doi.org/10.1017/s0031182005009029). ISSN [0031-1820](#).
 - 16. ↑ Jump up to:[16.0](#) [16.1](#) [16.2](#) Stear, M.J.; Park, M.; Bishop, S.C. (1996-11). "The key components of resistance to *Ostertagia circumcincta* in lambs". *Parasitology Today* **12** (11): 438–441. doi:[10.1016/0169-4758\(96\)10069-7](https://doi.org/10.1016/0169-4758(96)10069-7). ISSN [0169-4758](#).
 - 17. ↑ Jump up to:[17.0](#) [17.1](#) Bishop, S. C.; Bairden, K.; McKellar, Q. A.; Park, M.; Stear, M. J. (1996-12). "Genetic parameters for faecal egg count following mixed, natural, predominantly *Ostertagia circumcincta* infection and relationships with live weight in young lambs". *Animal Science* **63** (03): 423–428. doi:[10.1017/s1357729800015319](https://doi.org/10.1017/s1357729800015319). ISSN [1357-7298](#).
 - 18. ↑ *Introduction to Quantitative Genetics*. By D. S. Falconer and T. F. C. Mackay. Fourth Edition. Longman. 1996. 464 pages.. 1996. doi:[10.1017/s001667230003408x](https://doi.org/10.1017/s001667230003408x). ISSN [0016-6723](#).
 - 19. ↑ M., Lynch, (1998). *Genetics and analysis of quantitative traits*. Sinauer. ISBN [0878934812](#). OCLC [709410500](#).
 - 20. ↑ Woolaston, R. R.; Windon, R. G. (2001-08). "Selection of sheep for response to *Trichostrongylus colubriformis* larvae: genetic parameters". *Animal Science* **73** (01): 41–48. doi:[10.1017/s1357729800058033](https://doi.org/10.1017/s1357729800058033). ISSN [1357-7298](#).
 - 21. ↑ Bisset, SA; Morris, CA; McEwan, JC; Vlassof, A (2001-12). "Breeding sheep in New Zealand that are less reliant on anthelmintics to maintain health and productivity". *New Zealand Veterinary Journal* **49** (6): 236–246. doi:[10.1080/00480169.2001.36238](https://doi.org/10.1080/00480169.2001.36238). ISSN [0048-0169](#).
 - 22. ↑ Jump up to:[22.0](#) [22.1](#) Karlsson, L. J. E.; Greeff, J. C. (2006). "Selection response in fecal worm egg counts in the Rylington Merino parasite resistant flock". *Australian Journal of Experimental Agriculture* **46** (7): 809. doi:[10.1071/ea05367](https://doi.org/10.1071/ea05367). ISSN [0816-1089](#).
 - 23. ↑ Stear, M.J.; Bairden, K.; Bishop, S.C.; Buitkamp, J.; Duncan, J.L.; Gettinby, G.; McKellar, Q.A.; Park, M. et al. (1997-09). "The genetic basis of resistance to *Ostertagia circumcincta* in lambs". *The Veterinary Journal* **154** (2): 111–119. doi:[10.1016/s1090-0233\(97\)80049-4](https://doi.org/10.1016/s1090-0233(97)80049-4). ISSN [1090-0233](#).
 - 24. ↑ Jump up to:[24.0](#) [24.1](#) Bishop SC, Jackson F, Coop RL, Stear MJ. Genetic parameters for resistance to nematode infections in Texel lambs and their utility in breeding programmes. *Anim Sci*2004;78:185-194.
 - 25. ↑ Riggio, V; Matika, O; Pong-Wong, R; Stear, M J; Bishop, S C (2013-03-20). "Genome-wide association and regional heritability mapping to identify loci underlying variation in nematode resistance and body weight in Scottish Blackface lambs". *Heredity* **110** (5): 420–429. doi:[10.1038/hdy.2012.90](https://doi.org/10.1038/hdy.2012.90). ISSN [0018-067X](#).
 - 26. ↑ Jump up to:[26.0](#) [26.1](#) Benavides, Magda Vieira; Sonstegard, Tad S.; Van Tassell, Curtis (2016-06). "Genomic Regions Associated with Sheep Resistance to Gastrointestinal Nematodes". *Trends in Parasitology* **32** (6): 470–480. doi:[10.1016/j.pt.2016.03.007](https://doi.org/10.1016/j.pt.2016.03.007). ISSN [1471-4922](#).
 - 27. ↑ Stear, M.J.; Innocent, G.T.; Buitkamp, J. (2005-10). "The evolution and maintenance of polymorphism in the major histocompatibility complex". *Veterinary Immunology and Immunopathology* **108** (1-2): 53–57. doi:[10.1016/j.vetimm.2005.07.005](https://doi.org/10.1016/j.vetimm.2005.07.005). ISSN [0165-2427](#).
 - 28. ↑ Schwaiger, F.-W.; Gostomski, D.; Stear, M.J.; Duncan, J.L.; McKellar, Q.A.; Epplen, J.T.; Buitkamp, J. (1995-07). "An ovine Major histocompatibility complex DRB1 allele is associated with low faecal egg counts following natural, predominantly *Ostertagia circumcincta* infection". *International Journal for Parasitology* **25** (7): 815–822. doi:[10.1016/0020-7519\(94\)00216-b](https://doi.org/10.1016/0020-7519(94)00216-b). ISSN [0020-7519](#).
 - 29. ↑ Jump up to:[29.0](#) [29.1](#) Sweeney T, Hanrahan J, Ryan M, Good B. Immunogenomics of gastrointestinal nematode infection in ruminants—breeding for resistance to produce food sustainably and safely. *Parasite Immunol*. 2016;38(9):569-586.
 - 30. ↑ Hassan, Musa; Good, Barbara; Hanrahan, James P; Campion, Deirdre; Sayers, Gearoid; Mulcahy, Grace; Sweeney, Torres (2011). "The dynamic influence of the DRB1*1101 allele on the resistance of sheep to experimental *Teladorsagia circumcincta* infection". *Veterinary Research* **42** (1): 46. doi:[10.1186/1297-9716-42-46](https://doi.org/10.1186/1297-9716-42-46). ISSN [1297-9716](#).
 - 31. ↑ SAYERS, G.; GOOD, B.; HANRAHAN, J. P.; RYAN, M.; ANGLES, J. M.; SWEENEY, T. (2005-06-08). "Major Histocompatibility Complex DRB1 gene: its role in nematode resistance in Suffolk and Texel sheep breeds". *Parasitology* **131** (03): 403. doi:[10.1017/s0031182005007778](https://doi.org/10.1017/s0031182005007778). ISSN [0031-1820](#).

32. ↑ COLTMAN, D. W.; WILSON, K.; PILKINGTON, J. G.; STEAR, M. J.; PEMBERTON, J. M. (2001-05). "A microsatellite polymorphism in the gamma interferon gene is associated with resistance to gastrointestinal nematodes in a naturally-parasitized population of Soay sheep". *Parasitology* **122** (05). doi:[10.1017/s0031182001007570](https://doi.org/10.1017/s0031182001007570). ISSN 0031-1820.
33. ↑ Hutchings, M.R; Knowler, K.J; McAnulty, R; McEwan, J.C (2007-08-07). "Genetically resistant sheep avoid parasites to a greater extent than do susceptible sheep". *Proceedings of the Royal Society B: Biological Sciences* **274** (1620): 1839–1844. doi:[10.1098/rspb.2007.0398](https://doi.org/10.1098/rspb.2007.0398). ISSN 0962-8452.
34. ↑ STRAIN, S. A. J.; BISHOP, S. C.; HENDERSON, N. G.; KERR, A.; MCKELLAR, Q. A.; MITCHELL, S.; STEAR, M. J. (2002-05). "The genetic control of IgA activity against *Teladorsagia circumcincta* and its association with parasite resistance in naturally infected sheep". *Parasitology* **124** (05). doi:[10.1017/s0031182002001531](https://doi.org/10.1017/s0031182002001531). ISSN 0031-1820.
35. ↑ MARTINEZ-VALLADARES, M.; VARA-DEL RIO, M. P.; CRUZ-ROJO, M. A.; ROJO-VAZQUEZ, F. A. (2005-06). "Genetic resistance to *Teladorsagia circumcincta*: IgA and parameters at slaughter in Churra sheep". *Parasite Immunology* **27** (6): 213–218. doi:[10.1111/j.1365-3024.2005.00769.x](https://doi.org/10.1111/j.1365-3024.2005.00769.x). ISSN 0141-9838.
36. ↑ STEAR, M. J.; HENDERSON, N. G.; KERR, A.; MCKELLAR, Q. A.; MITCHELL, S.; SEELEY, C.; BISHOP, S. C. (2002-05). "Eosinophilia as a marker of resistance to *Teladorsagia circumcincta* in Scottish Blackface lambs". *Parasitology* **124** (05). doi:[10.1017/s0031182002001580](https://doi.org/10.1017/s0031182002001580). ISSN 0031-1820.
37. ↑ González, Jorge F.; Hernández, Álvaro; Meeusen, Els N.T.; Rodríguez, Francisco; Molina, José M.; Jaber, José R.; Raadsma, Herman W.; Piedrafita, David (2011-06). "Fecundity in adult *Haemonchus contortus* parasites is correlated with abomasal tissue eosinophils and γδ T cells in resistant Canaria Hair Breed sheep". *Veterinary Parasitology* **178** (3-4): 286–292. doi:[10.1016/j.vetpar.2011.01.005](https://doi.org/10.1016/j.vetpar.2011.01.005). ISSN 0304-4017.
38. ↑ Meeusen, E.N.T; Balic, A (2000-03). "Do Eosinophils have a Role in the Killing of Helminth Parasites?". *Parasitology Today* **16** (3): 95–101. doi:[10.1016/s0169-4758\(99\)01607-5](https://doi.org/10.1016/s0169-4758(99)01607-5). ISSN 0169-4758.
39. ↑ Urban, J. F.; Katona, I. M.; Paul, W. E.; Finkelman, F. D. (1991-07-01). "Interleukin 4 is important in protective immunity to a gastrointestinal nematode infection in mice.". *Proceedings of the National Academy of Sciences* **88** (13): 5513–5517. doi:[10.1073/pnas.88.13.5513](https://doi.org/10.1073/pnas.88.13.5513). ISSN 0027-8424.
40. ↑ van Egmond, Marjolein; Damen, Cora A; van Spriel, Annemiek B; Vidarsson, Gestur; van Garderen, Evert; van de Winkel, Jan G.J (2001-04). "IgA and the IgA Fc receptor". *Trends in Immunology* **22** (4): 205–211. doi:[10.1016/s1471-4906\(01\)01873-7](https://doi.org/10.1016/s1471-4906(01)01873-7). ISSN 1471-4906.
41. ↑ Sinski, E.; Bairden, K.; Duncan, J.L.; Eisler, M.C.; Holmes, P.H.; McKellar, Q.A.; Murray, Max; Stear, M.J. (1995-09). "Local and plasma antibody responses to the parasitic larval stages of the abomasal nematode *Ostertagia circumcincta*". *Veterinary Parasitology* **59**(2): 107–118. doi:[10.1016/0304-4017\(94\)00761-z](https://doi.org/10.1016/0304-4017(94)00761-z). ISSN 0304-4017.
42. ↑ Miller, H.R.P. (1984-05). "The protective mucosal response against gastrointestinal nematodes in ruminants and laboratory animals". *Veterinary Immunology and Immunopathology* **6** (1-2): 167–259. doi:[10.1016/0165-2427\(84\)90051-5](https://doi.org/10.1016/0165-2427(84)90051-5). ISSN 0165-2427.
43. ↑ MURPHY, L.; ECKERSALL, P. D.; BISHOP, S. C.; PETTIT, J. J.; HUNTLEY, J. F.; BURCHMORE, R.; STEAR, M. J. (2010-03-17). "Genetic variation among lambs in peripheral IgE activity against the larval stages of *Teladorsagia circumcincta*". *Parasitology* **137** (08): 1249–1260. doi:[10.1017/s0031182010000028](https://doi.org/10.1017/s0031182010000028). ISSN 0031-1820.
44. ↑ Ellis, Samantha; Matthews, Jacqueline B.; Shaw, Darren J.; Paterson, Steve; McWilliam, Hamish E.G.; Inglis, Neil F.; Nisbet, Alasdair J. (2014-09). "Ovine IgA-reactive proteins from *Teladorsagia circumcincta* infective larvae". *International Journal for Parasitology* **44** (10): 743–750. doi:[10.1016/j.ijpara.2014.05.007](https://doi.org/10.1016/j.ijpara.2014.05.007). ISSN 0020-7519.
45. ↑ [Jump up to:45.1 45.2](#) Stear, M. J.; Bishop, S. C.; Henderson, N. G.; Scott, I. (2003-06). "A key mechanism of pathogenesis in sheep infected with the nematode *Teladorsagia circumcincta*". *Animal Health Research Reviews* **4** (01): 45–52. doi:[10.1079/ahrr200351](https://doi.org/10.1079/ahrr200351). ISSN 1466-2523.
46. ↑ Greer, A.W.; Huntley, J.F.; Mackellar, A.; McAnulty, R.W.; Jay, N.P.; Green, R.S.; Stankiewicz, M.; Sykes, A.R. (2008-12). "The effect of corticosteroid treatment on local immune responses, intake and performance in lambs infected with *Teladorsagia circumcincta*". *International Journal for Parasitology* **38** (14): 1717–1728. doi:[10.1016/j.ijpara.2008.05.010](https://doi.org/10.1016/j.ijpara.2008.05.010). ISSN 0020-7519.
47. ↑ [Jump up to:47.0 47.1](#) Stear, M.J; Bairden, K; Duncan, J.L; Eckersall, P.D; Fishwick, G; Graham, P.A; Holmes, P.H; McKellar, Q.A *et al.* (2000-12). "The influence of relative resistance and urea-supplementation on deliberate infection with *Teladorsagia circumcincta* during winter". *Veterinary Parasitology* **94** (1-2): 45–54. doi:[10.1016/s0304-4017\(00\)00370-8](https://doi.org/10.1016/s0304-4017(00)00370-8). ISSN 0304-4017.

48. ↑ WALLACE, D. S.; BAIRDEN, K.; DUNCAN, J. L.; ECKERSALL, P. D.; FISHWICK, G.; GILL, M.; HOLMES, P. H.; McKELLAR, Q. A. et al. (1998-01). "The influence of dietary supplementation with urea on resilience and resistance to infection with *Haemonchus contortus*". *Parasitology* **116** (1): 67–72. doi:[10.1017/s0031182097001947](https://doi.org/10.1017/s0031182097001947). ISSN [0031-1820](#).
49. ↑ Coop, R.L.; Kyriazakis, I. (1999-08). "Nutrition-parasite interaction". *Veterinary Parasitology* **84** (3-4): 187–204. doi:[10.1016/s0304-4017\(99\)00070-9](https://doi.org/10.1016/s0304-4017(99)00070-9). ISSN [0304-4017](#).
50. ↑ Smith, G.; Grenfell, B.T. (1994-08). "Modelling of parasite populations: gastrointestinal nematode models". *Veterinary Parasitology* **54** (1-3): 127–143. doi:[10.1016/0304-4017\(94\)90087-6](https://doi.org/10.1016/0304-4017(94)90087-6). ISSN [0304-4017](#).
51. ↑ Bishop, S. C.; Stear, M. J. (1997-06). "Modelling responses to selection for resistance to gastrointestinal parasites in sheep". *Animal Science* **64** (03): 469–478. doi:[10.1017/s1357729800016088](https://doi.org/10.1017/s1357729800016088). ISSN [1357-7298](#).
52. ↑ Doeschl-Wilson, Andrea; Vagenas, Dimitrios; Kyriazakis, Ilias; Bishop, Stephen (2008). "Exploring the assumptions underlying genetic variation in host nematode resistance (Open Access publication)". *Genetics Selection Evolution* **40** (3): 241. doi:[10.1186/1297-9686-40-3-241](https://doi.org/10.1186/1297-9686-40-3-241). ISSN [1297-9686](#).
53. ↑ Jump up to [53.1](#) Prada Jimenez de Cisneros, J.; Stear, M. J.; Mair, C.; Singleton, D.; Stefan, T.; Stear, A.; Marion, G.; Matthews, L. (2014-08-13). "An explicit immunogenetic model of gastrointestinal nematode infection in sheep". *Journal of The Royal Society Interface* **11**(99): 20140416–20140416. doi:[10.1098/rsif.2014.0416](https://doi.org/10.1098/rsif.2014.0416). ISSN [1742-5689](#).
54. ↑ Jump up to [54.1](#) Laurenson, Yan C.S.M.; Kyriazakis, Ilias; Bishop, Stephen C. (2013-10). "Can we use genetic and genomic approaches to identify candidate animals for targeted selective treatment". *Veterinary Parasitology* **197** (1-2): 379–383. doi:[10.1016/j.vetpar.2013.04.039](https://doi.org/10.1016/j.vetpar.2013.04.039). ISSN [0304-4017](#).
55. ↑ DE CISNEROS, JOAQUÍN PRADA JIMÉNEZ; MATTHEWS, LOUISE; MAIR, COLETTE; STEFAN, THORSTEN; STEAR, MICHAEL J. (2014-02-13). "The transfer of IgA from mucus to plasma and the implications for diagnosis and control of nematode infections". *Parasitology* **141** (07): 875–879. doi:[10.1017/s0031182013002321](https://doi.org/10.1017/s0031182013002321). ISSN [0031-1820](#).
56. ↑ STEAR, M. J.; DOLIGALSKA, M.; DONSKOW-SCHMELTER, K. (2006-11-01). "Alternatives to anthelmintics for the control of nematodes in livestock". *Parasitology* **134**(02): 139. doi:[10.1017/s0031182006001557](https://doi.org/10.1017/s0031182006001557). ISSN [0031-1820](#).
57. ↑ Sayers, G.; Sweeney, T. (2005-12). "Gastrointestinal nematode infection in sheep – a review of the alternatives to anthelmintics in parasite control". *Animal Health Research Reviews* **6** (02): 159–171. doi:[10.1079/ahr2005108](https://doi.org/10.1079/ahr2005108). ISSN [1466-2523](#).
58. ↑ Bartley, D.J.; Jackson, F.; Jackson, E.; Sargison, N. (2004-09). "Characterisation of two triple resistant field isolates of *Teladorsagia* from Scottish lowland sheep farms". *Veterinary Parasitology* **123** (3-4): 189–199. doi:[10.1016/j.vetpar.2004.06.018](https://doi.org/10.1016/j.vetpar.2004.06.018). ISSN [0304-4017](#).
59. ↑ Gruner, Lucas; Cortet, Jacques; Sauvage, Christine; Hoste, Hervé (2004-01). [<http://dx.doi.org/10.1051/vetres:2003043>] "Regulation of *Teladorsagia circumcincta* and *Trichostrongylus colubriformis* worm populations by grazing sheep with differing resistance status". *Veterinary Research* **35** (1): 91–101. doi:[10.1051/vetres:2003043](https://doi.org/10.1051/vetres:2003043). ISSN [0928-4249](#).
60. ↑ Waller, P.J.; Schwan, O.; Ljungström, B.-L.; Rydzik, A.; Yeates, G.W. (2004-12). "Evaluation of biological control of sheep parasites using *Duddingtonia flagrans* under commercial farming conditions on the island of Gotland, Sweden". *Veterinary Parasitology* **126** (3): 299–315. doi:[10.1016/j.vetpar.2004.08.008](https://doi.org/10.1016/j.vetpar.2004.08.008). ISSN [0304-4017](#).
61. ↑ Coop, R.L.; Holmes, P.H. (1996-08). "Nutrition and parasite interaction". *International Journal for Parasitology* **26** (8-9): 951–962. doi:[10.1016/s0020-7519\(96\)80070-1](https://doi.org/10.1016/s0020-7519(96)80070-1). ISSN [0020-7519](#).
62. ↑ Nisbet, Alasdair J.; McNeilly, Tom N.; Greer, Andrew W.; Bartley, Yvonne; Oliver, E. Margaret; Smith, Stephen; Palarea-Albaladejo, Javier; Matthews, Jacqueline B. (2016-09). "Protection of ewes against *Teladorsagia circumcincta* infection in the periparturient period by vaccination with recombinant antigens". *Veterinary Parasitology* **228**: 130–136. doi:[10.1016/j.vetpar.2016.09.002](https://doi.org/10.1016/j.vetpar.2016.09.002). ISSN [0304-4017](#).