

# z-Trans-7: Comparison with the Laurent Series : Residue Integral View Point


20180207

from Complex Analysis in plain view  
Laurent Series and z-Transform Note  
- Properties of a Geometric Series  
20170831

Copyright (c) 2016 - 2017 Young W. Lim.

Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.2 or any later version published by the Free Software Foundation; with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts. A copy of the license is included in the section entitled "GNU Free Documentation License".

# General Series Expansion at $z=0$



$$f(z) = \sum_{n=\eta_1}^{\infty} a_n z^n$$

$$a_n = \frac{1}{2\pi i} \oint_C \frac{f(z)}{z^{n+1}} dz$$
$$= \sum_k \operatorname{Res} \left( \frac{f(z)}{z^{n+1}}, z_k \right)$$

$$a_n = \frac{1}{2\pi i} \oint_C \frac{f(z)}{z^{n+1}} dz$$

$z_k$ : Poles of  $\frac{f(z)}{z^{n+1}}$

$$n \geq 0 \quad z_1, z_2, z_3, \circ$$

$$n < 0 \quad z_1, z_2, z_3$$





To compare with the  $z$ -Transform,

consider only the Laurent Series


expanded at  $z=0$

# Laurent Series ROC's

General Series Expansion at  $z=0$  becomes  
a Laurent Series Expansion at these Co-centric ROC's



$$x[n] = \frac{1}{2\pi i} \int_C X(z) z^{n-1} dz$$
$$= \sum_k \text{Res}(X(z) z^{n-1}, z_k)$$



$z$ -transform

## \* General Series Expansion at $z=0$





$$f(z) = \sum_{n=n_1}^{\infty} a_n z^n$$

$$\begin{aligned} a_n &= \frac{1}{2\pi i} \oint_C \frac{f(z)}{z^{n+1}} dz \\ &= \sum_k \operatorname{Res}\left(\frac{f(z)}{z^{n+1}}, z_k\right) \end{aligned}$$

## \* $z$ -transform

$$X(z) = \sum_{k=0}^{\infty} x_k z^{-k}$$


$$\begin{aligned} x_n &= \frac{1}{2\pi i} \oint_C X(z) z^{n+1} dz \\ &= \sum_k \operatorname{Res}(X(z) z^{n+1}, z_k) \end{aligned}$$



$f_i(z)$


# ROC's of the L.S.'s

Case ①  $f_1(z) = \sum a_n z^n$




$$a_n = \frac{1}{2\pi i} \oint_C \frac{f(z)}{z^{n+1}} dz$$

Case ②  $f_2(z) = \sum b_n z^n$




$$b_n = \frac{1}{2\pi i} \oint_C \frac{f(z)}{z^{n+1}} dz$$

Case ③  $f_3(z) = \sum c_n z^n$



$$c_n = \frac{1}{2\pi i} \oint_C \frac{f(z)}{z^{n+1}} dz$$

Case ④  $f_4(z) = \sum d_n z^n$



$$d_n = \frac{1}{2\pi i} \oint_C \frac{f(z)}{z^{n+1}} dz$$



\*  $f_i(z)$  valid in its ROC  
converge

$a_n$

# ROC's for the Cauchy's Residue Theorem



Case A  $f_1(z) = \sum a_n z^n$

Case B  $f_2(z) = \sum b_n z^n$



Case C  $f_3(z) = \sum c_n z^n$

Case D  $f_4(z) = \sum d_n z^n$



\* determine which poles are used in the Residue Computations

# Signal Processing Applications

$$x_k \rightarrow X(z)$$

$\boxed{z.T.}$

$$a_n \rightarrow f(z)$$

$\boxed{L.S.}$

↑

Given signal sequence

Causal signal


$$\sum_{n=0}^{\infty}$$

$n_1 = 0$

Anti-causal signal

$$\sum_{n=-\infty}^0$$

$n_1 = -\infty \rightarrow n_1 = 0$





# Inverse z-Transform $x[n] = \frac{1}{2\pi i} \int_C X(z) z^{n+1} dz$

$$X(z) = \sum_{k=0}^{\infty} x_k z^{-k}$$

$$z^{n+1} X(z) = \left( \sum_{k=0}^{\infty} x_k z^{-k} \right) z^{n+1} \quad \int z^{n+1} \text{LHS} dz = \int \text{RHS} z^{n+1} dz$$

$$= \sum_{k=0}^{\infty} x_k z^{-k+n+1} \quad [0, \infty) = [0, n-1] \cup [n] \cup [n+1, \infty)$$

$$= \sum_{k=0}^{n-1} x_k z^{-k+n+1} + \sum_{k=n}^n x_k z^{-k+n+1} + \sum_{k=n+1}^{\infty} x_k z^{-k+n+1}$$

$$= \sum_{k=0}^{n-1} x_k z^{-k+n+1} + \frac{x_n}{z^1} + \sum_{k=n+1}^{\infty} \frac{x_k}{z^{k-n+1}}$$

$$\begin{aligned} \int_C X(z) z^{n+1} dz &= \int_C \sum_{k=0}^{n-1} x_k z^{-k+n+1} dz + \int_C \frac{x_n}{z^1} dz + \int_C \sum_{k=n+1}^{\infty} \frac{x_k}{z^{k-n+1}} dz \\ &= \sum_{k=0}^{n-1} x_k \int_C z^{-k+n+1} dz + x_n \int_C \frac{1}{z^1} dz + \sum_{k=n+1}^{\infty} x_k \int_C \frac{1}{z^{k-n+1}} dz \\ &= \sum_{k=0}^{n-1} x_k \cdot 0 + x_n \cdot 2\pi i + \sum_{k=n+1}^{\infty} x_k \cdot 0 \\ &= 2\pi i \cdot x_n \end{aligned}$$

$$x[n] = \frac{1}{2\pi i} \int_C X(z) z^{n+1} dz$$

$z$ -Transform  $X(z)$   
Laurent Series  $f(z)$

$z$ -Transform  $X(z) \longleftrightarrow x_n$   
Laurent Series  $f(z) \longleftrightarrow a_n$

$$X(z) = f(z^{-1}) \longleftrightarrow x_n = a_n$$

$z$ -Transform  $X(z) \longleftrightarrow x_n$   
Laurent Series  $f(z) \longleftrightarrow a_n$


$$X(z) = f(z) \longleftrightarrow x_n = a_{-n}$$

$$X(z) = f(z^{-1}), \quad x_n = a_n$$

$$f(z) = \dots + a_2 z^{-2} + a_1 z^{-1} + a_0 z^0 + a_1 z^1 + a_2 z^2 + \dots$$

$$\left[ \begin{aligned} f(z^{-1}) &= \dots + a_2 z^2 + a_1 z^1 + a_0 z^0 + a_1 z^{-1} + a_2 z^{-2} + \dots \\ X(z) &= \dots + x_{-2} z^2 + x_{-1} z^1 + x_0 z^0 + x_1 z^{-1} + x_2 z^{-2} + \dots \end{aligned} \right.$$

$$f(z^{-1}) = X(z) \quad \longleftrightarrow \quad a_n = x_n$$



$$X(z) = f(z^{-1}), \quad x_n = a_n$$

$$f(z) = \dots + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0z^0 + a_1z^1 + a_2z^2 + \dots$$

$$f(z^{-1}) = \dots + a_{-2}z^2 + a_{-1}z^1 + a_0z^0 + a_1z^{-1} + a_2z^{-2} + \dots$$

$$f(z^{-1}) = \dots + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0z^0 + a_1z^1 + a_2z^2 + \dots$$

$$f(z) \quad \dots \quad z^{-2} \quad z^{-1} \quad z^0 \quad z^1 \quad z^2 \quad \dots$$

...
 $a_{-2}$ 
 $a_{-1}$ 
 $a_0$ 
 $a_1$ 
 $a_2$ 
...

$$f(z^{-1}) \quad \dots \quad a_2 \quad a_1 \quad a_0 \quad a_1 \quad a_2 \quad \dots$$

...
 $a_2$ 
 $a_1$ 
 $a_0$ 
 $a_1$ 
 $a_2$ 
...

$$X(z) = \dots + x_{-2}z^{-2} + x_{-1}z^{-1} + x_0z^0 + x_1z^1 + x_2z^2 + \dots$$

$$X(z) = \dots + x_{-2}z^2 + x_{-1}z^1 + x_0z^0 + x_1z^{-1} + x_2z^{-2} + \dots$$

$$X(z) \quad \dots \quad z^{-2} \quad z^{-1} \quad z^0 \quad z^1 \quad z^2 \quad \dots$$

...
 $x_{-2}$ 
 $x_{-1}$ 
 $x_0$ 
 $x_1$ 
 $x_2$ 
...

z-Transform

$$X(z) \longleftrightarrow x_n$$

Laurent Series

$$f(z) \longleftrightarrow a_n$$

$$X(z) = f(z^{-1}) \longleftrightarrow x_n = a_n$$

$$a_n = x_n \quad \longrightarrow \quad X(z) = f(z^{-1})$$

$$f(z) = \sum_{n=n_1}^{\infty} a_n z^n$$

$$X(z) = \sum_{k=0}^{\infty} x_k z^{-k}$$

$$\begin{aligned} a_n &= \frac{1}{2\pi i} \oint_C \frac{f(z)}{z^{n+1}} dz \\ &= \sum_k \operatorname{Res}\left(\frac{f(z)}{z^{n+1}}, z_k\right) \end{aligned}$$

$$\begin{aligned} x_n &= \frac{1}{2\pi i} \oint_C X(z) z^{n+1} dz \\ &= \sum_k \operatorname{Res}(X(z) z^{n+1}, z_k) \end{aligned}$$

$$a_n = x_n \quad \longrightarrow \quad X(z) = f(z^{-1})$$

$$a_n = \sum_k \operatorname{Res}\left(\frac{f(z)}{z^{n+1}}, z_k\right)$$

$$\sum_k \operatorname{Res}(X(z) z^{n+1}, z_k)$$

conformal  
mapping

$$w = z^{-1}$$

$$\sum_k \operatorname{Res}(f(w^{-1}) w^{n+1}, w_k)$$

$$\therefore X(z) = f(z^{-1})$$

$$a_n = x_n \quad \leftarrow \quad X(z) = f(z^{-1})$$

$$f(z) = \sum_{n=n_1}^{\infty} a_n z^n$$

$$X(z) = \sum_{k=0}^{\infty} x_k z^{-k}$$

$$\begin{aligned} a_n &= \frac{1}{2\pi i} \oint_C \frac{f(z)}{z^{n+1}} dz \\ &= \sum_k \operatorname{Res}\left(\frac{f(z)}{z^{n+1}}, z_k\right) \end{aligned}$$

$$\begin{aligned} x_n &= \frac{1}{2\pi i} \oint_C X(z) z^{n+1} dz \\ &= \sum_k \operatorname{Res}(X(z) z^{n+1}, z_k) \end{aligned}$$

$$a_n = x_n \quad \leftarrow \quad X(z) = f(z^{-1})$$

$$\therefore a_n = x_n \quad a_n = \sum_k \operatorname{Res}\left(\frac{f(w)}{w^{n+1}}, w_k\right) = \sum_k \operatorname{Res}(X(z) z^{n+1}, z_k) = x_n$$

conformal mapping

$$\begin{array}{ccc} & \omega = z^{-1} & \\ & \nearrow & \\ & \sum_k \operatorname{Res}(f(z^{-1}) z^{n+1}, z_k) & \\ & \searrow & \\ & & X(z) = f(z^{-1}) \\ & & = f(\omega) \end{array}$$

$$X(z) = f(z^{-1}), \quad x_n = \boxed{a_n}$$

$a_n$   $\longleftrightarrow$   $f(z)$

$$a_n = \frac{1}{2\pi i} \oint_C \frac{f(z)}{z^{n+1}} dz$$

$$\boxed{z = w^{-1}} \quad \boxed{dz = -w^{-2} dw}$$

$$= \frac{-1}{2\pi i} \oint_{C'} \frac{f(w^{-1})}{w^{-(n+1)}} w^{-2} dw$$

$$= \frac{-1}{2\pi i} \oint_{C'} f(w^{-1}) w^{n-1} dw$$


$$a_n = \frac{1}{2\pi i} \oint_C f(w^{-1}) w^{n-1} dw$$

$$x_n = \frac{1}{2\pi i} \oint_C X(z) z^{n+1} dz$$


$$a_n = x_n \quad \longrightarrow \quad X(z) = f(z^{-1})$$

$$a_n = x_n \quad \longleftarrow \quad X(z) = f(z^{-1})$$

$f(z)$



$$X(z) = f(z^{-1}) \quad \int_{C'} = - \int_C$$



$$X(z) = f(z^{-1}), \quad \boxed{x_n} = a_n$$

$$x_n \longleftrightarrow X(z)$$

$$x_n = \frac{1}{2\pi i} \oint_C X(z) z^{n+1} dz$$

$$\boxed{z = w^{-1}} \quad \boxed{dz = -w^{-2} dw}$$

$$= \frac{-1}{2\pi i} \oint_{C'} X(w^{-1}) w^{-(n+1)} w^{-2} dw$$

$$= \frac{-1}{2\pi i} \oint_{C'} X(w^{-1}) w^{-(n+1)} dw$$


$$x_n = \frac{1}{2\pi i} \oint_C \frac{X(w^{-1})}{w^{n+1}} dw$$

$$a_n = \frac{1}{2\pi i} \oint_C \frac{f(z)}{z^{n+1}} dz$$


$$a_n = x_n \quad \longrightarrow \quad X(z^{-1}) = f(z)$$

$$a_n = x_n \quad \longleftarrow \quad X(z^{-1}) = f(z)$$

$f(z)$



$$X(z) = f(z^{-1}) \quad \int_{C'} = - \int_C$$





$$X(z) = f(z), \quad x_n = a_{-n}$$

$$X(z) = \dots + x_2 z^2 + x_1 z^1 + x_0 z^0 + x_{-1} z^{-1} + x_{-2} z^{-2} + \dots$$

$$\left[ \begin{array}{l} X(z) = \dots + x_2 z^2 + x_1 z^1 + x_0 z^0 + x_{-1} z^{-1} + x_{-2} z^{-2} + \dots \\ f(z) = \dots + a_{-2} z^{-2} + a_{-1} z^{-1} + a_0 z^0 + a_1 z^1 + a_2 z^2 + \dots \end{array} \right.$$

$$f(z) = X(z) \iff a_{-n} = x_n$$

$$X(z) = f(z), \quad x_n = a_{-n}$$

$$f(z) = \dots + a_{-2}z^{-2} + a_{-1}z^{-1} + a_0z^0 + a_1z^1 + a_2z^2 + \dots$$

$$f(z) \quad \begin{array}{cccccc} \dots & z^{-2} & z^{-1} & z^0 & z^1 & z^2 & \dots \\ \dots & a_{-2} & a_{-1} & a_0 & a_1 & a_2 & \dots \end{array}$$

$$X(z) = \dots + x_{-2}z^2 + x_{-1}z^1 + x_0z^0 + x_1z^{-1} + x_2z^{-2} + \dots$$

$$X(z) = \dots x_2z^{-2} + x_1z^{-1} + x_0z^0 + x_{-1}z^1 + x_{-2}z^2 + \dots$$

$$X(z) \quad \begin{array}{cccccc} \dots & z^2 & z^1 & z^0 & z^{-1} & z^{-2} & \dots \\ \dots & x_{-2} & x_{-1} & x_0 & x_1 & x_2 & \dots \end{array}$$

$$X(z) \quad \begin{array}{cccccc} \dots & z^{-2} & z^{-1} & z^0 & z^1 & z^2 & \dots \\ \dots & x_2 & x_1 & x_0 & x_{-1} & x_{-2} & \dots \end{array}$$

z-Transform

$$X(z) \longleftrightarrow x_n$$

Laurent Series

$$f(z) \longleftrightarrow a_n$$

$$X(z) = f(z) \longleftrightarrow x_n = a_{-n}$$

$$X(z) = f(z), \quad x_n = a_{-n}$$

$$f(z) = X(z) \quad \longleftrightarrow \quad a_{-n} = x_n$$

$$a_n = \frac{1}{2\pi i} \oint_C \frac{X(z)}{z^{n+1}} dz = \sum_k \operatorname{Res}\left(\frac{X(z)}{z^{n+1}}, z_k\right) \quad \text{L.T.}$$





$$x_n = a_{-n} = \frac{1}{2\pi i} \oint_C \frac{X(z)}{z^{-n+1}} dz = \sum_k \operatorname{Res}\left(\frac{X(z)}{z^{-n+1}}, z_k\right)$$

$$x_n = \frac{1}{2\pi i} \oint_C X(z) z^{n-1} dz = \sum_k \operatorname{Res}(X(z) z^{n-1}, z_k) \quad \text{Z-T.}$$

causal ( $n \geq 0$ )

anti-causal ( $n \leq 0$ )

two-sided



$f(z) + \text{ROC?}$

$f(z) + \text{ROC?}$



$f(z) + \text{ROC?}$

$X(z) + \text{ROC?}$

$X(z) + \text{ROC?}$

$X(z) + \text{ROC?}$

# Causal & Anti-causal Signals






$a_n, x_n$	L.S.	Z.T.
causal $(n \geq 0)$ $(\frac{1}{2})^n$ 	$f(z) = \frac{2}{2-z}$	$X(z) = \frac{z}{z-0.5}$
anti-causal $(n \leq 0)$ $(\frac{1}{2})^{-n}$ 	$f(z) = \frac{z}{z-0.5}$	$X(z) = \frac{2}{2-z}$
two-sided $(\frac{1}{2})^{-n}$ $(\frac{1}{2})^n$ 	$f(z) = \frac{2}{2-z} + \frac{z}{z-0.5}$	$X(z) = \frac{2}{2-z} + \frac{z}{z-0.5}$

L.S.

Z.T.




causal

$(n \geq 0)$






anticausal

$(n \leq 0)$




two-sided



\* no poles in ROC's

L.S.



$$f(z) = 1 + \left(\frac{1}{2}\right)^1 z^1 + \left(\frac{1}{2}\right)^2 z^2 + \dots$$

$$a_0 = 1$$


$$a_1 = \left(\frac{1}{2}\right)$$

$$a_2 = \left(\frac{1}{2}\right)^2$$

$$a_3 = \left(\frac{1}{2}\right)^3 \quad a_n = \left(\frac{1}{2}\right)^n$$

$$f(z) = \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n z^n$$

Z.T.



$$\chi(z) = 1 + \left(\frac{1}{2}\right) z^{-1} + \left(\frac{1}{2}\right)^2 z^{-2} + \dots$$


$$x_0 = 1$$

$$x_1 = \left(\frac{1}{2}\right)$$

$$x_2 = \left(\frac{1}{2}\right)^2$$

$$x_3 = \left(\frac{1}{2}\right)^3 \quad x_n = \left(\frac{1}{2}\right)^n$$

$$\chi(z) = \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n z^{-n}$$



$$f(z) = 1 + \left(\frac{1}{2}\right) z^{-1} + \left(\frac{1}{2}\right)^2 z^{-2} + \dots$$


$$a_0 = 1$$

$$a_{-1} = \left(\frac{1}{2}\right)^1 = \left(\frac{1}{2}\right)^{-(-1)}$$

$$a_{-2} = \left(\frac{1}{2}\right)^2 = \left(\frac{1}{2}\right)^{-(-2)}$$

$$a_{-3} = \left(\frac{1}{2}\right)^3 = \left(\frac{1}{2}\right)^{-(-3)} \quad a_n = \left(\frac{1}{2}\right)^{-n}$$

$$f(z) = \sum_{n=-\infty}^0 \left(\frac{1}{2}\right)^{-n} z^n$$



$$\chi(z) = 1 + \left(\frac{1}{2}\right) z^1 + \left(\frac{1}{2}\right)^2 z^2 + \dots$$

$$x_0 = 1$$


$$x_{-1} = \left(\frac{1}{2}\right)^1 = \left(\frac{1}{2}\right)^{-(-1)}$$

$$x_{-2} = \left(\frac{1}{2}\right)^2 = \left(\frac{1}{2}\right)^{-(-2)}$$

$$x_{-3} = \left(\frac{1}{2}\right)^3 = \left(\frac{1}{2}\right)^{-(-3)} \quad x_n = \left(\frac{1}{2}\right)^{-n}$$

$$\chi(z) = \sum_{n=-\infty}^0 \left(\frac{1}{2}\right)^{-n} z^{-n}$$

## Causal



$$f(z) = X(z^{-1}) = \frac{z^1}{z^1 - 0.5}$$

$$= \frac{1}{1 - 0.5z} = \frac{2}{2 - z}$$


$$a_n = \text{Res}\left(\frac{f(z)}{z^{n+1}}, 0\right)$$

$$= \text{Res}\left(\frac{2}{z^{n+1}(2-z)}, 0\right)$$

$$= \left(\frac{1}{2}\right)^n \quad (n \geq 0)$$

$$f(z) = 1 + \left(\frac{1}{2}\right)^1 z^1 + \left(\frac{1}{2}\right)^2 z^2 + \dots$$

$$= \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n z^n = \frac{2}{2 - z}$$



$$\dots 0, 0, 0, 1, b, b^2, b^3, \dots$$

$$X(z) = \sum_{n=-\infty}^{\infty} x_n z^{-n} = \sum_{n=0}^{\infty} \left(\frac{b}{z}\right)^n$$

$$= \frac{1}{1 - \frac{b}{z}} = \frac{z}{z - b} = \frac{z}{z - 0.5}$$

$$x_n = \text{Res}\left(X(z) z^{n+1}, 0.5\right)$$

$$= \text{Res}\left(\frac{z^n}{z - 0.5}, 0.5\right)$$

$$= \left(\frac{1}{2}\right)^n \quad (n \geq 0)$$

$$X(z) = 1 + \left(\frac{1}{2}\right) z^{-1} + \left(\frac{1}{2}\right)^2 z^{-2} + \dots$$

$$= \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n z^{-n} = \frac{z}{z - 0.5}$$



causal

$$n \geq 0$$

$$a_n = \text{Res}\left(\frac{f(z)}{z^{n+1}}, 0\right) = \text{Res}\left(\frac{2}{z^{n+1}(2-z)}, 0\right)$$

 $n \geq 0$   
causal

$$a_n = \text{Res}\left(\frac{2}{z^{n+1}(2-z)}, 0\right) = \left(\frac{1}{2}\right)^n$$

$$a_0 = \text{Res}\left(\frac{2}{z^1(2-z)}, 0\right) = 1$$

 $n=0$ 

$$a_1 = \text{Res}\left(\frac{2}{z^2(2-z)}, 0\right) = \frac{2}{1!} \frac{d}{dz} \frac{1}{2-z} \Big|_{z=0} = \frac{2}{(2-z)^2} = \left(\frac{1}{2}\right)^1$$

 $n=1$ 



$$a_2 = \text{Res}\left(\frac{2}{z^3(2-z)}, 0\right) = \frac{2}{2!} \frac{d^2}{dz^2} \frac{1}{2-z} \Big|_{z=0} = \frac{2}{(2-z)^3} = \left(\frac{1}{2}\right)^2$$

 $n=2$ 

$$a_3 = \text{Res}\left(\frac{2}{z^4(2-z)}, 0\right) = \frac{2}{3!} \frac{d^3}{dz^3} \frac{1}{2-z} \Big|_{z=0} = \frac{2}{(2-z)^4} = \left(\frac{1}{2}\right)^3$$

 $n=3$ 

$$a_4 = \text{Res}\left(\frac{2}{z^5(2-z)}, 0\right) = \frac{2}{4!} \frac{d^4}{dz^4} \frac{1}{2-z} \Big|_{z=0} = \frac{2}{(2-z)^5} = \left(\frac{1}{2}\right)^4$$

 $n=4$ 


the finite number of poles

$$n \geq 0$$

$$x_n = \text{Res}(X(z)z^{n-1}, 0.5)$$

$$X(z) = \frac{z}{z-0.5}$$

$$X(z)z^{n-1} = \frac{z^n}{z-0.5} \quad \text{pole: } 0.5$$



$$x_n = \text{Res}(X(z)z^{n-1}, 0.5) = \text{Res}\left(\frac{z^n}{z-0.5}, 0.5\right)$$

$$x_0 = \text{Res}\left(\frac{z^0}{z-0.5}, 0.5\right) = 1$$

$$x_1 = \text{Res}\left(\frac{z^1}{z-0.5}, 0.5\right) = \left(\frac{1}{2}\right)^1$$



$$x_2 = \text{Res}\left(\frac{z^2}{z-0.5}, 0.5\right) = \left(\frac{1}{2}\right)^2$$

$$x_3 = \text{Res}\left(\frac{z^3}{z-0.5}, 0.5\right) = \left(\frac{1}{2}\right)^3$$



the finite number of poles

## Anti-causal



$\dots, b^3, b^2, b^1, 1, 0, 0, 0, \dots$

$$f(z) = X(z^{-1}) = \frac{2}{2 - z^{-1}}$$

$$= \frac{2z}{2z - 1} = \frac{z}{z - 0.5}$$

$$X(z) = \sum_{n=-\infty}^0 x_n z^{-n} = \sum_{n=-\infty}^0 b^n z^{-n}$$

$$= \sum_{n=0}^{\infty} (bz)^n = \frac{1}{1 - bz} = \frac{2}{2 - z}$$

$$a_n = \text{Res}\left(\frac{f(z)}{z^{n+1}}, 0.5\right) \quad n \leq 0$$

$$= \text{Res}\left(\frac{z}{z^{n+1}(z-0.5)}, 0.5\right)$$

$$= \left(\frac{1}{2}\right)^{-n} \quad (n \leq 0)$$

$$x_n = \text{Res}\left(X(z)z^{n+1}, 0\right)$$

$$= \text{Res}\left(\frac{2}{z^{1-n}(2-z)}, 0\right)$$

$$= \left(\frac{1}{2}\right)^{-n} \quad (n \leq 0)$$

$$f(z) = 1 + \left(\frac{1}{2}\right)^{-1} z^{-1} + \left(\frac{1}{2}\right)^{-2} z^{-2} + \left(\frac{1}{2}\right)^{-3} z^{-3} + \dots$$

$$X(z) = 1 + \left(\frac{1}{2}\right) z^1 + \left(\frac{1}{2}\right)^2 z^2 + \left(\frac{1}{2}\right)^3 z^3 + \dots$$

$$= \sum_{n=-\infty}^0 \left(\frac{1}{2}\right)^{-n} z^n = \frac{z}{z - 0.5}$$

$$= \sum_{n=-\infty}^0 \left(\frac{1}{2}\right)^n z^{-n} = \frac{2}{2 - z}$$

$n \leq 0$ 

$$a_n = \text{Res}\left(\frac{f(z)}{z^{n+1}}, 0.5\right) = \text{Res}\left(\frac{z}{z^{n+1}(z-0.5)}, 0.5\right)$$

 $n \leq 0$ 

\*  $z=0$  ~~pole~~ for  $n \leq 0$

$$\frac{z}{z^{n+1}(z-0.5)} = \frac{1}{z^n(z-0.5)}$$

$$\frac{1}{z^0}, \frac{1}{z^{-1}}, \frac{1}{z^{-2}}, \dots = 1, z, z^2, \dots$$

\*  $z = \frac{1}{2}$  the only pole

$$a_n = \text{Res}\left(\frac{1}{z^n(z-0.5)}, 0.5\right) = \left(\frac{1}{2}\right)^{-n} \quad n \leq 0$$



$$a_0 = \text{Res}\left(\frac{1}{z^0(z-0.5)}, 0.5\right) = 1 \quad n=0$$

$$a_{-1} = \text{Res}\left(\frac{1}{z^{-1}(z-0.5)}, 0.5\right) = \left(\frac{1}{2}\right)^1 \quad n=-1$$

$$a_{-2} = \text{Res}\left(\frac{1}{z^{-2}(z-0.5)}, 0.5\right) = \left(\frac{1}{2}\right)^2 \quad n=-2$$

$$a_{-3} = \text{Res}\left(\frac{1}{z^{-3}(z-0.5)}, 0.5\right) = \left(\frac{1}{2}\right)^3 \quad n=-3$$

$$a_{-4} = \text{Res}\left(\frac{1}{z^{-4}(z-0.5)}, 0.5\right) = \left(\frac{1}{2}\right)^4 \quad n=-4$$



$n \leq 0$ 

$$x_n = \text{Res}(X(z)z^{n+1}, 0)$$

$$X(z) = \frac{2}{2-z}$$

$$X(z)z^{n+1} = \frac{2z^{n+1}}{2-z} \quad \text{pole: } 2$$

$$x_n = \text{Res}\left(\frac{2z^{n+1}}{2-z}, 0\right) = \text{Res}\left(\frac{2}{z^{1-n}(2-z)}, 0\right) \quad (n \leq 0)$$



$$x_0 = \text{Res}\left(\frac{2}{z^1(2-z)}, 0\right) = 1 \quad n=0$$

$$x_{-1} = \text{Res}\left(\frac{2}{z^2(2-z)}, 0\right) = \frac{2}{1!} \frac{d}{dz} \frac{1}{2-z} \Big|_{z=0} = \frac{2}{(2-z)^2} = \left(\frac{1}{2}\right)^1 \quad n=1$$


$$x_{-2} = \text{Res}\left(\frac{2}{z^3(2-z)}, 0\right) = \frac{2}{2!} \frac{d^2}{dz^2} \frac{1}{2-z} \Big|_{z=0} = \frac{2}{(2-z)^3} = \left(\frac{1}{2}\right)^2 \quad n=2$$

$$x_{-3} = \text{Res}\left(\frac{2}{z^4(2-z)}, 0\right) = \frac{2}{3!} \frac{d^3}{dz^3} \frac{1}{2-z} \Big|_{z=0} = \frac{2}{(2-z)^4} = \left(\frac{1}{2}\right)^3 \quad n=3$$

$$x_{-4} = \text{Res}\left(\frac{2}{z^5(2-z)}, 0\right) = \frac{2}{4!} \frac{d^4}{dz^4} \frac{1}{2-z} \Big|_{z=0} = \frac{2}{(2-z)^5} = \left(\frac{1}{2}\right)^4 \quad n=4$$



# Summary




$$X(z^{-1}) = \frac{z^{-1}}{z^{-1} - 0.5} = \frac{1}{1 - (z/2)}$$

$$f(z) = \frac{2}{2-z} = \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n z^n$$

$$a_n = \left(\frac{1}{2}\right)^n \quad (n \geq 0)$$

$$= p^{-n} \quad (n \geq 0) \quad p=2$$




... 0, 0, 0, 1, b, b<sup>2</sup>, b<sup>3</sup>, ...

$$X(z) = \frac{z}{z - 0.5} = \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n z^{-n}$$

$$x_n = \left(\frac{1}{2}\right)^n \quad (n \geq 0)$$

$$= p^n \quad (n \geq 0) \quad p = \frac{1}{2}$$




$$X(z^{-1}) = \frac{2}{2 - z^{-1}}$$

$$f(z) = \frac{z}{z - 0.5} = \sum_{n=-\infty}^0 \left(\frac{1}{2}\right)^{-n} z^n$$

$$a_n = \left(\frac{1}{2}\right)^{-n} \quad (n \leq 0)$$

$$= p^{-n} \quad (n \leq 0) \quad p = \frac{1}{2}$$




... b<sup>3</sup>, b<sup>2</sup>, b<sup>1</sup>, 1, 0, 0, 0, ...

$$X(z) = \frac{2}{2-z} = \sum_{n=-\infty}^0 \left(\frac{1}{2}\right)^{-n} z^{-n}$$

$$x_n = \left(\frac{1}{2}\right)^{-n} \quad (n \leq 0)$$

$$= p^n \quad (n \leq 0) \quad p = 2$$




$$X(z^{-1}) = \frac{z^{-1}}{z^{-1} - b} = \frac{1}{1 - bz}$$

$$= (bz)^0 + (bz)^1 + (bz)^2 + \dots$$

$$f(z) = \frac{b^{-1}}{b^{-1} - z} = \sum_{n=0}^{\infty} b^n z^n$$

$$a_n = b^n \quad (n \geq 0)$$

$$p = b^{-1} \quad |z| < p \quad |pz| < 1$$




$$\dots, 0, 0, 0, 1, b, b^2, b^3, \dots$$

$$\left(\frac{b}{z}\right)^0 + \left(\frac{b}{z}\right)^1 + \left(\frac{b}{z}\right)^2 + \dots$$

$$X(z) = \frac{1}{1 - \frac{b}{z}} = \frac{z}{z - b}$$

$$x_n = b^n \quad (n \geq 0)$$

$$p = b \quad |z| > p \quad \left|\frac{p}{z}\right| < 1$$




$$X(z^{-1}) = \frac{b^{-1}}{b^{-1} - z^{-1}} = \frac{1}{1 - \frac{b}{z}}$$

$$= \left(\frac{b}{z}\right)^0 + \left(\frac{b}{z}\right)^1 + \left(\frac{b}{z}\right)^2 + \dots$$

$$f(z) = \frac{z}{z - b} = \sum_{n=-\infty}^0 b^{-n} z^n$$

$$a_n = b^{-n} \quad (n \leq 0)$$

$$p = b$$



$$\dots, b^3, b^2, b^1, 1, 0, 0, 0, \dots$$

$$(bz)^0 + (bz)^1 + (bz)^2 + \dots$$

$$X(z) = \frac{1}{1 - bz} = \frac{b^{-1}}{b^{-1} - z}$$

$$x_n = b^{-n} \quad (n \leq 0)$$

$$p = b^{-1} \quad |z| < p \quad \left|\frac{z}{p}\right| < 1$$

L.S.

Z.T.

causal

$(n \geq 0)$

$|z| < p$        $|p^{-1}z| < 1$

$$f(z) = \frac{p}{p-z} = \sum_{n=0}^{\infty} p^{-n} z^n$$

$$= \sum_{n=0}^{\infty} b^n z^n$$

$p = b'$

$|z| > p$        $|p z^{-1}| < 1$

$$X(z) = \frac{z}{z-p} = \sum_{n=0}^{\infty} p^n z^{-n}$$

$$= \sum_{n=0}^{\infty} b^n z^{-n}$$

$p = b$

anticausal

$(n \leq 0)$

$|z| > p$        $|p z^{-1}| < 1$

$$f(z) = \frac{z}{z-p} = \sum_{n=0}^{\infty} p^n z^{-n}$$

$$= \sum_{n=0}^{-\infty} p^{-n} z^n = \sum_{n=0}^{-\infty} b^{-n} z^n$$

$p = b$

$|z| < p$        $|p^{-1}z| < 1$

$$X(z) = \frac{p}{p-z} = \sum_{n=0}^{\infty} p^{-n} z^n$$

$$= \sum_{n=0}^{-\infty} p^n z^{-n} = \sum_{n=0}^{-\infty} b^{-n} z^{-n}$$


$p = b'$



L. S.

Z. T.

causal


 $(n \geq 0)$ 

$$f(z) = \frac{p}{p-z} = \frac{2}{2-z}$$

$$a_n = \text{Res}\left(\frac{f(z)}{z^{n+1}}, 0\right)$$

$$= \text{Res}\left(\frac{2}{z^{n+1}(2-z)}, 0\right)$$

$$= \left(\frac{1}{2}\right)^n \quad (n \geq 0)$$




$$X(z) = \frac{z}{z-p} = \frac{z}{z-0.5}$$

$$x_n = \text{Res}\left(X(z)z^{n-1}, 0.5\right)$$

$$= \text{Res}\left(\frac{z^n}{z-0.5}, 0.5\right)$$

$$= \left(\frac{1}{2}\right)^n \quad (n \geq 0)$$

anticausal


 $(n \leq 0)$ 

$$f(z) = \frac{z}{z-p} = \frac{z}{z-0.5}$$

$$a_n = \text{Res}\left(\frac{f(z)}{z^{n+1}}, 0.5\right)$$

$$= \text{Res}\left(\frac{z}{z^{n+1}(z-0.5)}, 0.5\right)$$

$$= \left(\frac{1}{2}\right)^{-n} \quad (n \leq 0)$$




$$X(z) = \frac{p}{p-z} = \frac{2}{2-z}$$

$$x_n = \text{Res}\left(X(z)z^{n-1}, 0\right)$$

$$= \text{Res}\left(\frac{2}{z^{1-n}(2-z)}, 0\right)$$


$$= \left(\frac{1}{2}\right)^{-n} \quad (n \leq 0)$$



$$X(z^{-1}) = \frac{z^{-1}}{z^{-1} - 0.5} \quad |z| < 2$$


$$f(z) = \frac{z}{z - 0.5} = \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n z^n$$

$$a_n = \left(\frac{1}{2}\right)^n = p^{-n} \quad \boxed{p=2}$$



$$X(z) = \frac{z}{z - 0.5} = \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n z^{-n} \quad |z| > \frac{1}{2}$$


$$x_n = \left(\frac{1}{2}\right)^n = p^n \quad \boxed{p=\frac{1}{2}}$$



$$X(z^{-1}) = \frac{z}{z - 0.5} \quad |z| > \frac{1}{2}$$

$$f(z) = \frac{z}{z - 0.5} = \sum_{n=-\infty}^0 \left(\frac{1}{2}\right)^{-n} z^n = \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n z^{-n}$$

$$a_n = \left(\frac{1}{2}\right)^{-n} = p^{-n} \quad \boxed{p=\frac{1}{2}}$$




$$X(z) = \frac{z}{z - 0.5} = \sum_{n=-\infty}^0 \left(\frac{1}{2}\right)^{-n} z^{-n} \quad |z| < 2$$

$$= \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n z^n$$

$$x_n = \left(\frac{1}{2}\right)^{-n} = p^n \quad \boxed{p=2}$$

$$(bz)^n \quad (n \geq 0)$$

$$|z| < b^{-1}$$




$$f(z) = \frac{1}{1 - bz} = \frac{b^{-1}}{b^{-1} - z}$$

$$a_n = b^n$$


$$= p^{-n}$$

$$p = b^{-1}$$



$$(bz^{-1})^n \quad (n \geq 0)$$

$$|z| > b$$




$$X(z) = \frac{1}{1 - b/z} = \frac{z}{z - b}$$

$$x_n = b^n$$


$$= p^{-n}$$

$$p = b$$



$$(n \leq 0) \quad (b^{-1}z)^n = (bz^{-1})^{-n}$$

$$|z| > b$$




$$f(z) = \frac{1}{1 - (bz^{-1})} = \frac{z}{z - b}$$

$$a_n = b^{-n}$$


$$= p^{-n}$$

$$p = b$$



$$(n \leq 0) \quad (b^{-1}z^{-1})^n = (bz)^{-n}$$

$$|z| < b^{-1}$$




$$X(z) = \frac{1}{1 - (bz)} = \frac{b^{-1}}{b^{-1} - z}$$

$$x_n = b^{-n}$$



$$= p^{-n}$$

$$p = b^{-1}$$





# Two-sided



$$\frac{1}{2} < |z| < 2 \Rightarrow \left| \frac{1}{2z} \right| < 1, \left| \frac{z}{2} \right| < 1$$

$$\begin{aligned} \frac{1}{1 - \frac{1}{2z}} + \frac{1}{1 - \frac{z}{2}} &= \frac{2z}{2z-1} + \frac{2}{2-z} \\ &= \frac{z}{z-0.5} - \frac{2}{z-2} \end{aligned} \quad \times$$

$$\boxed{\frac{1}{1 - \frac{1}{2z}} + \frac{1}{1 - \frac{z}{2}} - 1 = \frac{z}{z-0.5} - \frac{2}{z-2} - 1}$$

$$\frac{1}{1 - \frac{1}{2z}} = \left(\frac{1}{2z}\right)^0 + \left(\frac{1}{2z}\right)^1 + \left(\frac{1}{2z}\right)^2 + \left(\frac{1}{2z}\right)^3 + \dots = \frac{2z}{2z-1} = \frac{z}{z-0.5}$$

$$\textcircled{1} \left(\frac{1}{2z}\right)^1 + \left(\frac{1}{2z}\right)^2 - \left(\frac{1}{2z}\right)^3 + \dots = \frac{z}{z-0.5} - 1 = \frac{0.5}{z-0.5}$$

$$\frac{1}{1 - \frac{z}{2}} = \left(\frac{z}{2}\right)^0 + \left(\frac{z}{2}\right)^1 + \left(\frac{z}{2}\right)^2 + \left(\frac{z}{2}\right)^3 + \dots = \frac{2}{2-z}$$

$$f(z) = \frac{0.5}{z-0.5} - \frac{2}{z-2} = \frac{\cancel{\frac{1}{2}z} - 2\cancel{z}}{(z-0.5)(z-2)} = \frac{-\frac{3}{2}z}{(z-0.5)(z-2)}$$

$$\boxed{\frac{z}{z-0.5} + \frac{2}{2-z} - 1}$$

$$f(z) = \frac{0.5}{z-0.5} - \frac{2}{z-2} = \frac{\cancel{\frac{1}{2}z} - 2\cancel{z}}{(z-0.5)(z-2)} = \frac{-\frac{3}{2}z}{(z-0.5)(z-2)}$$

$$f(z^{-1}) = \frac{0.5}{z^{-1}-0.5} - \frac{2}{z^{-1}-2} = \frac{\cancel{\frac{1}{2}z} - 2\cancel{z}}{(z-0.5)(z-2)} = \frac{-\frac{3}{2}z}{(z-0.5)(z-2)}$$

$$= \frac{0.5z}{1-0.5z} - \frac{2z}{1-2z}$$

$$= \frac{z}{2-z} - \frac{z}{0.5-z}$$

$$= \frac{-z}{z-2} + \frac{z}{z-0.5} = \frac{-z^2 + 0.5z + z^2 - 2z}{(z-0.5)(z-2)} = \frac{-\frac{3}{2}z}{(z-0.5)(z-2)}$$

$$f(z) = f(z^{-1}) = X(z)$$

$$X(z) = X(z^{-1}) = f(z)$$

$$\underbrace{\left(\frac{1}{2z}\right)^1 + \left(\frac{1}{2z}\right)^2 + \left(\frac{1}{2z}\right)^3 + \dots}_{\text{blue arrow}} = \frac{0.5}{z-0.5} = \sum_{n=1}^{\infty} \left(\frac{1}{2z}\right)^n$$

$$\underbrace{\left(\frac{z}{2}\right)^0 + \left(\frac{z}{2}\right)^1 + \left(\frac{z}{2}\right)^2 + \left(\frac{z}{2}\right)^3 + \dots}_{\text{pink arrow}} = \frac{2}{2-z} = \sum_{n=0}^{\infty} \left(\frac{z}{2}\right)^n$$

$$\dots + \left(\frac{z}{2}\right)^3 + \left(\frac{z}{2}\right)^2 + \left(\frac{z}{2}\right)^1 + \left(\frac{z}{2}\right)^0 + \underbrace{\left(\frac{1}{2z}\right)^1 + \left(\frac{1}{2z}\right)^2 + \left(\frac{1}{2z}\right)^3 + \dots}_{\text{blue arrow}} = \boxed{\frac{2}{2-z}} + \boxed{\frac{0.5}{z-0.5}}$$


$$\begin{aligned} &= \frac{0.5}{z-0.5} + \frac{2}{2-z} \\ &= \frac{0.5}{z-0.5} - \frac{2}{z-2} \\ &= \frac{\cancel{\frac{1}{2}z} - 2\cancel{z}}{(z-0.5)(z-2)} \\ &= \frac{-\frac{3}{2}z}{(z-0.5)(z-2)} \end{aligned}$$

$$= \sum_{n=0}^{\infty} \left(\frac{1}{2z}\right)^n + \sum_{n=0}^{\infty} \left(\frac{z}{2}\right)^n - 1$$

$$= \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n z^{-n} + \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n z^n - 1$$


$$= \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n z^{-n} + \sum_{k=0}^{-\infty} \left(\frac{1}{2}\right)^k z^{-k} - 1$$

( $n \leq 0$ )      ( $n \geq 0$ )




$$\frac{z}{z-0.5} + \frac{2}{2-z} - 1$$

two-sided



$$\frac{1}{2} < |z| < 2$$

$$|\frac{1}{2}z^{-1}| < 1, |2^+z| < 1$$





$$\frac{1}{2} < |z| < 2$$


$$|\frac{1}{2}z^{-1}| < 1, |2^+z| < 1$$

$$\frac{z}{z-0.5} + \frac{2}{2-z} - 1 = \frac{z}{z-0.5} + \frac{2}{2-z} - 1$$







$$|2^{-1}z| < 1 \quad \frac{z}{2-z}$$



$$|\frac{1}{2}z^{-1}| < 1 \quad \frac{z}{z-0.5}$$



$$|\frac{1}{2}z^{-1}| < 1 \quad \frac{z}{z-0.5}$$



$$|2z^{-1}| < 1 \quad \frac{z}{2-z}$$

... 0, 0, 0, 1,  $(\frac{1}{2})$ ,  $(\frac{1}{2})^2$ ,  $(\frac{1}{2})^3$ , ...

$$X(z) = \frac{1}{1 - \frac{0.5}{z}} = \frac{z}{z - 0.5}$$

$$\left| \frac{0.5}{z} \right| < 1 \quad |z| > 0.5$$

... 0, 0, 0,  $(\frac{1}{2})^1$ ,  $(\frac{1}{2})^2$ ,  $(\frac{1}{2})^3$ ,  $(\frac{1}{2})^4$ , ...

$$X(z) = \frac{0.5}{1 - \frac{0.5}{z}} = \frac{0.5z}{z - 0.5}$$

$$\left| \frac{0.5}{z} \right| < 1 \quad |z| > 0.5$$

⊙ ... 0, 0, 0, 0,  $(\frac{1}{2})$ ,  $(\frac{1}{2})^2$ ,  $(\frac{1}{2})^3$ , ...

$$X(z) = \frac{0.5}{1 - \frac{0.5}{z}} \cdot z^{-1} = \frac{0.5}{z - 0.5}$$

$$\left| \frac{0.5}{z} \right| < 1 \quad |z| > 0.5$$

⊙ ...  $(\frac{1}{2})^3$ ,  $(\frac{1}{2})^2$ ,  $(\frac{1}{2})$ , 1, 0, 0, 0, ...

$$X(z) = \frac{1}{1 - \frac{z}{2}} = \frac{2}{2 - z}$$

$$\left| \frac{z}{2} \right| < 1 \quad |z| < 2$$

$b^3, b^2, b^1, 1, a, a^2, a^3$

... 0, 0, 0, 0, a, a^2, a^3, ...





...  $b^3, b^2, b^1, 1, 0, 0, 0, \dots$

$$\begin{aligned} X(z) &= \frac{0.5}{z - 0.5} + \frac{2}{2 - z} \\ &= \frac{0.5}{z - 0.5} - \frac{2}{z - 2} \\ &= \frac{\cancel{\frac{1}{2}z} - 2\cancel{z}}{(z - 0.5)(z - 2)} \\ &= \frac{-\frac{3}{2}z}{(z - 0.5)(z - 2)} \end{aligned}$$

...  $b^3, b^2, b^1, 1, 0, 0, 0, \dots$

... 0, 0, 0, 1, a, a^2, a^3, ...

$$\begin{aligned} &\frac{z}{z - 0.5} + \frac{2}{2 - z} - 1 \\ &= \frac{z^2 - 2z - 2z + 1}{(z - 0.5)(z - 2)} + \\ &= \frac{\cancel{z^2} - 4z + \cancel{1} - \cancel{z^2} + 2.5z - \cancel{1}}{(z - 0.5)(z - 2)} \\ &= \frac{-1.5z}{(z - 0.5)(z - 2)} \end{aligned}$$

$a_n$  $x_n$  $f(z)$   
 $X(z)$   
ROCRes  
ROC $(n \geq 0)$ 

$$a_n = \text{Res}\left(\frac{f(z)}{z^{n+1}}, 0\right) + \text{Res}\left(\frac{f(z)}{z^{n+1}}, \frac{1}{2}\right)$$

 $x_n = \text{Res}\left(X(z)z^{n+1}, \frac{1}{2}\right)$  ( $n \geq 0$ ) $x_n = \text{Res}\left(X(z)z^{n+1}, 0\right)$  ( $n < 0$ ) $(n < 0)$ 

$$a_n = \text{Res}\left(\frac{f(z)}{z^{n+1}}, \frac{1}{2}\right)$$

$$+ \text{Res}\left(X(z)z^{n+1}, \frac{1}{2}\right)$$

$$\begin{aligned} \frac{f(z)}{z^{n+1}} &= \frac{-1.5z}{(z-0.5)(z-2)z^{n+1}} \\ &= \frac{-1.5}{(z-0.5)(z-2)z^n} \end{aligned}$$

$$\begin{aligned} X(z)z^{n+1} &= \frac{-1.5z}{(z-0.5)(z-2)} z^{n+1} \\ &= \frac{-1.5z^n}{(z-0.5)(z-2)} \end{aligned}$$

$$a_n = \sum_k \operatorname{Res}\left(\frac{f(z)}{z^{n+1}}, z_k\right)$$

$$f(z) = \frac{-\frac{3}{2}z}{(z-0.5)(z-2)}$$

$$a_n = \begin{cases} \operatorname{Res}\left(X(z)z^{n+1}, \frac{1}{2}\right) & (n \geq 0) \\ \operatorname{Res}\left(X(z)z^{n+1}, \frac{1}{2}\right) + \operatorname{Res}\left(X(z)z^{n+1}, 0\right) & (n < 0) \end{cases}$$

$$a_n = \begin{cases} \operatorname{Res}\left(\frac{-\frac{3}{2}z^n}{(z-0.5)(z-2)}, \frac{1}{2}\right) & (n \geq 0) \\ \operatorname{Res}\left(\frac{-\frac{3}{2}z^n}{(z-0.5)(z-2)}, \frac{1}{2}\right) + \operatorname{Res}\left(\frac{-\frac{3}{2}z^n}{(z-0.5)(z-2)}, 0\right) & (n < 0) \end{cases}$$

$$a_3 = \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^{-3}}, \frac{1}{2}\right) = \left(\frac{1}{2}\right)^3 = \left(\frac{1}{2}\right)^{-(-3)}$$

$$a_2 = \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^{-2}}, \frac{1}{2}\right) = \left(\frac{1}{2}\right)^2 = \left(\frac{1}{2}\right)^{-(-2)}$$

$$a_1 = \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^{-1}}, \frac{1}{2}\right) = \left(\frac{1}{2}\right)^1 = \left(\frac{1}{2}\right)^{-(-1)}$$

$$a_0 = \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^0}, \frac{1}{2}\right) = \left(\frac{1}{2}\right)^0 = \left(\frac{1}{2}\right)^0$$

$$a_{-1} = \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^1}, \frac{1}{2}\right) + \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^1}, 0\right) = \left(\frac{1}{2}\right)^1 + \left(\frac{-3}{2}\right) = \left(\frac{1}{2}\right)$$

$$a_{-2} = \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^2}, \frac{1}{2}\right) + \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^2}, 0\right) = \left(\frac{1}{2}\right)^2 - \frac{6}{4} = \left(\frac{1}{2}\right)^2$$

$$a_{-3} = \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^3}, \frac{1}{2}\right) + \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^3}, 0\right) = \left(\frac{1}{2}\right)^3 - \frac{6}{8} = \left(\frac{1}{2}\right)^3$$

$$x_n = \sum_k \operatorname{Res}(X(z) z^{n-1}, z_k)$$

$$X(z) = \frac{-\frac{3}{2}z}{(z-0.5)(z-2)}$$

$$x_n = \begin{cases} \operatorname{Res}\left(\frac{f(z)}{z^{n+1}}, \frac{1}{2}\right) + \operatorname{Res}\left(\frac{f(z)}{z^{n+1}}, 0\right) & (n \geq 0) \\ \operatorname{Res}\left(\frac{f(z)}{z^{n+1}}, \frac{1}{2}\right) & (n < 0) \end{cases}$$

$$x_n = \begin{cases} \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^n}, \frac{1}{2}\right) + \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^n}, 0\right) & (n \geq 0) \\ \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^n}, \frac{1}{2}\right) & (n < 0) \end{cases}$$

$$x_{-3} = \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^{-3}}, \frac{1}{2}\right) = \left(\frac{1}{2}\right)^3 = \left(\frac{1}{2}\right)^{-(-3)}$$

$$x_{-2} = \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^{-2}}, \frac{1}{2}\right) = \left(\frac{1}{2}\right)^2 = \left(\frac{1}{2}\right)^{-(-2)}$$

$$x_{-1} = \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^{-1}}, \frac{1}{2}\right) = \left(\frac{1}{2}\right)^1 = \left(\frac{1}{2}\right)^{-(-1)}$$

$$x_0 = \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^0}, \frac{1}{2}\right) = \left(\frac{1}{2}\right)^0 = \left(\frac{1}{2}\right)^0$$

$$x_1 = \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^1}, \frac{1}{2}\right) + \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^1}, 0\right) = \left(\frac{1}{2}\right)^1 + \left(\frac{-3}{2}\right) = \left(\frac{1}{2}\right)$$

$$x_2 = \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^2}, \frac{1}{2}\right) + \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^2}, 0\right) = \left(\frac{1}{2}\right)^2 - \frac{1}{4} = \left(\frac{1}{2}\right)^2$$

$$x_3 = \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^3}, \frac{1}{2}\right) + \operatorname{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^3}, 0\right) = \left(\frac{1}{2}\right)^3 - \frac{1}{8} = \left(\frac{1}{2}\right)^3$$

# Residue Computations

$$\text{Res}(G(z), z_0) \left\{ \begin{array}{l} \lim_{z \rightarrow z_0} (z-z_0) G(z) = a_{-1} \quad \text{Simple pole } z_0 \\ \frac{1}{(n-1)!} \lim_{z \rightarrow z_0} \frac{d^{n-1}}{dz^{n-1}} (z-z_0)^n G(z) = a_{-1} \quad n\text{-th order pole } z_0 \end{array} \right.$$

$$\text{Res}\left(\frac{-\frac{3}{2}}{(z-0.5)(z-2)z^n}, 0\right) ?$$

$$\frac{-\frac{3}{2}}{(z-0.5)(z-2)} = \frac{A}{z-0.5} + \frac{B}{z-2}$$

$$A = \frac{-\frac{3}{2}}{z-2} \Big|_{z=0.5} = 1 \quad \frac{-1}{z-2}$$

$$B = \frac{-\frac{3}{2}}{z-0.5} \Big|_{z=2} = \frac{-\frac{3}{2}}{\frac{3}{2}} = -1$$

$$\frac{-\frac{3}{2}}{(z-0.5)(z-2)} = \frac{1}{z-0.5} - \frac{1}{z-2}$$

$$\frac{d}{dz} \frac{-\frac{3}{2}}{(z-0.5)(z-2)} = \frac{-1}{(z-0.5)^2} + \frac{1}{(z-2)^2}$$

$$\frac{d^2}{dz^2} \frac{-\frac{3}{2}}{(z-0.5)(z-2)} = \frac{2}{(z-0.5)^3} - \frac{2}{(z-2)^3}$$

$$\frac{d^3}{dz^3} \frac{-\frac{3}{2}}{(z-0.5)(z-2)} = \frac{6}{(z-0.5)^4} + \frac{6}{(z-2)^4}$$

$$\left\{ \begin{array}{l} \frac{1}{1!} \frac{d}{dz} \frac{-\frac{3}{2}}{(z-0.5)(z-2)} = \frac{-1}{(z-0.5)^2} + \frac{1}{(z-2)^2} \\ \frac{1}{2!} \frac{d^2}{dz^2} \frac{-\frac{3}{2}}{(z-0.5)(z-2)} = \frac{1}{(z-0.5)^3} - \frac{1}{(z-2)^3} \\ \frac{1}{3!} \frac{d^3}{dz^3} \frac{-\frac{3}{2}}{(z-0.5)(z-2)} = \frac{-1}{(z-0.5)^4} + \frac{1}{(z-2)^4} \end{array} \right.$$

$$\text{Res}\left(\frac{z^{-\frac{3}{2}}}{(z-0.5)(z-2)z}, 0\right) = \frac{z^{-\frac{3}{2}}}{(z-0.5)(z-2)} \Big|_{z=0} = \left[ \frac{1}{(z-0.5)} - \frac{1}{(z-2)} \right]_{z=0}$$

$$= -2 + \frac{1}{2} = -\frac{3}{2}$$

$$\text{Res}\left(\frac{z^{-\frac{3}{2}}}{(z-0.5)(z-2)z^2}, 0\right) = \frac{d}{dz} \frac{z^{-\frac{3}{2}}}{(z-0.5)(z-2)} \Big|_{z=0} = \left[ \frac{-1}{(z-0.5)^2} + \frac{1}{(z-2)^2} \right]_{z=0}$$

$$= -4 + \frac{1}{4} = -\frac{15}{4}$$

$$\text{Res}\left(\frac{z^{-\frac{3}{2}}}{(z-0.5)(z-2)z^3}, 0\right) = \frac{1}{2!} \frac{d^2}{dz^2} \frac{z^{-\frac{3}{2}}}{(z-0.5)(z-2)} \Big|_{z=0} = \left[ \frac{1}{(z-0.5)^3} - \frac{1}{(z-2)^3} \right]_{z=0}$$

$$= \left(-8 + \frac{1}{8}\right) = -\frac{63}{8}$$

$$a_n = \begin{cases} \text{Res}\left(\frac{z^{-\frac{3}{2}}}{(z-0.5)(z-2)z^n}, \frac{1}{2}\right) + \text{Res}\left(\frac{z^{-\frac{3}{2}}}{(z-0.5)(z-2)z^n}, 0\right) = \left(\frac{1}{2}\right)^n & (n \geq 0) \\ \text{Res}\left(\frac{z^{-\frac{3}{2}}}{(z-0.5)(z-2)z^n}, \frac{1}{2}\right) = \left(\frac{1}{2}\right)^{-n} & (n < 0) \end{cases}$$

$$a_n = \begin{cases} \left(\frac{1}{2}\right)^n & (n \geq 0) \\ \left(\frac{1}{2}\right)^{-n} & (n < 0) \end{cases}$$





