

Residue Integrals and Laurent Series with non-annular region

20170216

Copyright (c) 2016 - 2017 Young W. Lim.

Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.2 or any later version published by the Free Software Foundation; with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts. A copy of the license is included in the section entitled "GNU Free Documentation License".

Based on

T.J. Cavicchi, Digital Signal Processing

Complex Analysis for Mathematics and Engineering
J. Mathews

Residue Theorem

D: Simply connected domain


C: Simple closed contour (CCW) in D

if $f(z)$ is **analytic** inside C and on C
except at the points z_1, z_2, \dots, z_k in C

then

$$\frac{1}{2\pi i} \int_C f(z) dz = \sum_{j=1}^k \text{Res}(f(z), z_j)$$


Singular points of $f(z)$: z_1, z_2, \dots, z_k



Integration of a function of a complex var.


$$\oint_C f(z) dz = 2\pi i \sum_{k=1}^n \text{Res}(f(z), z_k)$$

finite number k of
singular points z_k
residue theorem



$$\oint_C f(z) dz = 0 \quad \text{if } f(z) \text{ is analytic within and on } C$$

no singularity



$$\oint_C f(z) dz = 0 \quad \text{if } f(z) = F'(z) \text{ on } C$$

: $F(z)$ is an antiderivative of $f(z)$
fundamental theorem of calculus

$\oint_C f(z) dz = 0$ if $f(z)$ is continuous in D and
 $f(z) = F'(z)$: $F(z)$ is an antiderivative of $f(z)$
fundamental theorem of calculus

Series Expansion

can expand $f(z)$ about any point z_m
over powers of $(z - z_m)$

whether or not $f(z)$ is singular at z_m
or at other points between z and z_m

$$f(z) = \sum_{n=\eta_1}^{\infty} a_n^{(m)} (z - z_m)^n$$

① Laurent Series Expansion of $f(z)$ at z_m
general η_1 - depend on $f(z)$ and z_m

② z -transform of $a_n^{(m)}$
general η_1 - depend on $f(z)$

$$z_m = 0$$

③ Taylor Series Expansion of $f(z)$ at z_m
positive η_1 - depend on $f(z)$ and z_m ($\eta_1 > 0$)

④ MacLaurin Series Expansion of $f(z)$ at z_m
positive η_1 - depend on $f(z)$ ($\eta_1 > 0$)

$$z_m = 0$$

$$f(z) = \sum_{n=n_1}^{\infty} a_n^{(m)} (z - z_m)^n$$

$n_1 > 0$ pos powers

$$z_m = 0$$

① Laurent Series

③ Taylor Series

② z-transform

④ MacLaurin Series

* Expansion of $f(z)$ about any point z_m
over powers of $(z - z_m)$

$$f(z) = \sum_{n=n_1}^{\infty} a_n^{(m)} (z - z_m)^n$$

$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz$$
$$a_n^{(m)} = \sum_k \operatorname{Res} \left(\frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

for general $f(z)$

for general $f(z)$

$$a_n^{(m)} = \frac{1}{n!} f^{(n)}(z_m) \quad n_1 \geq 0$$

for analytic $f(z)$ within C

analytic $f(z) \longrightarrow \frac{f(z)}{(z - z_m)^{n+1}}$ has a pole at z_m
order of $n+1$

$$f(z) = \sum_{n=n_1}^{\infty} a_n^{(m)} (z - z_m)^n$$

z_m : possible poles of $f(z)$
not necessarily poles

$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z')}{(z' - z_m)^{n+1}} dz'$$
$$= \sum_k \text{Res} \left(\frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$




z_k : poles of $\frac{f(z)}{(z - z_m)^{n+1}}$
within C

$$= \frac{1}{n!} f^{(n)}(z_m) \quad n_1 \geq 0$$

Residue Theorem and Laurent Series

assumed there are (K) singularities (poles) of $f(z)$ in a region


let C_k is taken to enclose only one pole z



$a_n^{(1)}$ expanded at z_1

C_1 encloses z_1 only


$$\tilde{a}_{-1}^{(1)} = \text{Res}(f(z), z_1)$$



$a_n^{(2)}$ expanded at z_2

C_2 encloses z_2 only

$$\tilde{a}_{-1}^{(2)} = \text{Res}(f(z), z_2)$$



$a_n^{(3)}$ expanded at z_3

C_3 encloses z_3 only

$$\tilde{a}_{-1}^{(3)} = \text{Res}(f(z), z_3)$$

Cauchy's Residue Theorem



$f(z)$: **analytic** on and within C
except a finite number of **singular points**
 z_1, z_2, \dots, z_n within C

then

$$\int_C f(z) dz = 2\pi i \sum_{k=1}^n \text{Res}(f(z), z_k)$$

D : a simply connected domain

C : a simple closed contour in D



z_1

$$f(z) = \sum_{k=-\infty}^{+\infty} a_k (z - z_1)^k$$

$$a_{-1}^{(z_1)} = \frac{1}{2\pi i} \oint_{C_1} f(s) ds = \text{Res}(f(z), z_1)$$

z_2

$$f(z) = \sum_{k=-\infty}^{+\infty} a_k (z - z_2)^k$$

$$a_{-1}^{(z_2)} = \frac{1}{2\pi i} \oint_{C_2} f(s) ds = \text{Res}(f(z), z_2)$$

z_3


$$f(z) = \sum_{k=-\infty}^{+\infty} a_k (z - z_3)^k$$

$$a_{-1}^{(z_3)} = \frac{1}{2\pi i} \oint_{C_3} f(s) ds = \text{Res}(f(z), z_3)$$

Laurent Series with Annular Region expanded at each pole of $f(z)$

z_1 Laurent series expansion at z_1


$$f(z) = \sum_{n=-\infty}^{+\infty} a_n^{(1)} (z - z_1)^n$$



$$\begin{aligned} \tilde{a}_{-1}^{(1)} &= \text{Res}(f(z), z_1) \\ &= \frac{1}{2\pi i} \oint_{C_1} f(z) dz \end{aligned}$$

z_2 Laurent series expansion at z_2


$$f(z) = \sum_{n=-\infty}^{+\infty} a_n^{(2)} (z - z_2)^n$$




$$\begin{aligned} \tilde{a}_{-1}^{(2)} &= \text{Res}(f(z), z_2) \\ &= \frac{1}{2\pi i} \oint_{C_2} f(z) dz \end{aligned}$$

z_3 Laurent series expansion at z_3

$$f(z) = \sum_{n=-\infty}^{+\infty} a_n^{(3)} (z - z_3)^n$$




$$\begin{aligned} \tilde{a}_{-1}^{(3)} &= \text{Res}(f(z), z_3) \\ &= \frac{1}{2\pi i} \oint_{C_3} f(z) dz \end{aligned}$$



$$\int_C f(z) dz = 2\pi i \sum_{k=1}^n \text{Res}(f(z), z_k)$$

Residue Theorem + Laurent Series

* Whether z_m is singular or not







$$f(z) = \sum_{n=n_1}^{\infty} a_n^{(m)} (z - z_m)^n$$

$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz$$

$$= \sum_k \text{Res} \left(\frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

$$a_{-1}^{(m)} = \frac{1}{2\pi i} \oint_C f(z) dz$$

$$= \sum_k \text{Res} (f(z), z_k)$$



$$\tilde{a}_{-1}^{(1)} = \text{Res}(f(z), z_1)$$

$$\tilde{a}_{-1}^{(2)} = \text{Res}(f(z), z_2)$$

$$\tilde{a}_{-1}^{(3)} = \text{Res}(f(z), z_3)$$

$$a_{-1}^{(m)} = \tilde{a}_{-1}^{(1)} + \tilde{a}_{-1}^{(2)} + \tilde{a}_{-1}^{(3)}$$

$$a_{-1}^{(m)} = \text{Res}(f(z), z_1) + \text{Res}(f(z), z_2) + \text{Res}(f(z), z_3)$$

← Laurent Series coefficient $a_{-1}^{(2)}$

- Singular center z_i
- Annular region

This can be a residue if

- Singular center z_m
- Annular region

Laurent Series — Annular Region of Convergence
 — no singularity in this region


Residue — Laurent Series expanded at a pole

a punctured open disk


- Annular
- Isolated Singularity




$$\tilde{a}_{-1}^{\{1\}} = \text{Res}(f(z), z_1)$$




$$\tilde{a}_{-1}^{\{2\}} = \text{Res}(f(z), z_2)$$



$$\tilde{a}_{-1}^{\{3\}} = \text{Res}(f(z), z_3)$$



$$\tilde{a}_{-1}^{\{m\}} = \text{Res}(f(z), z_m)$$



Computing $a_n^{\{m\}}$

$$f(z) = \sum_{n=\nu_1}^{\infty} a_n^{\{m\}} (z - z_m)^n \quad \boxed{n \leftarrow k}$$

$$f(z) = \sum_{k=\nu_1}^{\infty} a_k^{\{m\}} (z - z_m)^k$$

for a given n $\frac{f(z)}{(z - z_m)^{n+1}} = \sum_{k=\nu_1}^{\infty} a_k^{\{m\}} (z - z_m)^{k-n-1}$ k : index variable
 n : fixed value

$$\oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz = \oint_C \sum_{k=\nu_1}^{\infty} a_k^{\{m\}} (z - z_m)^{k-n-1} dz$$

$$= \sum_{k=\nu_1}^{\infty} \oint_C a_k^{\{m\}} (z - z_m)^{k-n-1} dz \quad \boxed{k=n}$$

$$\oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz = \oint_C a_n^{\{m\}} \frac{1}{(z - z_m)} dz = 2\pi i \cdot a_n^{\{m\}}$$

$$\boxed{a_n^{\{m\}} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz}$$

$$\oint_C \left[\dots (z - z_m)^{-3} + (z - z_m)^{-2} + \frac{1}{(z - z_m)} + 1 + (z - z_m) + (z - z_m)^2 + \dots \right] dz$$

$$= \oint_C \frac{1}{(z - z_m)} dz = 2\pi i$$

Computing $a_n^{\{m\}}$ using Residues

expansion at z_m

$$\eta = -1 \quad \eta + 1 = 0 \quad (z - z_m)^{\eta+1} = 1$$

$$a_n^{\{m\}} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz$$


$$= \sum_k \operatorname{Res} \left(\frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

$$a_{-1}^{\{m\}} = \frac{1}{2\pi i} \oint_C f(z) dz$$

$$= \sum_k \operatorname{Res} (f(z), z_k)$$

$$a_{-1}^{\{m\}} = \frac{1}{2\pi i} \oint_C f(z) dz = \sum_k \operatorname{Res} (f(z), z_k)$$

$$a_{-1}^{\{m\}} = \cancel{\operatorname{Res} (f(z), z_m)}$$



$$f(z) = \sum_{n=n_1}^{\infty} a_n^{\{m\}} (z - z_m)^n$$

$$a_n^{\{m\}} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz$$

$$= \sum_k \operatorname{Res} \left(\frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

Residue \rightarrow Laurent series \rightarrow annular region \rightarrow expanded at a pole \star) a punctured open disk

$$f(z) = \sum_{n=n_1}^{\infty} a_n^{(m)} (z - z_m)^n$$

$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz$$
$$= \sum_k \operatorname{Res} \left(\frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

$$a_{-1}^{(m)} = \frac{1}{2\pi i} \oint_C f(z) dz$$
$$= \sum_k \operatorname{Res} (f(z), z_k)$$

⋮

$$a_{-3}^{(m)} = \sum_k \operatorname{Res} (f(z)(z - z_m)^2, z_k)$$

$$a_{-2}^{(m)} = \sum_k \operatorname{Res} (f(z)(z - z_m)^1, z_k)$$

$$a_{-1}^{(m)} = \sum_k \operatorname{Res} (f(z), z_k)$$

$$a_0^{(m)} = \sum_k \operatorname{Res} \left(\frac{f(z)}{(z - z_m)^1}, z_k \right)$$

$$a_1^{(m)} = \sum_k \operatorname{Res} \left(\frac{f(z)}{(z - z_m)^2}, z_k \right)$$

$$a_2^{(m)} = \sum_k \operatorname{Res} \left(\frac{f(z)}{(z - z_m)^3}, z_k \right)$$

⋮

Poles for Residue Computation

$$f(z) = \sum_{n=-\infty}^{\infty} a_n^{(m)} (z - z_m)^n$$


$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z')}{(z' - z_m)^{n+1}} dz'$$

$$= \sum_k \operatorname{Res} \left(\frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$


z_k within C : singularities of $\boxed{\frac{f(z)}{(z - z_m)^{n+1}}}$

$n \geq 0$ { poles of $f(z)$ } \cup { $z = z_m$ } $n = 0, 1, 2, \dots$


$n < 0$ { poles of $f(z)$ } $n = -1, -2, \dots$




$$\int_C f(z) dz = 2\pi i \operatorname{Res}(f(z), z_1) + 2\pi i \operatorname{Res}(f(z), z_2) + 2\pi i \operatorname{Res}(f(z), z_3)$$



$$\int_{C'} f(z) dz = 2\pi i \operatorname{Res}(f(z), z_1) + 2\pi i \operatorname{Res}(f(z), z_2)$$




$$\int_{C''} f(z) dz = 2\pi i \operatorname{Res}(f(z), z_1)$$



$$\int_{C'''} f(z) dz = 0$$

Series Expansion at z_m



$$f(z) = \sum_{n=n_1}^{\infty} a_n^{[m]} (z - z_m)^n$$


$$a_n^{[m]} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz$$

$$= \sum_k \text{Res} \left(\frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

$$a_{-1}^{[m]} = \frac{1}{2\pi i} \oint_C f(z) dz$$

$$= \sum_k \text{Res} (f(z), z_k)$$

[Annular Region]



* even for a nonsingular z_m

z_m can be a pole of


$$\frac{f(z)}{(z - z_m)^{n+1}} \text{ only if } n \geq 0$$

When computing

$$a_n^{[m]} = \sum_k \text{Res} \left(\frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

[Annular Region] & [z_m : isolated singularity]

A punctured open disk



$$f(z) = \sum_{n=n_1}^{\infty} a_n^{(m)} (z - z_m)^n$$

$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz$$


$$= \sum_k \text{Res} \left(\frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

$$a_{-1}^{(m)} = \frac{1}{2\pi i} \oint_C f(z) dz$$

$$= \sum_k \text{Res} (f(z), z_k)$$

$$= \text{Res} (f(z), z_m)$$

Series Expansion at $z=0$



$$f(z) = \sum_{n=n_1}^{\infty} a_n^{(m)} z^n$$


$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{z^{n+1}} dz$$
$$= \sum_k \text{Res} \left(\frac{f(z)}{z^{n+1}}, z_k \right)$$

Poles z_k


$$n \geq 0 \quad z_1, z_2, z_3, \circ$$

$$n < 0 \quad z_1, z_2, z_3$$

Series Expansion at z_m no annular region.



$$f(z) = \sum_{n=n_1}^{\infty} a_n^{(m)} (z - z_m)^n$$



$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz$$

$$= \sum_k \text{Res} \left(\frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

Let z_1, z_2, z_3 poles of $f(z)$

Then the poles of $\frac{f(z)}{(z - z_m)^{n+1}}$

| | |
|------------|----------------------|
| $n \geq 0$ | z_1, z_2, z_3, z_m |
| $n < 0$ | z_1, z_2, z_3 |

if C encloses only one pole z_0 ,
and the expansion at that pole z_0 is assumed,
then



$$a_{-1}^{\{0\}} = \frac{1}{2\pi i} \oint_{C_0} f(z) dz = \text{Res}(f(z), z_0)$$

Let



$$\tilde{a}_{-1}^{\{m\}} = \text{Res}(f(z), z_m)$$

notation \sim

the **residue** of $f(z)$ at z_m

using C_m which is in the **annulus** ROC

$$f(z) = \sum_{n=-\infty}^{+\infty} a_n^{\{m\}} (z - z_m)^n$$



$$\oint_C f(z) dz = 2\pi j \sum_{k=1}^M \tilde{a}_{-1}^{(k)} = 2\pi j \sum_{k=1}^M \text{Res}(f(z), z_k)$$

residue theorem

$$a_n = \sum_{k=1}^M \text{Res} \left(\frac{f(z)}{(z-z_k)^{n+1}}, z_k \right)$$

Laurent coefficient

\$C\$ encloses \$k\$ poles


\$C_k\$ encloses only the \$k\$-th pole

\$\tilde{a}_{-1}^{(k)}\$ the residue of the \$k\$-th pole enclosed by \$C\$, \$z_k\$

Non-annular region

$$f(z) = \sum_{n=n_1}^{\infty} a_n^{f, m} (z - z_m)^n$$

$$a_n^{f, m} = \frac{1}{2\pi i} \oint_C \frac{f(z')}{(z' - z_m)^{n+1}} dz'$$
$$= \sum_k \operatorname{Res} \left(\frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$



C is in the same **region of analyticity** of $f(z)$
typically a circle centered on z_m **non-annular ok**

z_k within C : **singularities** of $\frac{f(z)}{(z - z_m)^{n+1}}$

$n_1 = n_{f, m}$ depends on $f(z)$, z_m

$a_n^{f, m}$ depends on $f(z)$, z_m , **region of analyticity**

Whether $f(z)$ is **singular** at $z = z_m$ or not
or at other points between z and z_m

We can expand $f(z)$ about **any point** z_m
over powers of $(z - z_m)$.

Laurent's Theorem

f : analytic within the **annular** domain D

$$r < |z - z_0| < R$$

then

$$f(z) = \sum_{k=-\infty}^{+\infty} a_k (z - z_0)^k,$$

valid for $r < |z - z_0| < R$

The coefficients a_k are given by




$$a_k = \frac{1}{2\pi i} \oint_C \frac{f(s)}{(s - z_0)^{k+1}} ds, \quad k = 0, \pm 1, \pm 2, \dots$$

C : a simple closed curve
that lies entirely within D
that encloses z_0


Curve C & Domain D of the Laurent Series

$$f(z) = \sum_{n=-\infty}^{\infty} a_n^{(m)} (z - z_m)^n$$

$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z')}{(z' - z_m)^{n+1}} dz'$$
$$= \sum_k \operatorname{Res} \left(\frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$



Expansion Points and Evaluation Points



• z_0 : expansion point

z : evaluation point

•

which poles of $f(z)$ lie between the point of evaluation z and the point z_0 about which the expansion is formed

$\frac{f(z')}{(z' - z_0)}$ is analytic between C_1 & C_2

deformation theorem

$C_1 - C_2$ coincide

common contour \curvearrowright


Residues

$$a_{-1} = \frac{1}{2\pi i} \oint_C f(s) ds \quad \rightarrow \quad \oint_C f(s) ds = 2\pi i \cdot a_{-1}$$

$$a_{-1} = \frac{1}{2\pi i} \oint_C f(s) ds = \text{Res}(f(z), z_0)$$

$$= \begin{cases} \lim_{z \rightarrow z_0} (z - z_0) f(z) & \text{(simple)} \\ \frac{1}{(n-1)!} \lim_{z \rightarrow z_0} \frac{d^{n-1}}{dz^{n-1}} (z - z_0)^n f(z) & \text{(order } n) \end{cases}$$

Series Expansion at $z=0$




$$f(z) = \sum_{n=\eta_1}^{\infty} a_n^{(mf)} z^n$$

$$a_n^{(mf)} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{z^{n+1}} dz$$
$$= \sum_k \text{Res} \left(\frac{f(z)}{z^{n+1}}, z_k \right)$$


Poles z_k

$$n \geq 0 \quad z_1, z_2, z_3, \circ$$


$$n < 0 \quad z_1, z_2, z_3$$




$\text{Res}(f, z_m)$




$\text{Res}(f, z_1)$
 $\text{Res}(f, z_m)$




$\text{Res}(f, z_1) + \text{Res}(f, z_2) + \text{Res}(f, z_m)$




$\text{Res}(f, z_1) + \text{Res}(f, z_2) + \text{Res}(f, z_3)$
 $+ \text{Res}(f, z_m)$




$$\text{Res}(f, z_m) \quad (n \geq 0)$$



$$\begin{aligned} &\text{Res}(f, z_1) \\ &\text{Res}(f, z_m) \quad (n \geq 0) \end{aligned}$$



$$\begin{aligned} &\text{Res}(f, z_1) + \text{Res}(f, z_2) + \text{Res}(f, z_m) \\ &\quad (n \geq 0) \end{aligned}$$



$$\begin{aligned} &\text{Res}(f, z_1) + \text{Res}(f, z_2) + \text{Res}(f, z_3) \\ &+ \text{Res}(f, z_m) \quad (n \geq 0) \end{aligned}$$

Inverse z-Transform $x[n] = \frac{1}{2\pi i} \int_C X(z) z^{n-1} dz$

$$X(z) = \sum_{k=0}^{\infty} x_k z^{-k}$$

$$z^{n-1} X(z) = \left(\sum_{k=0}^{\infty} x_k z^{-k} \right) z^{n-1}$$

$$\int z^{n-1} \text{LHS} dz = \int \text{RHS} z^{n-1} dz$$

$$= \sum_{k=0}^{\infty} x_k z^{-k+n-1}$$

$$[0, \infty) = [0, n-1] \cup [n] \cup [n+1, \infty)$$

$$= \sum_{k=0}^{n-1} x_k z^{-k+n-1} + \sum_{k=n}^n x_k z^{-k+n-1} + \sum_{k=n+1}^{\infty} x_k z^{-k+n-1}$$

$$= \sum_{k=0}^{n-1} x_k z^{-k+n-1} + \frac{x_n}{z^1} + \sum_{k=n+1}^{\infty} \frac{x_k}{z^{k-n+1}}$$

$$\int_C X(z) z^{n-1} dz = \int_C \sum_{k=0}^{n-1} x_k z^{-k+n-1} dz + \int_C \frac{x_n}{z^1} dz + \int_C \sum_{k=n+1}^{\infty} \frac{x_k}{z^{k-n+1}} dz$$

$$= \sum_{k=0}^{n-1} x_k \int_C z^{-k+n-1} dz + x_n \int_C \frac{1}{z^1} dz + \sum_{k=n+1}^{\infty} x_k \int_C \frac{1}{z^{k-n+1}} dz$$

$$= \sum_{k=0}^{n-1} x_k \cdot 0 + x_n \cdot 2\pi i + \sum_{k=n+1}^{\infty} x_k \cdot 0$$

$$x[n] = \frac{1}{2\pi i} \int_C X(z) z^{n-1} dz$$

Z-transform

$$z_m = 0$$

$$\begin{aligned}
 x[n] &= \frac{1}{2\pi i} \oint_C f(z) z^{n-1} dz \\
 &= \sum_k \operatorname{Res}(f(z) z^{n-1}, z_k)
 \end{aligned}$$

$n > 0$ z_k : poles of $f(z)$

$n = 0$ z_k : poles of $f(z)$ + $z = 0$
 $z^{n-1} = z^{-1} = \frac{1}{z}$

$x[n]$ includes $u[n] \rightarrow X[z]$ contains z on its numerator

Also, think about modified partial fraction $\frac{X[z]}{z}$

Laurent Expansion

expansion at z_m

$$\begin{aligned}
 a_n^{\{m\}} &= \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z-z_m)^{n+1}} dz \\
 &= \sum_k \operatorname{Res}\left(\frac{f(z)}{(z-z_m)^{n+1}}, z_k\right)
 \end{aligned}$$





$$z_m = 0$$

$$\begin{aligned}
 a_n^{\{0\}} &= \frac{1}{2\pi i} \oint_C \frac{f(z)}{z^{n+1}} dz \\
 &= \sum_k \operatorname{Res}\left(\frac{f(z)}{z^{n+1}}, z_k\right)
 \end{aligned}$$

$$\begin{aligned}
 a_{-n}^{\{0\}} &= \frac{1}{2\pi i} \oint_C f(z) z^{n-1} dz \\
 &= \sum_k \operatorname{Res}(f(z) z^{n-1}, z_k)
 \end{aligned}$$


$$\begin{aligned}
 a_{-n}^{\{0\}} &= \frac{1}{2\pi i} \oint_C \frac{f(z)}{z^{-n+1}} dz \\
 &= \sum_k \operatorname{Res}\left(\frac{f(z)}{z^{-n+1}}, z_k\right)
 \end{aligned}$$

Different D, Different Laurent Series



$$x[n] = \frac{1}{2\pi i} \int_C X(z) z^{n-1} dz$$

$$= \sum_{z_k} \text{Res}(X(z) z^{n-1}, z_k)$$



z-transform

$$f(z) = \frac{-1}{(z-1)(z-2)}$$

Complex Variables and Ap
Brown & Churchill

$$f(z) = \frac{-1}{(z-1)(z-2)} = \frac{1}{z-1} - \frac{1}{z-2}$$

$$D_1: |z| < 1$$

$$D_2: 1 < |z| < 2$$

$$D_3: 2 < |z|$$

$$\textcircled{1} D_1 \quad |z| < 1, \quad \left|\frac{z}{2}\right| < 1$$

$$\begin{aligned} f(z) &= \frac{1}{z-1} - \frac{1}{z-2} = \frac{-1}{1-z} + \frac{1}{1-\left(\frac{z}{2}\right)} \\ &= -\sum_{n=0}^{\infty} z^n + \sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} = \sum_{n=0}^{\infty} (2^{-n-1} - 1)z^n \quad |z| < 1 \end{aligned}$$

$$\textcircled{2} D_2 \quad 1 < |z| < 2 \Rightarrow \left|\frac{1}{z}\right| < 1, \quad \left|\frac{z}{2}\right| < 1$$


$$\begin{aligned} f(z) &= \frac{1}{z-1} - \frac{1}{z-2} = \frac{1}{z} \cdot \frac{1}{1-\left(\frac{1}{z}\right)} + \frac{1}{2} \frac{1}{1-\left(\frac{z}{2}\right)} \\ &= \sum_{n=0}^{\infty} \frac{1}{z^{n+1}} + \sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} \\ &= \sum_{n=1}^{\infty} \frac{1}{z^n} + \sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} \end{aligned}$$

$$\textcircled{3} D_3 \quad 2 < |z| \quad \left|\frac{z}{2}\right| < 1 \quad \left|\frac{1}{z}\right| < 1$$

$$\begin{aligned} f(z) &= \frac{1}{z-1} - \frac{1}{z-2} = \frac{1}{z} \frac{1}{1-\left(\frac{1}{z}\right)} - \frac{1}{z} \frac{1}{1-\left(\frac{z}{2}\right)} \\ &= \sum_{n=0}^{\infty} \frac{1}{z^{n+1}} - \sum_{n=0}^{\infty} \frac{z^n}{z^{n+1}} = \sum_{n=0}^{\infty} \frac{1-2^n}{z^{n+1}} \\ &= \sum_{n=1}^{\infty} \frac{1-2^{n-1}}{z^n} \end{aligned}$$

$$f(z) = \frac{-1}{(z-1)(z-2)}$$

① $D_1 \quad |z| < 1, \quad \left|\frac{z}{2}\right| < 1$




$$\frac{f(z)}{z^{n+1}} = \frac{-1}{(z-1)(z-2)z^{n+1}}$$

$$\begin{aligned} f(z) &= \frac{1}{z-1} - \frac{1}{z-2} = \frac{-1}{1-z} + \frac{1}{1-\left(\frac{z}{2}\right)} \\ &= -\sum_{n=0}^{\infty} z^n + \sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} = \sum_{n=0}^{\infty} (2^{-n-1} - 1)z^n \quad |z| < 1 \end{aligned}$$

$$a_n = \frac{f(z)}{z^{n+1}} = \frac{1}{(z-1)(z-2)z^{n+1}} \quad \frac{1}{z-1} - \frac{1}{z-2}$$

$$a_n = \sum_{k=1}^M \operatorname{Res} \left(\frac{f(z)}{(z-z_k)^{n+1}}, z_k \right) = \operatorname{Res} \left(\frac{-1}{(z-1)(z-2)z^{n+1}}, 0 \right)$$



$$a_n = \sum_{k=1}^M \operatorname{Res} \left(\frac{f(z)}{(z-z_k)^{n+1}}, z_k \right) = \operatorname{Res} \left(\frac{-1}{(z-1)(z-2)z^{n+1}}, 0 \right)$$

$n \geq 0$ then the pole $z=0$

$$\frac{1}{(n-1)!} \lim_{z \rightarrow z_0} \frac{d^{n-1}}{dz^{n-1}} (z-z_0)^n f(z) \quad (\text{order } n)$$

$$\frac{d}{dz} ((z-1)^{-1} - (z-2)^{-1}) = (-1) ((z-1)^{-2} - (z-2)^{-2})$$

$$\frac{d^2}{dz^2} ((z-1)^{-1} - (z-2)^{-1}) = (-1)(-2) ((z-1)^{-3} - (z-2)^{-3})$$

$$\frac{d^3}{dz^3} ((z-1)^{-1} - (z-2)^{-1}) = (-1)(-2)(-3) ((z-1)^{-4} - (z-2)^{-4})$$

$$\frac{d^n}{dz^n} ((z-1)^{-1} - (z-2)^{-1}) = (-1)^n n! ((z-1)^{-n-1} - (z-2)^{-n-1})$$


$$\begin{aligned} \frac{1}{n!} \lim_{z \rightarrow 0} \frac{d^n}{dz^n} ((z-1)^{-1} - (z-2)^{-1}) &= (-1)^n \lim_{z \rightarrow 0} ((z-1)^{-n-1} - (z-2)^{-n-1}) \\ &= (-1)^n ((-1)^{-n-1} - (-2)^{-n-1}) \\ &= -1 + 2^{-n-1} \end{aligned}$$

$$a_n = -1 + 2^{-n-1} \quad (n \geq 0)$$

$$f(z) = \sum_{n=-n_1}^{\infty} a_n z^n = \sum_{n=0}^{\infty} (2^{-n-1} - 1) z^n$$


$$f(z) = \frac{-1}{(z-1)(z-2)}$$

$$\textcircled{2} \quad D_2 \quad 1 < |z| < 2 \Rightarrow \left| \frac{1}{z} \right| < 1, \quad \left| \frac{z}{2} \right| < 1$$



$$\begin{aligned} f(z) &= \frac{1}{z-1} - \frac{1}{z-2} = \frac{1}{z} \cdot \frac{1}{1-\left(\frac{1}{z}\right)} + \frac{1}{2} \frac{1}{1-\left(\frac{z}{2}\right)} \\ &= \sum_{n=0}^{\infty} \frac{1}{z^{n+1}} + \sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} \\ &= \sum_{n=1}^{\infty} \frac{1}{z^n} + \sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} \end{aligned}$$

$$\begin{aligned} a_n &= \sum_{k=1}^M \operatorname{Res} \left(\frac{f(z)}{(z-z_k)^{n+1}}, z_k \right) = \operatorname{Res} \left(\frac{-1}{(z-1)(z-2)z^{n+1}}, 0 \right) \\ &\quad + \operatorname{Res} \left(\frac{-1}{(z-1)(z-2)z^{n+1}}, 1 \right) \end{aligned}$$



$$a_n = \sum_{k=1}^M \operatorname{Res} \left(\frac{f(z)}{(z-z_k)^{n+1}}, z_k \right) = \operatorname{Res} \left(\frac{-1}{(z-1)(z-2)z^{n+1}}, 0 \right) + \operatorname{Res} \left(\frac{-1}{(z-1)(z-2)z^{n+1}}, 1 \right)$$

$$\frac{1}{(n-1)!} \lim_{z \rightarrow z_0} \frac{d^{n-1}}{dz^{n-1}} (z-z_0)^n f(z) \quad (\text{order } n)$$

$$\begin{aligned} \frac{1}{n!} \lim_{z \rightarrow 0} \frac{d^n}{dz^n} ((z-1)^{-1} - (z-2)^{-1}) &= (-1)^n \lim_{z \rightarrow 0} ((z-1)^{-n-1} - (z-2)^{-n-1}) \\ &= (-1)^n ((-1)^{-n-1} - (-2)^{-n-1}) \\ &= -1 + 2^{-n-1} \end{aligned}$$

$$\operatorname{Res} \left(\frac{-1}{(z-1)(z-2)z^{n+1}}, 0 \right) = -1 + 2^{-n-1} \quad (n \geq 0)$$

$$\operatorname{Res} \left(\frac{-1}{(z-1)(z-2)z^{n+1}}, 1 \right) = \lim_{z \rightarrow 1} (z-1) \frac{-1}{(z-1)(z-2)z^{n+1}} = 1$$


| $n=-3$ | $n=-2$ | $n=-1$ | $n=0$ | $n=1$ | $n=2$ | |
|--------|--------|--------|-------------|-------------|-------------|---|
| 0 | 0 | 0 | $-1+2^{-1}$ | $-1+2^{-2}$ | $-1+2^{-3}$ | $\operatorname{Res} \left(\frac{f(z)}{z^{n+1}}, 0 \right)$ |
| 1 | 1 | 1 | 1 | 1 | 1 | $\operatorname{Res} \left(\frac{f(z)}{z^{n+1}}, 1 \right)$ |
| 1 | 1 | 1 | 2^{-1} | 2^{-2} | 2^{-3} | |

$$\begin{cases} a_n = 2^{-n-1} & n \geq 0 \\ a_n = 1 & n < 0 \end{cases} \quad \begin{cases} 2^{-n-1} z^n \\ z^{-n} \end{cases}$$

$$f(z) = \sum_{n=1}^{\infty} \frac{1}{z^n} + \sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}}$$


$$f(z) = \frac{-1}{(z-1)(z-2)}$$

③ $D_3 \quad 2 < |z| \quad \left| \frac{2}{z} \right| < 1 \quad \left| \frac{1}{z} \right| < 1$



$$\begin{aligned} f(z) &= \frac{1}{z-1} - \frac{1}{z-2} = \frac{1}{z} \frac{1}{1-(\frac{1}{z})} - \frac{1}{z} \frac{1}{1-(\frac{2}{z})} \\ &= \sum_{n=0}^{\infty} \frac{1}{z^{n+1}} - \sum_{n=0}^{\infty} \frac{2^n}{z^{n+1}} = \sum_{n=0}^{\infty} \frac{1-2^n}{z^{n+1}} \\ &= \sum_{n=1}^{\infty} \frac{1-2^{n-1}}{z^n} \end{aligned}$$

$$\begin{aligned} a_n &= \sum_{k=1}^M \operatorname{Res} \left(\frac{f(z)}{(z-z_k)^{n+1}}, z_k \right) = \operatorname{Res} \left(\frac{-1}{(z-1)(z-2)z^{n+1}}, 0 \right) \\ &\quad + \operatorname{Res} \left(\frac{-1}{(z-1)(z-2)z^{n+1}}, 1 \right) \\ &\quad + \operatorname{Res} \left(\frac{-1}{(z-1)(z-2)z^{n+1}}, 2 \right) \end{aligned}$$



$$\text{Res} \left(\frac{-1}{(z-1)(z-2)z^{n+1}}, 0 \right) = -1 + 2^{-n+1} \quad (n \geq 0)$$

$$\text{Res} \left(\frac{-1}{(z-1)(z-2)z^{n+1}}, 1 \right) = \lim_{z \rightarrow 1} (z-1) \frac{-1}{(z-1)(z-2)z^{n+1}} = 1$$


$$\text{Res} \left(\frac{-1}{(z-1)(z-2)z^{n+1}}, 2 \right) = \lim_{z \rightarrow 2} (z-2) \frac{-1}{(z-1)(z-2)z^{n+1}} = -\frac{1}{2^{n+1}}$$

| $n=-3$ | $n=-2$ | $n=-1$ | $n=0$ | $n=1$ | $n=2$ | |
|---------|--------|--------|----------|----------|----------|---|
| 0 | 0 | 0 | $-1+2^1$ | $-1+2^2$ | $-1+2^3$ | $\text{Res} \left(\frac{f(z)}{z^{n+1}}, 0 \right)$ |
| 1 | 1 | 1 | 1 | 1 | 1 | $\text{Res} \left(\frac{f(z)}{z^{n+1}}, 1 \right)$ |
| -2^2 | -2 | -1 | -2^1 | -2^2 | -2^3 | $\text{Res} \left(\frac{f(z)}{z^{n+1}}, 2 \right)$ |
| $1-2^2$ | $1-2$ | 0 | 0 | 0 | 0 | |

$$a_n = 1 - 2^{-n+1} \quad n < 0 = \sum_{n=1}^{\infty} \frac{1-2^{n+1}}{z^n}$$

$$f(z) = \sum_{n=-1}^{-\infty} (1-2^{-n+1}) z^n = \sum_{n=1}^{\infty} \frac{1-2^{n+1}}{z^n}$$

$$f(z) = \frac{-1}{(z-1)(z-2)}$$



$$x[n]$$

$$= \frac{1}{2\pi i} \int_C \boxed{X(z) z^{n-1}} dz$$

$$= \sum_{j=1}^k \text{Res}(\boxed{X(z) z^{n-1}}, z_j)$$


$$X(z) = \frac{-1}{(z-1)(z-2)}$$

$$X(z) z^{n-1} = \frac{-1}{(z-1)(z-2)} z^{n-1}$$

$$\text{Res}(\boxed{X(z) z^{n-1}}, 1) = (z-2) \frac{-1}{(z-1)(z-2)} z^{n-1} \Big|_{z=1} = 1$$

$$\text{Res}(\boxed{X(z) z^{n-1}}, 2) = (z-1) \frac{-1}{(z-1)(z-2)} z^{n-1} \Big|_{z=2} = -2^{n-1}$$

$$x[n] = 1 - 2^{n-1}$$



ROC (Region of Convergence)

$$|z| > 2 \Rightarrow \frac{2}{|z|} < 1$$

$$\left(\frac{2}{z}\right)^0 + \left(\frac{2}{z}\right)^1 + \left(\frac{2}{z}\right)^2 + \dots \longrightarrow \frac{1}{1 - \frac{2}{z}}$$

Converge

$$|z| > 2 \Rightarrow \frac{1}{|z|} < 1$$

$$\left(\frac{1}{z}\right)^0 + \left(\frac{1}{z}\right)^1 + \left(\frac{1}{z}\right)^2 + \dots \longrightarrow \frac{1}{1 - \frac{1}{z}}$$

Converge

$$\begin{aligned} f(z) &= \frac{1}{z-1} - \frac{1}{z-2} = \frac{1}{z} \frac{1}{1 - (\frac{1}{z})} - \frac{1}{z} \frac{1}{1 - (\frac{2}{z})} \\ &= \sum_{n=0}^{\infty} \frac{1}{z^{n+1}} - \sum_{n=0}^{\infty} \frac{2^n}{z^{n+1}} = \sum_{n=0}^{\infty} \frac{1 - 2^n}{z^{n+1}} \\ &= \sum_{n=1}^{\infty} \frac{1 - 2^{n-1}}{z^n} \end{aligned}$$

$$\left(\frac{1}{z}\right) + \left(\frac{1}{z}\right)^2 + \left(\frac{1}{z}\right)^3 + \dots + \frac{1}{z} \left\{ \left(\frac{2}{z}\right) + \left(\frac{2}{z}\right)^2 + \left(\frac{2}{z}\right)^3 + \dots \right\} \longrightarrow \frac{1}{z-1} - \frac{1}{z-2} = \frac{-1}{(z-1)(z-2)}$$

Converge

$$(1-2^0)z^1 + (1-2^1)z^2 + (1-2^2)z^3 + \dots \longrightarrow \frac{-1}{(z-1)(z-2)} \quad (|z| > 2)$$

Converge

$$x[n] = 1 - 2^n \quad \longleftrightarrow \quad X(z) = \frac{-1}{(z-1)(z-2)} \quad (|z| > 2)$$





