

# Residue Integrals and Laurent Series with non-annular region

20170217

Copyright (c) 2016 - 2017 Young W. Lim.

Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.2 or any later version published by the Free Software Foundation; with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts. A copy of the license is included in the section entitled "GNU Free Documentation License".

Based on

T.J. Cavicchi, Digital Signal Processing

Complex Analysis for Mathematics and Engineering  
J. Mathews

# Residue Theorem

D: Simply connected domain

C: Simple closed contour (CCW) in D


if  $f(z)$  is **analytic** inside C and on C  
except at the points  $z_1, z_2, \dots, z_k$  in C

then


$$\frac{1}{2\pi i} \int_C f(z) dz = \sum_{j=1}^k \text{Res}(f(z), z_j)$$

Singular points of  $f(z)$  :  $z_1, z_2, \dots, z_k$


D



C




C



# Integration of a function of a complex var.


$$\oint_C f(z) dz = 2\pi i \sum_{k=1}^n \text{Res}(f(z), z_k)$$

finite number  $k$  of  
singular points  $z_k$   
residue theorem



$$\oint_C f(z) dz = 0 \quad \text{if } f(z) \text{ is analytic within and on } C$$

no singularity



$$\oint_C f(z) dz = 0 \quad \text{if } f(z) = F'(z) \text{ on } C$$

:  $F(z)$  is an antiderivative of  $f(z)$   
fundamental theorem of calculus

$\oint_C f(z) dz = 0$  if  $f(z)$  is continuous in  $D$  and  
 $f(z) = F'(z)$  :  $F(z)$  is an antiderivative of  $f(z)$   
fundamental theorem of calculus

# Series Expansion

can expand  $f(z)$  about any point  $z_m$   
over powers of  $(z - z_m)$

whether or not  $f(z)$  is singular at  $z_m$   
or at other points between  $z$  and  $z_m$

$$f(z) = \sum_{n=\eta_1}^{\infty} a_n^{(m)} (z - z_m)^n$$

① Laurent Series Expansion of  $f(z)$  at  $z_m$   
general  $\eta_1$  - depend on  $f(z)$  and  $z_m$

②  $z$ -transform of  $a_n^{(m)}$   
general  $\eta_1$  - depend on  $f(z)$

$$z_m = 0$$

③ Taylor Series Expansion of  $f(z)$  at  $z_m$   
positive  $\eta_1$  - depend on  $f(z)$  and  $z_m$  ( $\eta_1 > 0$ )

④ MacLaurin Series Expansion of  $f(z)$  at  $z_m$   
positive  $\eta_1$  - depend on  $f(z)$  ( $\eta_1 > 0$ )

$$z_m = 0$$

$$f(z) = \sum_{n=n_1}^{\infty} a_n^{(m)} (z - z_m)^n$$

$n_1 > 0$  pos powers

$$z_m = 0$$

① Laurent Series

③ Taylor Series

② z-transform

④ MacLaurin Series

\* Expansion of  $f(z)$  about any point  $z_m$   
over powers of  $(z - z_m)$

$$f(z) = \sum_{n=n_1}^{\infty} a_n^{(m)} (z - z_m)^n$$

$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz$$
$$a_n^{(m)} = \sum_k \operatorname{Res} \left( \frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

for general  $f(z)$

for general  $f(z)$

$$a_n^{(m)} = \frac{1}{n!} f^{(n)}(z_m) \quad n_1 \geq 0$$

for analytic  $f(z)$  within  $C$

analytic  $f(z) \longrightarrow \frac{f(z)}{(z - z_m)^{n+1}}$  has a pole at  $z_m$   
order of  $n+1$



$$f(z) = \sum_{n=n_1}^{\infty} a_n^{(m)} (z - z_m)^n$$

$z_m$ : possible poles of  $f(z)$   
not necessarily poles

$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z')}{(z' - z_m)^{n+1}} dz'$$
$$= \sum_k \text{Res} \left( \frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$




$z_k$ : poles of  $\frac{f(z)}{(z - z_m)^{n+1}}$   
within  $C$

$$= \frac{1}{n!} f^{(n)}(z_m) \quad n_1 \geq 0$$

# Residue Theorem and Laurent Series

assumed there are  $(K)$  singularities (poles) of  $f(z)$  in a region


let  $\{C_k\}$  be taken to enclose only one pole  $z_k$



$a_n^{(1)}$  expanded at  $z_1$

$C_1$  encloses  $z_1$  only


$$\tilde{a}_{-1}^{(1)} = \text{Res}(f(z), z_1)$$



$a_n^{(2)}$  expanded at  $z_2$

$C_2$  encloses  $z_2$  only

$$\tilde{a}_{-1}^{(2)} = \text{Res}(f(z), z_2)$$



$a_n^{(3)}$  expanded at  $z_3$

$C_3$  encloses  $z_3$  only

$$\tilde{a}_{-1}^{(3)} = \text{Res}(f(z), z_3)$$

# Cauchy's Residue Theorem



$f(z)$ : **analytic** on and within  $C$   
except a finite number of **singular points**  
 $z_1, z_2, \dots, z_n$  within  $C$

then

$$\int_C f(z) dz = 2\pi i \sum_{k=1}^n \text{Res}(f(z), z_k)$$

$D$ : a simply connected domain

$C$ : a simple closed contour in  $D$



$C_1$   $(z_1)$

$$f(z) = \sum_{k=-\infty}^{+\infty} a_k (z - z_1)^k$$

$$a_{-1}^{(1)} = \frac{1}{2\pi i} \oint_{C_1} f(s) ds = \text{Res}(f(z), z_1)$$

$C_2$   $(z_2)$

$$f(z) = \sum_{k=-\infty}^{+\infty} a_k (z - z_2)^k$$

$$a_{-1}^{(2)} = \frac{1}{2\pi i} \oint_{C_2} f(s) ds = \text{Res}(f(z), z_2)$$

$C_3$   $(z_3)$

$$f(z) = \sum_{k=-\infty}^{+\infty} a_k (z - z_3)^k$$

$$a_{-1}^{(3)} = \frac{1}{2\pi i} \oint_{C_3} f(s) ds = \text{Res}(f(z), z_3)$$

# Laurent Series with Annular Region expanded at each pole of $f(z)$

$z_1$  Laurent series expansion at  $z_1$


$$f(z) = \sum_{n=-\infty}^{+\infty} a_n^{(1)} (z - z_1)^n$$



$$\begin{aligned} \tilde{a}_{-1}^{(1)} &= \text{Res}(f(z), z_1) \\ &= \frac{1}{2\pi i} \oint_{C_1} f(z) dz \end{aligned}$$

$z_2$  Laurent series expansion at  $z_2$


$$f(z) = \sum_{n=-\infty}^{+\infty} a_n^{(2)} (z - z_2)^n$$




$$\begin{aligned} \tilde{a}_{-1}^{(2)} &= \text{Res}(f(z), z_2) \\ &= \frac{1}{2\pi i} \oint_{C_2} f(z) dz \end{aligned}$$

$z_3$  Laurent series expansion at  $z_3$

$$f(z) = \sum_{n=-\infty}^{+\infty} a_n^{(3)} (z - z_3)^n$$




$$\begin{aligned} \tilde{a}_{-1}^{(3)} &= \text{Res}(f(z), z_3) \\ &= \frac{1}{2\pi i} \oint_{C_3} f(z) dz \end{aligned}$$



$$\int_C f(z) dz = 2\pi i \sum_{k=1}^n \text{Res}(f(z), z_k)$$

# Residue Theorem + Laurent Series

\* Whether  $z_m$  is singular or not






$$f(z) = \sum_{n=n_1}^{\infty} a_n^{(m)} (z - z_m)^n$$

$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz$$

$$= \sum_k \text{Res} \left( \frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

$$a_{-1}^{(m)} = \frac{1}{2\pi i} \oint_C f(z) dz$$

$$= \sum_k \text{Res} (f(z), z_k)$$



$$\tilde{a}_{-1}^{(1)} = \text{Res}(f(z), z_1)$$

$$\tilde{a}_{-1}^{(2)} = \text{Res}(f(z), z_2)$$

$$\tilde{a}_{-1}^{(3)} = \text{Res}(f(z), z_3)$$

$$a_{-1}^{(m)} = \tilde{a}_{-1}^{(1)} + \tilde{a}_{-1}^{(2)} + \tilde{a}_{-1}^{(3)}$$

$$a_{-1}^{(m)} = \text{Res}(f(z), z_1) + \text{Res}(f(z), z_2) + \text{Res}(f(z), z_3)$$

← Laurent Series coefficient  $a_{-1}^{(2)}$

- singular center  $z_i$
- punctured open disk

This cannot be a residue because it is not

isolated singular center nor punctured open disk

Laurent Series — Annular Region of Convergence  
 — no singularity in this region


Residue — Laurent Series expanded at a pole

a punctured open disk

- Annular
- Isolated Singularity




$$\tilde{a}_{-1}^{\{1\}} = \text{Res}(f(z), z_1)$$




$$\tilde{a}_{-1}^{\{2\}} = \text{Res}(f(z), z_2)$$



$$\tilde{a}_{-1}^{\{3\}} = \text{Res}(f(z), z_3)$$



$$\tilde{a}_{-1}^{\{m\}} = \text{Res}(f(z), z_m)$$



# Computing $a_n^{\{m\}}$

$$f(z) = \sum_{n=\nu_1}^{\infty} a_n^{\{m\}} (z - z_m)^n \quad \boxed{n \leftarrow k}$$

$$f(z) = \sum_{k=\nu_1}^{\infty} a_k^{\{m\}} (z - z_m)^k$$

for a given  $n$

$$\frac{f(z)}{(z - z_m)^{n+1}} = \sum_{k=\nu_1}^{\infty} a_k^{\{m\}} (z - z_m)^{k-n-1} \quad \begin{array}{l} k: \text{index variable} \\ n: \text{fixed value} \end{array}$$

$$\oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz = \oint_C \sum_{k=\nu_1}^{\infty} a_k^{\{m\}} (z - z_m)^{k-n-1} dz$$

$$= \sum_{k=\nu_1}^{\infty} \oint_C a_k^{\{m\}} (z - z_m)^{k-n-1} dz \quad \boxed{k=n}$$

$$\oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz = \oint_C a_n^{\{m\}} \frac{1}{(z - z_m)} dz = 2\pi i \cdot a_n^{\{m\}}$$

$$a_n^{\{m\}} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz$$

$$\oint_C \left[ \dots (z - z_m)^{-3} + (z - z_m)^{-2} + \frac{1}{(z - z_m)} + 1 + (z - z_m) + (z - z_m)^2 + \dots \right] dz$$

$$= \oint_C \frac{1}{(z - z_m)} dz = 2\pi i$$

# Computing $a_n^{\{m\}}$ using Residues

expansion at  $z_m$

$$\eta = -1 \quad \eta + 1 = 0 \quad (z - z_m)^{\eta+1} = 1$$

$$a_n^{\{m\}} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz$$


$$= \sum_k \text{Res} \left( \frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

$$a_{-1}^{\{m\}} = \frac{1}{2\pi i} \oint_C f(z) dz$$

$$= \sum_k \text{Res} (f(z), z_k)$$

$$a_{-1}^{\{m\}} = \frac{1}{2\pi i} \oint_C f(z) dz = \sum_k \text{Res} (f(z), z_k)$$

$$a_{-1}^{\{m\}} = \cancel{\text{Res} (f(z), z_m)}$$



$$f(z) = \sum_{n=n_1}^{\infty} a_n^{\{m\}} (z - z_m)^n$$

$$a_n^{\{m\}} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz$$

$$= \sum_k \text{Res} \left( \frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

Residue  $\rightarrow$  Laurent series  $\rightarrow$  annular region  $\rightarrow$  expanded at a pole  $\star$  ) a punctured open disk



$$f(z) = \sum_{n=n_1}^{\infty} a_n^{(m)} (z - z_m)^n$$

$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz$$
$$= \sum_k \operatorname{Res} \left( \frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

$$a_{-1}^{(m)} = \frac{1}{2\pi i} \oint_C f(z) dz$$
$$= \sum_k \operatorname{Res} (f(z), z_k)$$

⋮

$$a_{-3}^{(m)} = \sum_k \operatorname{Res} (f(z)(z - z_m)^2, z_k)$$

$$a_{-2}^{(m)} = \sum_k \operatorname{Res} (f(z)(z - z_m)^1, z_k)$$

$$a_{-1}^{(m)} = \sum_k \operatorname{Res} (f(z), z_k)$$

$$a_0^{(m)} = \sum_k \operatorname{Res} \left( \frac{f(z)}{(z - z_m)^1}, z_k \right)$$

$$a_1^{(m)} = \sum_k \operatorname{Res} \left( \frac{f(z)}{(z - z_m)^2}, z_k \right)$$

$$a_2^{(m)} = \sum_k \operatorname{Res} \left( \frac{f(z)}{(z - z_m)^3}, z_k \right)$$

⋮

# Poles for Residue Computation

$$f(z) = \sum_{n=\eta_1}^{\infty} a_n^{(m)} (z - z_m)^n$$

$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz$$

$$= \sum_k \operatorname{Res} \left( \frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

$z_k$  within  $C$  : singularities of

$$\frac{f(z)}{(z - z_m)^{n+1}}$$


Ⓘ non-singular  $z_m$

$$\begin{array}{lll} n \geq 0 & \{ \text{poles of } f(z) \} \cup \{ z_m \} & n = 0, 1, 2, \dots \\ n < 0 & \{ \text{poles of } f(z) \} & n = -1, -2, \dots \end{array}$$


Ⓡ singular  $z_m$

$$\begin{array}{ll} n \geq 0 & \{ \text{poles of } f(z) \} \\ n < 0 & \{ \text{poles of } f(z) \} \end{array}$$


↖  $z_m$  included




$$\int_C f(z) dz = 2\pi i \operatorname{Res}(f(z), z_1) + 2\pi i \operatorname{Res}(f(z), z_2) + 2\pi i \operatorname{Res}(f(z), z_3)$$



$$\int_{C'} f(z) dz = 2\pi i \operatorname{Res}(f(z), z_1) + 2\pi i \operatorname{Res}(f(z), z_2)$$




$$\int_{C''} f(z) dz = 2\pi i \operatorname{Res}(f(z), z_1)$$



$$\int_{C'''} f(z) dz = 0$$

# Series Expansion at $z_m$



$$f(z) = \sum_{n=n_1}^{\infty} a_n^{[m]} (z - z_m)^n$$

$$a_n^{[m]} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz$$


$$= \sum_k \text{Res} \left( \frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

$$a_{-1}^{[m]} = \frac{1}{2\pi i} \oint_C f(z) dz$$

$$= \sum_k \text{Res} (f(z), z_k)$$

$$a_{-1}^{[m]} \neq \text{Res}(f(z), z_m)$$

## [Annular Region]



$$a_{-1}^{[m]} \neq \text{Res}(f(z), z_m)$$

\* for a nonsingular  $z_m$   
 $z_m$  can be a pole of


$$\frac{f(z)}{(z - z_m)^{n+1}} \quad \text{if } n \geq 0$$

When computing

$$a_n^{[m]} = \sum_k \text{Res} \left( \frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

[Annular Region] & [ $z_m$  : isolated singularity]

A punctured open disk



$$f(z) = \sum_{n=n_1}^{\infty} a_n^{(m)} (z - z_m)^n$$

$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz$$


$$= \sum_k \text{Res} \left( \frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

$$a_{-1}^{(m)} = \frac{1}{2\pi i} \oint_C f(z) dz$$

$$= \sum_k \text{Res} (f(z), z_k)$$

$$a_{-1}^{(m)} = \text{Res} (f(z), z_m)$$

# Series Expansion at $z=0$



$$f(z) = \sum_{n=n_1}^{\infty} a_n^{(m)} z^n$$

$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{z^{n+1}} dz$$
$$= \sum_k \text{Res} \left( \frac{f(z)}{z^{n+1}}, z_k \right)$$


Poles  $z_k$

$$n \geq 0 \quad z_1, z_2, z_3, \circ$$

$$n < 0 \quad z_1, z_2, z_3$$

# A punctured open disk

if  $C$  encloses only one pole  $z_0$ ,  
and the expansion at that pole  $z_0$  is assumed,  
then



$$\boxed{a_{-1}^{\{0\}}} = \frac{1}{2\pi i} \oint_{C_0} f(z) dz = \text{Res}(f(z), z_0)$$

Let



$$\boxed{\tilde{a}_{-1}^{\{m\}} = \text{Res}(f(z), z_m)}$$

notation  $\sim$

the residue of  $f(z)$  at  $z_m$

using  $C_m$  which is in the punctured open disk ROC

$$\boxed{f(z) = \sum_{n=-\infty}^{+\infty} a_n^{\{m\}} (z - z_m)^n}$$



$$\oint_C f(z) dz = 2\pi j \sum_{k=1}^M \tilde{a}_{-1}^{(k)} = 2\pi j \sum_{k=1}^M \text{Res}(f(z), z_k)$$

residue theorem

$$a_n = \sum_{k=1}^M \text{Res} \left( \frac{f(z)}{(z-z_k)^{n+1}}, z_k \right)$$

Laurent coefficient

$C$  encloses  $k$  poles

$C_k$  encloses only the  $k$ -th pole


$\tilde{a}_{-1}^{(k)}$  the residue of the  $k$ -th pole enclosed by  $C$ ,  $z_k$



# Non-annular region

$$f(z) = \sum_{n=n_1}^{\infty} a_n^{f, m} (z - z_m)^n$$

$$a_n^{f, m} = \frac{1}{2\pi i} \oint_C \frac{f(z')}{(z' - z_m)^{n+1}} dz'$$
$$= \sum_k \operatorname{Res} \left( \frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$



$C$  is in the same region of analyticity of  $f(z)$

typically a circle centered on  $z_m$

non-annular ok

$z_k$  within  $C$  : singularities of  $\frac{f(z)}{(z - z_m)^{n+1}}$

$n_1 = n_{f, m}$  depends on  $f(z)$ ,  $z_m$

$a_n^{f, m}$  depends on  $f(z)$ ,  $z_m$ , region of analyticity

Whether  $f(z)$  is singular at  $z = z_m$  or not

or at other points between  $z$  and  $z_m$

We can expand  $f(z)$  about any point  $z_m$

over powers of  $(z - z_m)$ .

# Laurent's Theorem

$f$  : analytic within the **annular** domain  $D$

$$r < |z - z_0| < R$$

then

$$f(z) = \sum_{k=-\infty}^{+\infty} a_k (z - z_0)^k,$$

valid for  $r < |z - z_0| < R$

The coefficients  $a_k$  are given by




$$a_k = \frac{1}{2\pi i} \oint_C \frac{f(s)}{(s - z_0)^{k+1}} ds, \quad k = 0, \pm 1, \pm 2, \dots$$

$C$  : a simple closed curve  
that lies entirely within  $D$   
that encloses  $z_0$


# Curve $C$ & Domain $D$ of the Laurent Series

$$f(z) = \sum_{n=-\infty}^{\infty} a_n^{(m)} (z - z_m)^n$$

$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z')}{(z' - z_m)^{n+1}} dz'$$
$$= \sum_k \operatorname{Res} \left( \frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$



# Expansion Points and Evaluation Points



•  $z_0$ : expansion point

$z$ : evaluation point

•

which poles of  $f(z)$  lie between the point of evaluation  $z$  and the point  $z_0$  about which the expansion is formed

$\frac{f(z')}{(z' - z_0)}$  is analytic between  $C_1$  &  $C_2$

deformation theorem

$C_1 - C_2$  coincide

common contour  $\curvearrowright$


# Residues

$$a_{-1} = \frac{1}{2\pi i} \oint_C f(s) ds \quad \rightarrow \quad \oint_C f(s) ds = 2\pi i \cdot a_{-1}$$

$$a_{-1} = \frac{1}{2\pi i} \oint_C f(s) ds = \text{Res}(f(z), z_0)$$

$$= \begin{cases} \lim_{z \rightarrow z_0} (z - z_0) f(z) & \text{(simple)} \\ \frac{1}{(n-1)!} \lim_{z \rightarrow z_0} \frac{d^{n-1}}{dz^{n-1}} (z - z_0)^n f(z) & \text{(order } n) \end{cases}$$

# Series Expansion at $z=0$




$$f(z) = \sum_{n=\eta_1}^{\infty} a_n^{(mf)} z^n$$

$$a_n^{(mf)} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{z^{n+1}} dz$$
$$= \sum_k \text{Res} \left( \frac{f(z)}{z^{n+1}}, z_k \right)$$


Poles  $z_k$

$$n \geq 0 \quad z_1, z_2, z_3, \circ$$


$$n < 0 \quad z_1, z_2, z_3$$




$\text{Res}(f, z_m)$




$\text{Res}(f, z_1)$   
 $\text{Res}(f, z_m)$




$\text{Res}(f, z_1) + \text{Res}(f, z_2) + \text{Res}(f, z_m)$




$\text{Res}(f, z_1) + \text{Res}(f, z_2) + \text{Res}(f, z_3)$   
 $+ \text{Res}(f, z_m)$




$$\text{Res}(f, z_m) \quad (n \geq 0)$$



$$\begin{aligned} &\text{Res}(f, z_1) \\ &\text{Res}(f, z_m) \quad (n \geq 0) \end{aligned}$$



$$\begin{aligned} &\text{Res}(f, z_1) + \text{Res}(f, z_2) + \text{Res}(f, z_m) \\ &\quad (n \geq 0) \end{aligned}$$



$$\begin{aligned} &\text{Res}(f, z_1) + \text{Res}(f, z_2) + \text{Res}(f, z_3) \\ &+ \text{Res}(f, z_m) \quad (n \geq 0) \end{aligned}$$



# Inverse z-Transform $x[n] = \frac{1}{2\pi i} \int_C X(z) z^{n-1} dz$

$$X(z) = \sum_{k=0}^{\infty} x_k z^{-k}$$

$$z^{n-1} X(z) = \left( \sum_{k=0}^{\infty} x_k z^{-k} \right) z^{n-1}$$

$$\int z^{n-1} \text{LHS} dz = \int \text{RHS} z^{n-1} dz$$

$$= \sum_{k=0}^{\infty} x_k z^{-k+n-1}$$

$$[0, \infty) = [0, n-1] \cup [n] \cup [n+1, \infty)$$

$$= \sum_{k=0}^{n-1} x_k z^{-k+n-1} + \sum_{k=n}^n x_k z^{-k+n-1} + \sum_{k=n+1}^{\infty} x_k z^{-k+n-1}$$

$$= \sum_{k=0}^{n-1} x_k z^{-k+n-1} + \frac{x_n}{z^1} + \sum_{k=n+1}^{\infty} \frac{x_k}{z^{k-n+1}}$$

$$\int_C X(z) z^{n-1} dz = \int_C \sum_{k=0}^{n-1} x_k z^{-k+n-1} dz + \int_C \frac{x_n}{z^1} dz + \int_C \sum_{k=n+1}^{\infty} \frac{x_k}{z^{k-n+1}} dz$$

$$= \sum_{k=0}^{n-1} x_k \int_C z^{-k+n-1} dz + x_n \int_C \frac{1}{z^1} dz + \sum_{k=n+1}^{\infty} x_k \int_C \frac{1}{z^{k-n+1}} dz$$

$$= \sum_{k=0}^{n-1} x_k \cdot 0 + x_n \cdot 2\pi i + \sum_{k=n+1}^{\infty} x_k \cdot 0$$

$$x[n] = \frac{1}{2\pi i} \int_C X(z) z^{n-1} dz$$

## Z-transform

$$z_m = 0$$

$$\begin{aligned} x[n] &= \frac{1}{2\pi i} \oint_C f(z) z^{n-1} dz \\ &= \sum_k \operatorname{Res}(f(z) z^{n-1}, z_k) \end{aligned}$$

$n > 0$   $z_k$ : poles of  $f(z)$

$n = 0$   $z_k$ : poles of  $f(z)$  +  $z = 0$   
 $z^{n-1} = z^{-1} = \frac{1}{z}$

$x[n]$  includes  $u[n] \rightarrow X[z]$  contains  $z$  on its numerator

Also, think about modified partial fraction  $\frac{X[z]}{z}$

## Laurent Expansion

expansion at  $z_m$

$$\begin{aligned} a_n^{\{m\}} &= \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z-z_m)^{n+1}} dz \\ &= \sum_k \operatorname{Res}\left(\frac{f(z)}{(z-z_m)^{n+1}}, z_k\right) \end{aligned}$$





$$z_m = 0$$

$$\begin{aligned} a_n^{\{0\}} &= \frac{1}{2\pi i} \oint_C \frac{f(z)}{z^{n+1}} dz \\ &= \sum_k \operatorname{Res}\left(\frac{f(z)}{z^{n+1}}, z_k\right) \end{aligned}$$

$$\begin{aligned} a_{-n}^{\{0\}} &= \frac{1}{2\pi i} \oint_C f(z) z^{n-1} dz \\ &= \sum_k \operatorname{Res}(f(z) z^{n-1}, z_k) \end{aligned}$$


$$\begin{aligned} a_{-n}^{\{0\}} &= \frac{1}{2\pi i} \oint_C \frac{f(z)}{z^{-n+1}} dz \\ &= \sum_k \operatorname{Res}\left(\frac{f(z)}{z^{-n+1}}, z_k\right) \end{aligned}$$

# Different D, Different Laurent Series



$$x[n] = \frac{1}{2\pi i} \int_C X(z) z^{n-1} dz$$

$$= \sum_{z_k} \text{Res}(X(z) z^{n-1}, z_k)$$



z-transform

$$f(z) = \frac{-1}{(z-1)(z-2)}$$

Complex Variables and Ap  
Brown & Churchill

$$f(z) = \frac{-1}{(z-1)(z-2)} = \frac{1}{z-1} - \frac{1}{z-2}$$

$$D_1: |z| < 1$$

$$D_2: 1 < |z| < 2$$

$$D_3: 2 < |z|$$

$$\textcircled{1} D_1 \quad |z| < 1, \quad \left|\frac{z}{2}\right| < 1$$

$$\begin{aligned} f(z) &= \frac{1}{z-1} - \frac{1}{z-2} = \frac{-1}{1-z} + \frac{1}{2} \frac{1}{1-\left(\frac{z}{2}\right)} \\ &= -\sum_{n=0}^{\infty} z^n + \sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} = \sum_{n=0}^{\infty} (2^{-n-1} - 1)z^n \quad |z| < 1 \end{aligned}$$

$$\textcircled{2} D_2 \quad 1 < |z| < 2 \Rightarrow \left|\frac{1}{z}\right| < 1, \quad \left|\frac{z}{2}\right| < 1$$


$$\begin{aligned} f(z) &= \frac{1}{z-1} - \frac{1}{z-2} = \frac{1}{z} \cdot \frac{1}{1-\left(\frac{1}{z}\right)} + \frac{1}{2} \frac{1}{1-\left(\frac{z}{2}\right)} \\ &= \sum_{n=0}^{\infty} \frac{1}{z^{n+1}} + \sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} \\ &= \sum_{n=1}^{\infty} \frac{1}{z^n} + \sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} \end{aligned}$$

$$\textcircled{3} D_3 \quad 2 < |z| \quad \left|\frac{z}{2}\right| < 1 \quad \left|\frac{1}{z}\right| < 1$$

$$\begin{aligned} f(z) &= \frac{1}{z-1} - \frac{1}{z-2} = \frac{1}{z} \frac{1}{1-\left(\frac{1}{z}\right)} - \frac{1}{z} \frac{1}{1-\left(\frac{z}{2}\right)} \\ &= \sum_{n=0}^{\infty} \frac{1}{z^{n+1}} - \sum_{n=0}^{\infty} \frac{z^n}{z^{n+1}} = \sum_{n=0}^{\infty} \frac{1-2^n}{z^{n+1}} \\ &= \sum_{n=1}^{\infty} \frac{1-2^{n-1}}{z^n} \end{aligned}$$

$$f(z) = \frac{-1}{(z-1)(z-2)}$$

①  $D_1 \quad |z| < 1, \quad \left|\frac{z}{2}\right| < 1$




$$\frac{f(z)}{z^{n+1}} = \frac{-1}{(z-1)(z-2)z^{n+1}}$$

$$\begin{aligned} f(z) &= \frac{1}{z-1} - \frac{1}{z-2} = \frac{-1}{1-z} + \frac{1}{1-\left(\frac{z}{2}\right)} \\ &= -\sum_{n=0}^{\infty} z^n + \sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} = \sum_{n=0}^{\infty} (2^{-n-1} - 1)z^n \quad |z| < 1 \end{aligned}$$

$$a_n = \frac{f(z)}{z^{n+1}} = \frac{1}{(z-1)(z-2)z^{n+1}} \quad \frac{1}{z-1} - \frac{1}{z-2}$$

$$a_n = \sum_{k=1}^M \operatorname{Res} \left( \frac{f(z)}{(z-z_k)^{n+1}}, z_k \right) = \operatorname{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 0 \right)$$



$$a_n = \sum_{k=1}^M \operatorname{Res} \left( \frac{f(z)}{(z-z_k)^{n+1}}, z_k \right) = \operatorname{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 0 \right)$$

$n \geq 0$  then the pole  $z=0$

$$\frac{1}{(n-1)!} \lim_{z \rightarrow z_0} \frac{d^{n-1}}{dz^{n-1}} (z-z_0)^n f(z) \quad (\text{order } n)$$

$$\frac{d}{dz} ((z-1)^{-1} - (z-2)^{-1}) = (-1) ((z-1)^{-2} - (z-2)^{-2})$$

$$\frac{d^2}{dz^2} ((z-1)^{-1} - (z-2)^{-1}) = (-1)(-2) ((z-1)^{-3} - (z-2)^{-3})$$

$$\frac{d^3}{dz^3} ((z-1)^{-1} - (z-2)^{-1}) = (-1)(-2)(-3) ((z-1)^{-4} - (z-2)^{-4})$$

$$\frac{d^n}{dz^n} ((z-1)^{-1} - (z-2)^{-1}) = (-1)^n n! ((z-1)^{-n-1} - (z-2)^{-n-1})$$


$$\begin{aligned} \frac{1}{n!} \lim_{z \rightarrow 0} \frac{d^n}{dz^n} ((z-1)^{-1} - (z-2)^{-1}) &= (-1)^n \lim_{z \rightarrow 0} ((z-1)^{-n-1} - (z-2)^{-n-1}) \\ &= (-1)^n ((-1)^{-n-1} - (-2)^{-n-1}) \\ &= -1 + 2^{-n-1} \end{aligned}$$

$$a_n = -1 + 2^{-n-1} \quad (n \geq 0)$$

$$f(z) = \sum_{n=-n_1}^{\infty} a_n z^n = \sum_{n=0}^{\infty} (2^{-n-1} - 1) z^n$$


$$f(z) = \frac{-1}{(z-1)(z-2)}$$

$$\textcircled{2} \quad D_2 \quad 1 < |z| < 2 \Rightarrow \left| \frac{1}{z} \right| < 1, \quad \left| \frac{z}{2} \right| < 1$$



$$\begin{aligned} f(z) &= \frac{1}{z-1} - \frac{1}{z-2} = \frac{1}{z} \cdot \frac{1}{1-\left(\frac{1}{z}\right)} + \frac{1}{2} \frac{1}{1-\left(\frac{z}{2}\right)} \\ &= \sum_{n=0}^{\infty} \frac{1}{z^{n+1}} + \sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} \\ &= \sum_{n=1}^{\infty} \frac{1}{z^n} + \sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} \end{aligned}$$

$$\begin{aligned} a_n &= \sum_{k=1}^M \operatorname{Res} \left( \frac{f(z)}{(z-z_k)^{n+1}}, z_k \right) = \operatorname{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 0 \right) \\ &\quad + \operatorname{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 1 \right) \end{aligned}$$



$$a_n = \sum_{k=1}^M \operatorname{Res} \left( \frac{f(z)}{(z-z_k)^{n+1}}, z_k \right) = \operatorname{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 0 \right) + \operatorname{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 1 \right)$$

$$\frac{1}{(n-1)!} \lim_{z \rightarrow z_0} \frac{d^{n-1}}{dz^{n-1}} (z-z_0)^n f(z) \quad (\text{order } n)$$

$$\begin{aligned} \frac{1}{n!} \lim_{z \rightarrow 0} \frac{d^n}{dz^n} ((z-1)^{-1} - (z-2)^{-1}) &= (-1)^n \lim_{z \rightarrow 0} ((z-1)^{-n-1} - (z-2)^{-n-1}) \\ &= (-1)^n ((-1)^{-n-1} - (-2)^{-n-1}) \\ &= -1 + 2^{-n-1} \end{aligned}$$

$$\operatorname{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 0 \right) = -1 + 2^{-n-1} \quad (n \geq 0)$$

$$\operatorname{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 1 \right) = \lim_{z \rightarrow 1} (z-1) \frac{-1}{(z-1)(z-2)z^{n+1}} = 1$$

$n=-3$	$n=-2$	$n=-1$	$n=0$	$n=1$	$n=2$	
0	0	0	$-1+2^{-1}$	$-1+2^{-2}$	$-1+2^{-3}$	$\operatorname{Res} \left( \frac{f(z)}{z^{n+1}}, 0 \right)$
1	1	1	1	1	1	$\operatorname{Res} \left( \frac{f(z)}{z^{n+1}}, 1 \right)$
1	1	1	$2^{-1}$	$2^{-2}$	$2^{-3}$	


$$\begin{cases} a_n = 2^{-n-1} & n \geq 0 \\ a_n = 1 & n < 0 \end{cases} \quad \begin{cases} 2^{-n-1} z^n \\ z^{-n} \end{cases}$$

$$f(z) = \sum_{n=1}^{\infty} \frac{1}{z^n} + \sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}}$$




$$f(z) = \frac{-1}{(z-1)(z-2)}$$

③  $D_3 \quad 2 < |z| \quad \left| \frac{2}{z} \right| < 1 \quad \left| \frac{1}{z} \right| < 1$



$$\begin{aligned} f(z) &= \frac{1}{z-1} - \frac{1}{z-2} = \frac{1}{z} \frac{1}{1-(\frac{1}{z})} - \frac{1}{z} \frac{1}{1-(\frac{2}{z})} \\ &= \sum_{n=0}^{\infty} \frac{1}{z^{n+1}} - \sum_{n=0}^{\infty} \frac{2^n}{z^{n+1}} = \sum_{n=0}^{\infty} \frac{1-2^n}{z^{n+1}} \\ &= \sum_{n=1}^{\infty} \frac{1-2^{n-1}}{z^n} \end{aligned}$$

$$\begin{aligned} a_n &= \sum_{k=1}^M \text{Res} \left( \frac{f(z)}{(z-z_k)^{n+1}}, z_k \right) = \text{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 0 \right) \\ &\quad + \text{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 1 \right) \\ &\quad + \text{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 2 \right) \end{aligned}$$



$$\text{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 0 \right) = -1 + 2^{-n+1} \quad (n \geq 0)$$

$$\text{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 1 \right) = \lim_{z \rightarrow 1} (z-1) \frac{-1}{(z-1)(z-2)z^{n+1}} = 1$$


$$\text{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 2 \right) = \lim_{z \rightarrow 2} (z-2) \frac{-1}{(z-1)(z-2)z^{n+1}} = -\frac{1}{2^{n+1}}$$

$n=-3$	$n=-2$	$n=-1$	$n=0$	$n=1$	$n=2$	
0	0	0	$-1+2^1$	$-1+2^2$	$-1+2^3$	$\text{Res} \left( \frac{f(z)}{z^{n+1}}, 0 \right)$
1	1	1	1	1	1	$\text{Res} \left( \frac{f(z)}{z^{n+1}}, 1 \right)$
$-2^2$	$-2$	$-1$	$-2^1$	$-2^2$	$-2^3$	$\text{Res} \left( \frac{f(z)}{z^{n+1}}, 2 \right)$
$1-2^2$	$1-2$	0	0	0	0	

$$a_n = 1 - 2^{-n+1} \quad n < 0 = \sum_{n=1}^{\infty} \frac{1-2^{n-1}}{z^n}$$

$$f(z) = \sum_{n=-1}^{-\infty} (1-2^{-n+1}) z^n = \sum_{n=1}^{\infty} \frac{1-2^{n-1}}{z^n}$$

$$f(z) = \frac{-1}{(z-1)(z-2)}$$



$$x[n]$$

$$= \frac{1}{2\pi i} \int_C \boxed{X(z) z^{n-1}} dz$$

$$= \sum_{j=1}^k \text{Res}(\boxed{X(z) z^{n-1}}, z_j)$$


$$X(z) = \frac{-1}{(z-1)(z-2)}$$

$$X(z) z^{n-1} = \frac{-1}{(z-1)(z-2)} z^{n-1}$$

$$\text{Res}(\boxed{X(z) z^{n-1}}, 1) = (z-2) \frac{-1}{(z-1)(z-2)} z^{n-1} \Big|_{z=1} = 1$$

$$\text{Res}(\boxed{X(z) z^{n-1}}, 2) = (z-1) \frac{-1}{(z-1)(z-2)} z^{n-1} \Big|_{z=2} = -2^{n-1}$$

$$x[n] = 1 - 2^{n-1}$$



ROC (Region of Convergence)

$$|z| > 2 \Rightarrow \frac{2}{|z|} < 1$$

$$\left(\frac{2}{z}\right)^0 + \left(\frac{2}{z}\right)^1 + \left(\frac{2}{z}\right)^2 + \dots \longrightarrow \frac{1}{1 - \frac{2}{z}}$$

Converge

$$|z| > 2 \Rightarrow \frac{1}{|z|} < 1$$

$$\left(\frac{1}{z}\right)^0 + \left(\frac{1}{z}\right)^1 + \left(\frac{1}{z}\right)^2 + \dots \longrightarrow \frac{1}{1 - \frac{1}{z}}$$

Converge

$$\begin{aligned} f(z) &= \frac{1}{z-1} - \frac{1}{z-2} = \frac{1}{z} \frac{1}{1 - (\frac{1}{z})} - \frac{1}{z} \frac{1}{1 - (\frac{2}{z})} \\ &= \sum_{n=0}^{\infty} \frac{1}{z^{n+1}} - \sum_{n=0}^{\infty} \frac{2^n}{z^{n+1}} = \sum_{n=0}^{\infty} \frac{1 - 2^n}{z^{n+1}} \\ &= \sum_{n=1}^{\infty} \frac{1 - 2^{n-1}}{z^n} \end{aligned}$$

$$\left(\frac{1}{z}\right) + \left(\frac{1}{z}\right)^2 + \left(\frac{1}{z}\right)^3 + \dots + \frac{1}{2} \left\{ \left(\frac{2}{z}\right) + \left(\frac{2}{z}\right)^2 + \left(\frac{2}{z}\right)^3 + \dots \right\} \longrightarrow \frac{1}{z-1} - \frac{1}{z-2} = \frac{-1}{(z-1)(z-2)}$$

Converge

$$(1-2^0)z^{-1} + (1-2^1)z^{-2} + (1-2^2)z^{-3} + \dots \longrightarrow \frac{-1}{(z-1)(z-2)} \quad (|z| > 2)$$

Converge

$$x[n] = 1 - 2^n \quad \longleftrightarrow \quad X(z) = \frac{-1}{(z-1)(z-2)} \quad (|z| > 2)$$





