Laurent Series and z-Transform

Geometric Series Double Pole Examples B

7	\cap	1 (\cap	7	1	\cap
2	U	Ιì	31	U	Z	T	9

Copyright (c) 2016 - 2018 Young W. Lim.

Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.2 or any later version published by the Free Software Foundation; with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts. A copy of the license is included in the section entitled "GNU Free Documentation License".

2 formulas of z

$$\frac{-1}{(2-1)(2-2)} = \left(\frac{1}{\xi-1} - \frac{1}{\xi-2}\right)$$

$$\frac{-052^2}{(2-1)(2-0.5)} = \left(-\frac{2}{(2-1)} + \frac{0.52}{(2-0.5)}\right)$$

$$\frac{-1}{(2^{-1})(2^{-2})} = \left(\frac{1}{\xi - 1} - \frac{1}{\xi - 2}\right)$$

$$= \left(\frac{1}{\xi - 0.5} - \frac{1}{\xi - 2}\right)$$

$$= \left(\frac{1}{\xi^{-1} - 1} - \frac{1}{\xi^{-2}}\right)$$

$$= \left(\frac{\xi}{1 - \xi} - \frac{\xi}{1 - 2\xi}\right)$$

$$= \left(\frac{-\xi}{2 - 1} + \frac{0.5\xi}{2 - 0.5}\right)$$

$$= \xi \left(\frac{-1}{\xi - 1} + \frac{0.5\xi}{2 - 0.5}\right)$$

$$= \xi \left(\frac{-0.5\xi^{2}}{(\xi - 1)(\xi - 0.5)}\right)$$

$$\frac{-0.52^{2}}{(2-1)(2-0.5)} = \left(-\frac{2}{(2-1)} + \frac{0.52}{(2-0.5)}\right)$$

$$\frac{(3-1)(3-2)}{-1} = \left(\frac{\xi-1}{\xi-1} - \frac{\xi-2}{\xi}\right)$$

$$\frac{f(z)}{|z| > 2} \qquad f(z) = \frac{z^{-1}}{|-z^{-1}|} - \frac{z^{-1}}{|-z|^{-1}} \qquad + |^{n+1} - (\frac{1}{z})^{n+1}| \qquad (n < 0)$$

$$\frac{\chi(\xi)}{|\xi| > 2} \qquad \chi(\xi) = \frac{\xi^{-1}}{|-\xi^{-1}|} - \frac{\xi^{-1}}{|-2\xi^{-1}|} \qquad + \frac{|-1|}{|-2|^{n-1}} - \frac{|-1|}{|-2|}$$

$$\frac{-05\xi^2}{(2-1)(2-0.5)} = \left(-\frac{\xi}{(\xi-1)} + \frac{0.5\xi}{(\xi-0.5)}\right)$$

$$\frac{f(\xi)}{|\xi| > 2} \qquad f(\xi) = -\frac{1}{1-\xi^{-1}} + \frac{0.5}{1-0.5\xi^{-1}} \qquad -|\frac{n-1}{1} + 2^{n-1} \qquad (\cap < |)$$

$$2-B |\xi| < | \qquad \qquad |\xi| < | \qquad \qquad |\xi| < | \qquad \qquad |\xi| = | \qquad \qquad |\xi| - |\xi| - |\xi| + |\xi| - |\xi| - |\xi| + |\xi| - |\xi| -$$

$$\frac{\chi(\xi)}{|\xi| > 2} \qquad \chi(\xi) = -\frac{1}{|-\xi^{-1}|} + \frac{0.5}{|-0.5\xi^{-1}|} - \frac{1}{n+1} + \left(\frac{1}{2}\right)^{n+1} \qquad (\eta \geqslant 0)$$

		(1) (2-2)	$2 \frac{-0.52^{2}}{(2-1)(2-0.5)}$
(A)	2 < 1	$- ^{n+1}+(\frac{1}{2})^{n+1}$ (n>0)	$+ \mid \stackrel{n-l}{-} - 2^{n-l} (n \geqslant l)$
f(2)	2 > 2	$+ \frac{1}{n+1} - (\frac{1}{2})^{n+1} $ (n<0)	$-1^{n-1}+2^{n-1}(n<1)$
B	2 <	- n-1 + 2 n-1 (n<)	$+ \frac{z}{u+1} - (\frac{z}{1})_{U+1} (U < 0)$
Χ(₹)	2 > 2	+ n-1 - 2 n-1 (n>1)	$\frac{- \left \frac{2}{0+1} + \left(\frac{2}{1} \right)_{U+1} (U \geqslant 0) \right }{\left \frac{2}{0+1} + \left(\frac{2}{1} \right)_{U+1} \right }$

		(2-1)(2-2)	
	f(2)	$\frac{-1}{n+1} + \left(\frac{7}{1}\right)_{U+1} (U > 0)$	$+ \frac{n-1}{2} - 2^{n-1} (n \ge 1)$
	X(Z)	- n-1 + 2 n-1 (n<)	+ 1 - (1) (1 < 0)
	f(2)	$+ \frac{1}{n+1} - (\frac{2}{1})^{n+1} $ (n<0)	$-1^{n-1}+2^{n-1}(n<1)$
2 > 2	X(2)	+ n-1 - 2 n-1 (n>1)	$\frac{- \left(\frac{2}{1}\right)^{n+1} + \left(\frac{2}{1}\right)^{n+1} (u > 0)}{1}$

		(2-1)(2-2)	$2 \frac{-0.52^2}{(2-1)(2-0.5)}$
	f(2)	causal (n>0)	causal (N>1)
2 > 2	f(2)	anticausal (n<0)	anticausal (n<1)
2 <	X(₹)	anticausal (n<1)	anticausal (n<0)
 	X(2)	causal (n>1)	causal (n>0)

		(2-1)(2-2)	$2\frac{-052^{2}}{(2-1)(2-0.5)}$
2 <	f({ })	causal (n>0)	causal (n>1)
2 <	Χ(₹)	anticausal (N<1)	anticausal (n<0)
& > 2	f(2)	anticausal (n<0)	anticausal (N<1)
z - 2	X(3)	causal (n31)	causal (n > 0)

$$\frac{-1}{(2-1)(2-2)} \longrightarrow 2 \frac{-052^2}{(2-1)(2-0.5)}$$

$$\left(\frac{1}{\xi-0.5}-\frac{1}{\xi-2}\right) \qquad \left(\frac{0.5\xi}{(\xi-0.5)}-\frac{2\xi}{(\xi-1)}\right)$$

$$\frac{2}{1-2\xi} + \frac{0.5}{1-0.5\xi}$$
 $\frac{2}{1-2\xi} + \frac{\xi}{1-0.5\xi}$

$$\frac{z^{-1}}{1-0.5z^{-1}} - \frac{z^{-1}}{1-2z^{-1}}$$

$$\frac{z^{-1}}{1-0.5z^{-1}} - \frac{2}{1-2z^{-1}}$$

$$\frac{-1}{(2-1)(2-2)} \qquad 2 \qquad \frac{-05\xi^2}{(2-1)(2-0.5)}$$

$$\left(\frac{1}{\xi-0.5}-\frac{1}{\xi-2}\right)$$

$$\left(\frac{0.5\xi}{(\xi-0.5)}-\frac{2\xi}{(\xi-\Sigma)}\right)$$

$$\frac{2}{1-2z} + \frac{0.5}{1-0.5z}$$

$$\frac{2}{1-2z} + \frac{0.5}{1-0.5z}$$
 $\frac{2}{1-2z} + \frac{z}{1-0.5z}$

$$\frac{z^{-1}}{|-0.5z^{-1}|} - \frac{z^{-1}}{|-2z^{-1}|} = \frac{0.5}{|-0.5z^{-1}|} - \frac{2}{|-2z^{-1}|}$$

$$\frac{0.5}{1 - 0.5 z^{-1}} - \frac{2}{1 - 2 z^{-1}}$$

$$(n \ge 1)$$

$$X(2)$$
 causal $(n \ge 0)$

$$(n \ge 0)$$

$$\frac{-1}{(2-1)(2-2)} = \frac{z^{-1}}{(2-1)(2-0.5)}$$

$$\left(\frac{1}{\xi-0.5}-\frac{1}{\xi-2}\right) \qquad \left(\frac{0.5\xi}{(\xi-0.5)}-\frac{2\xi}{(\xi-1)}\right)$$


$$\frac{2}{1-2z} + \frac{0.5}{1-0.5z}$$
 $\frac{2}{1-2z} + \frac{z}{1-0.5z}$

$$|2|<0.5$$
 f(2) causal $(n>0)$ $|2|<0.5$ f(2) causal $(n>1)$

$$X(2)$$
 anticausal $(n < 1)$ $X(2)$ anticausal $(n < 0)$

$$\frac{\xi^{-1}}{|-0.5\xi^{-1}|} - \frac{\xi^{-1}}{|-2\xi^{-1}|} = \frac{0.5}{|-0.5\xi^{-1}|} - \frac{2}{|-2\xi^{-1}|}$$

$$|\vec{z}| \neq 1$$
 f(z) anticawal (n < 0)
$$|\vec{z}| \neq 2$$
 f(z) anticawal (n < 1)
$$|\vec{z}| \neq 2$$
 X(z) causal (n > 0)



$$-\frac{2}{1-2\xi}+\frac{0.5}{1-0.5\xi}$$

$$-\frac{\xi}{|-2\xi|}+\frac{\xi}{|-0.5\xi|}$$

$$\frac{f(z) = -\left[2 + 2^{2}z^{1} + 2^{3}z^{2} + \cdots\right] -2^{\frac{n+1}{2}}}{+\left[\left(\frac{1}{2}\right) + \left(\frac{1}{2}\right)^{2}z^{2} + \left(\frac{1}{2}\right)^{3}z^{2} + \cdots\right] + \left(\frac{1}{2}\right)^{n+1}}$$

$$\frac{f(z) = -\left[2^{0}z^{1} + 2^{1}z^{2} + 2^{2}z^{3} + \cdots\right] -2^{n+1}}{+\left[\left(\frac{1}{2}\right)^{0}z^{1} + \left(\frac{1}{2}\right)^{1}z^{2} + \left(\frac{1}{2}\right)^{2}z^{3} + \cdots\right] + \left(\frac{1}{2}\right)^{n+1}}$$

18172

$$\frac{\xi^{-1}}{1 - 0.5\xi^{-1}} = \frac{\xi^{-1}}{1 - 2\xi^{-1}}$$

$$\frac{0.5}{|-0.5 \, \epsilon^{-1}|} \frac{2}{|-2 \, \hat{\epsilon}^{-1}|}$$

$$\frac{(2)}{\sqrt{(2)}} = + \left[\left(\frac{1}{2} \right)^{6} z^{1} + \left(\frac{1}{2} \right)^{6} z^{2} + \left(\frac{1}{2} \right)^{3} z^{-3} + \cdots \right] + \left(\frac{1}{2} \right)^{n-1} - \left[2^{6} z^{-1} + 2^{1} z^{-2} + 2^{2} z^{-3} + \cdots \right] - 2^{n-1}$$

$$\frac{2}{\sqrt{(2)}} = \frac{1}{\sqrt{(2)}} \left[\frac{1}{\sqrt{(2)}} z^{1} + \left(\frac{1}{2} \right)^{3} z^{-3} + \cdots \right] - 2^{n-1}$$

$$\frac{(2)}{-\left(\frac{1}{2}\right)^{\frac{1}{2}}} = + \left[\left(\frac{1}{2}\right)^{\frac{1}{2}} z^{-\frac{1}{2}} + \left(\frac{1}{2}\right)^{\frac{3}{2}} z^{-\frac{1}{2}} + \cdots\right] + \left(\frac{1}{2}\right)^{\frac{3}{4}+1}}{-\left(\frac{1}{2}\right)^{\frac{3}{2}} z^{-\frac{1}{2}} + \cdots\right] - 2^{\frac{3}{4}+1}}$$

$$\frac{1}{2}$$

$$\frac{1}{2}$$

$$\frac{-1}{(2-1)(2-2)} \longrightarrow 2 \frac{-05\xi^2}{(2-1)(2-0.5)}$$

$$\frac{2}{1-2\xi}+\frac{0.5}{1-0.5\xi}$$

$$f(\xi) = -\left[2 + 2^{2}\xi^{1} + 2^{3}\xi^{2} + \cdots\right] -2^{n+1} + \left[\left(\frac{1}{2}\right) + \left(\frac{1}{2}\right)^{2}\xi^{2} + \left(\frac{1}{2}\right)^{3}\xi^{2} + \cdots\right] + \left(\frac{1}{2}\right)^{n+1}$$

$$X (Z) = -\left[\left(\frac{1}{2} \right)^{-1} + \left(\frac{1}{2} \right)^{-2} z^{1} + \left(\frac{1}{2} \right)^{-3} z^{2} + \cdots \right] - \left(\frac{1}{2} \right)^{n-1} + \left[2^{-1} + 2^{-2} z^{1} + 2^{-3} z^{2} + \cdots \right] + 2^{n-1}$$

18172

$2 = \left(\frac{1}{2}\right)^{-1} \quad f(x) = x^{-1}$

$$f(z) = + \left[2^{\circ} z^{1} + 2^{-1} z^{-2} + 2^{-2} z^{-3} + \cdots \right] + 2^{n+1}$$

$$- \left[\left(\frac{1}{2} \right)^{\circ} z^{4} + \left(\frac{1}{2} \right)^{-1} z^{-2} + \left(\frac{1}{2} \right)^{-2} z^{-3} + \cdots \right] - \left(\frac{1}{2} \right)^{n+1}$$

121<0.5

$$-\frac{\xi}{|-2\xi|} + \frac{\xi}{|-0.5\xi|}$$

$$f(z) = -\left[2^{0}z^{1} + 2^{1}z^{2} + 2^{2}z^{3} + \cdots\right] -2^{n-1} + \left[\left(\frac{1}{2}\right)^{0}z^{1} + \left(\frac{1}{2}\right)^{1}z^{2} + \left(\frac{1}{2}\right)^{2}z^{3} + \cdots\right] + \left(\frac{1}{2}\right)^{n-1}$$

$$\frac{0.5}{1-0.5\,\epsilon^{-1}} - \frac{2}{1-2\,\epsilon^{-1}}$$

$$f(z) = + \left[2^{4} z^{6} + 2^{-2} z^{-1} + 2^{-3} z^{-2} + \cdots \right] + 2^{n-1}$$

$$- \left[\left(\frac{1}{2} \right)^{\frac{1}{2}} {}^{6} + \left(\frac{1}{2} \right)^{-\frac{3}{2}} z^{-1} + \left(\frac{1}{2} \right)^{\frac{3}{2}} z^{-2} + \cdots \right] - \left(\frac{1}{2} \right)^{\frac{3}{2}-1}$$

$$\begin{array}{c} \chi \left(\xi \right) = + \left[\left(\frac{1}{2} \right)^{1} \xi^{0} + \left(\frac{1}{2} \right)^{3} \xi^{-1} + \left(\frac{1}{2} \right)^{3} \xi^{-2} + \cdots \right] & \uparrow \left(\frac{1}{2} \right)^{n+1} \\ - \left[2^{1} \xi^{0} + 2^{3} \xi^{-1} + 2^{3} \xi^{-2} + \cdots \right] & -2^{n+1} \end{array}$$

$$\frac{z^{-1}}{(z-1)(z-0.5)}$$

121<0.5

$$-\frac{2}{1-2\xi}+\frac{0.5}{1-0.5\xi}$$

$$f(\xi) = -\left[2 + 2^{3}\xi + 2^{3}\xi^{3} + \cdots\right]$$
$$+\left[\left(\frac{1}{2}\right) + \left(\frac{1}{2}\right)^{3}\xi + \left(\frac{1}{2}\right)^{3}\xi^{3} + \cdots\right]$$

$$(n) = -2^{n+1} + (\frac{1}{2})^{n+1} \quad (n > 0)$$

$$\frac{(2) = -\left[\left(\frac{1}{2}\right)^{-1} + \left(\frac{1}{2}\right)^{-2} z^{1} + \left(\frac{1}{2}\right)^{-3} z^{2} + \cdots\right]}{+\left[2^{-1} + 2^{-2} z^{1} + 2^{-3} z^{2} + \cdots\right]}$$

$$(n = -(\frac{1}{2})^{n-1} + 2^{n-1} \quad (n < [)$$

151 < 0.5

$$\frac{z}{|-2z|} + \frac{z}{|-0.5z|}$$

$$f(z) = -\left[2^{0}z^{1} + 2^{1}z^{2} + 2^{2}z^{3} + \cdots\right] + \left[\left(\frac{1}{2}\right)^{0}z^{1} + \left(\frac{1}{2}\right)^{1}z^{2} + \left(\frac{1}{2}\right)^{2}z^{3} + \cdots\right]$$

$$\Delta_n = -2^{n-1} + \left(\frac{1}{2}\right)^{n-1} \quad (n > 1)$$

$$Q_n = -\left(\frac{1}{2}\right)^{n+1} + 2^{n+1} \quad (\cap < 0)$$

18172

$$f(Z) = + \left[2^{\circ} \xi^{-1} + 2^{-1} \xi^{-2} + 2^{-2} \xi^{-3} + \cdots \right] - \left[\left(\frac{1}{2} \right)^{\circ} \xi^{-4} + \left(\frac{1}{2} \right)^{-1} \xi^{-2} + \left(\frac{1}{2} \right)^{-2} \xi^{-3} + \cdots \right]$$

$$a_n = +2^{n+1} - \left(\frac{1}{2}\right)^{n+1} \quad (n < 0)$$

18/72

$$\frac{0.5}{|-0.5\xi^{-1}|} - \frac{2}{|-2\xi^{-1}|}$$

$$f(z) = + \left[2^4 z^6 + 2^{-2} z^{-1} + 2^{-3} z^{-2} + \cdots \right] - \left[\left(\frac{1}{2} \right)^{-1} z^6 + \left(\frac{1}{2} \right)^{-2} z^{-1} + \left(\frac{1}{2} \right)^{-3} z^{-2} + \cdots \right]$$

$$\alpha_n = \pm 2^{n-1} - \left(\frac{1}{2}\right)^{n-1} \quad (n < 1)$$

$$\alpha_n = + \left(\frac{1}{2}\right)^{n-1} - 2^{n-1} \qquad (n > 1)$$

$$\Delta_n = t(\frac{1}{2})^{n+1} - 2^{n+1} \quad (n \ge 0)$$

		1 (2-0.5) (3-2)	$2^{\frac{3}{2}} \frac{-\xi^2}{(2-2)(2-0.5)}$
라 < 글	f(2)	$-2^{n+1}+(\frac{1}{2})^{n+1}$ (n>0)	$-2^{n-1}+(\frac{1}{2})^{n-1}(n \ge 1)$
161 7 2			
121 2 2	f(2)	$+ 5_{0+1} - (\frac{7}{1})_{0+1} $ ($1 < 0$)	$+2^{n_1}-(\frac{1}{2})^{n_1}(n<1)$
2 > 2			

		1 (2-0.2) (3-2)	2 3 22 (2-0.5)
z < \frac{1}{2}			
101 > 2	X(3)	$-(\frac{1}{2})^{n-1}+2^{n-1}$ (n<1)	$-\left(\frac{1}{2}\right)^{n+1} + 2^{n+1} (n < 0)$
151 5 3			-1
2 > 2	X(2)	$+(\frac{2}{1})^{n-1}-2^{n-1}$ (n>1)	$+\left(\frac{1}{2}\right)^{n+1}-2^{n+1} (n \ge 0)$

$$2^{-n+1} = \left(\frac{1}{2}\right)^{n} \cdot 2 = \left(\frac{1}{2}\right)^{n-1} \qquad \left(\frac{1}{2}\right)^{-n-1} = 2^{n} \cdot 2 = 2^{n+1}$$

$$\left(\frac{1}{2}\right)^{-n+1} = 2^{n} \cdot \frac{1}{2} = 2^{n-1} \qquad 2^{-n-1} = \left(\frac{1}{2}\right)^{n} \cdot \frac{1}{2} = \left(\frac{1}{2}\right)^{n+1}$$

			1 (2-0.5) (2-2)	$2^{\frac{3}{2}} \frac{-\xi^2}{(2-2)(2-0.5)}$
	2 < 1/2	f(2)	$-2^{n+1}+(\frac{1}{2})^{n+1}$ (n>0)	$-2^{n-1}+(\frac{1}{2})^{n-1}$ $(n\geqslant 1)$
	161 > 2		-n	1
٤'		f(2)	$+2^{n+1}-(\frac{1}{2})^{n+1}$ (n<0)	$+2^{n1}-(\frac{1}{2})^{n-1}(n<1)$
	 			

27

			1 (2-0.5) (2-2)	$2^{\frac{1}{2}}\frac{(2-2)(2-0.5)}{(2-2)(2-0.5)}$
	151 4 1			
77	라 < 코	X(2)	$-\left(\frac{1}{2}\right)^{n-1}+2^{n-1}$ (n<1)	$-\left(\frac{1}{2}\right)^{\eta+1}+2^{\eta+1}$ ($\eta<0$)
ξ'			-n	-1)
	2 > 2	X(2)	$+\left(\frac{2}{1}\right)^{n-1} - 2^{n-1} (n \ge 1)$	$+\left(\frac{1}{2}\right)^{n+1}-2^{n+1} (n \ge 0)$

	_		
		1 (2-0.5) (2-2)	$2^{\frac{3}{2}} \frac{-\xi^2}{(2-2)(2-0.5)}$
라 < 1	f(2)	$-2^{n+1}+(\frac{1}{2})^{n+1}$ (n>0)	$-2^{n-1}+(\frac{1}{2})^{n-1}$ $(n>1)$
161 \ 2		- N	-1
121 2 2	f(2)	$+ 5_{0+1} - (\frac{7}{1})_{0+1} $ ($1 < 0$)	$+2^{n1}-(\frac{1}{2})^{n-1}(n<1)$
2 > 2		-n,	1

₹-<u>]</u>

	_		
		1 2 (2-0.2) (3-2)	2 = -2 ² (2-2)(2-0.5)
라 < 1		ر٣-	
161 , 7	Χ(₹)	$-\left(\frac{1}{2}\right)^{n-1}+2^{n-1} (n<1)$	$-\left(\frac{1}{2}\right)^{\eta+1}+2^{\eta+1}$ (n<0)
		ر٣-	-1
121 2 2			
2 > 2	X(2)	$+\left(\frac{2}{1}\right)^{n-1}2^{n-1} (n>1)$	$+\frac{1}{2}^{n+1} - 2^{n+1} (n \ge 0)$

		1 (2-0.5) (2-2)	2 3 - 2² (2-2)(2-0.5)
라 < 1	f(₹)	$-2^{n+1}+(\frac{1}{2})^{n+1}$ (n>0)	
161 7 2	X(2)	$-\left(\frac{1}{2}\right)^{n-1}+2^{n-1}$ (n<1)	
151 5 3	f(})	$+2^{\frac{2}{n+1}}-(\frac{2}{1})^{n+1}$ (n<0)	-
2 > 2	X(2)	$+\left(\frac{7}{7}\right)_{U-I}-5_{U-I} (U \gg I)$	

		(3 (2-0.2) (5-2)	2 = -22 (2-0.5)
2 < 1	f(2)		$-2^{n-1}+(\frac{1}{2})^{n-1}$ $(n \ge 1)$
	X(£)		$-\left(\frac{1}{2}\right)^{n+1} + 2^{n+1} (n < 0)$
& > 2	f({ })		$+2^{n_1}-(\frac{1}{2})^{n_1}(n<1)$
	X(£)		$+\left(\frac{1}{2}\right)^{n+1}-2^{n+1} (n \ge 0)$

$$f(z)$$
 $|z| < 0.5$ $|z| > 2$

Causal anticausal

$$|\xi| < 0.5$$
 $f(\xi) = -\frac{2}{1-2\xi} + \frac{0.5}{1-0.5\xi} -2^{n+1} + (\frac{1}{2})^{n+1}$ $(N > 0)$

$$\frac{1-\alpha\xi}{\alpha} \frac{\xi^{-1}}{\alpha^{4}\xi^{4}-1} \frac{-\left(2+2^{3}\xi+2^{3}\xi^{2}+\cdots\right)+\left(\frac{1}{2}+\frac{1}{2}\xi+\frac{1}{2}\xi+\frac{1}{2}\xi+\frac{1}{2}\xi^{2}\xi^{2}+\cdots\right)}{n=0 \quad n=1 \quad n=2}$$

$$|\xi| > 2 \qquad f(\xi) = \frac{\xi^{-1}}{|-0.5\xi^{-1}|} - \frac{\xi^{-1}}{|-2\xi^{-1}|} + 2^{n+1} - (\frac{1}{2})^{n+1} \qquad (n < 0)$$

$$(\xi^{-1} + (\frac{1}{2})^{1} \xi^{-2} + (\frac{1}{2})^{2} \xi^{-3} + \cdots) - (\xi^{-1} + 2 \xi^{-2} + 2^{2} \xi^{-3} + \cdots)$$

$$\left(2^{5}\xi^{-1} + 2^{-1}\xi^{-2} + 2^{-2}\xi^{-5} + \cdots\right) - \left(\left(\frac{1}{2}\right)^{6}\xi^{-1} + \left(\frac{1}{2}\right)^{-1}\xi^{-1} + \left(\frac{1}{2}\right)^{2}\xi^{-3} + \cdots\right)$$

$$N = -1 \qquad N = -2 \qquad N = -3$$

$$N = -1 \qquad N = -2 \qquad N = -3$$

$$-A = \frac{3}{2} \frac{-2^2}{(2-2)(2-0.5)} = \left(\frac{0.52}{(2-0.5)} - \frac{22}{(2-2)}\right)$$

$$|\xi| < 0.5$$
 $f(\xi) = -\frac{\xi}{1-2\xi} + \frac{\xi}{1-0.5\xi} -2^{n-1} + (\frac{1}{2})^{n-1} \quad (n > 1)$

$$|\xi| > 2$$
 $f(\xi) = \frac{0.5}{|-a.5\xi^{-1}|} - \frac{2}{|-2\xi^{-1}|} + 2^{n_1} - (\frac{1}{2})^{n_{-1}}$ $(n < 1)$

$$\frac{\left(\left(\frac{1}{2}\right) + \left(\frac{1}{2}\right)^{2} \xi^{-1} + \left(\frac{1}{2}\right)^{3} \xi^{-2} + \cdots\right) + \left(\left(\frac{1}{2}\right)^{-1} + \left(\frac{1}{2}\right)^{-2} \xi^{-1} + \left(\frac{1}{2}\right)^{-3} \xi^{-2} + \cdots\right)}{\left(2 + 2^{2} \xi^{-1} + 2^{-3} \xi^{-2} + \cdots\right) + \left(2 + 2^{2} \xi^{-1} + 2^{3} \xi^{-2} + \cdots\right)}$$

$$(z)$$
 $|z| < 0.5$ $|z| > 2$ anticausal causal

$$|\xi| < 0.5$$
 $\chi(\xi) = -\frac{2}{1-2\xi} + \frac{0.5}{1-0.5\xi} -(\frac{1}{2})^{n-1} + 2^{n-1}$ $(n < 1)$

$$-\left(2^{i}\xi^{0}+2^{2}\xi^{1}+2^{3}\xi^{2}+\cdots\right)+\left(\frac{1}{2}\xi^{0}+\left(\frac{1}{2}\xi^{0}+\left(\frac{1}{2}\xi^{0}+2^{3}\xi^{1}+\cdots\right)\right)\right)$$
$$-\left(\left(\frac{1}{2}\xi^{0}+\left(\frac{1}{2}\xi^{0}+\left(\frac{1}{2}\xi^{0}+2^{3}\xi^{1}+\cdots\right)+\left(2^{-1}\xi^{0}+2^{-2}\xi^{1}+2^{3}\xi^{0}+\cdots\right)\right)\right)$$

n=0 n=-1 n=-2 n=0 n=-1 n=-2

$$|\xi| > 2$$
 $\chi(\xi) = \frac{\xi^{-1}}{|-\rho.5\xi^{-1}|} - \frac{\xi^{-1}}{|-\rho.5\xi^{-1}|} + (\frac{1}{2})^{n-1} - 2^{n-1}$ $(n \ge 1)$

$$\frac{U=1}{\left(\frac{7}{17} \right)_{0}^{2} \xi_{-1} + \left(\frac{7}{17} \right)_{1}^{2} \xi_{-2} + \left(\frac{7}{17} \right)_{2}^{2} \xi_{-2} + \cdots } - \left(\frac{7}{17} \right)_{0}^{2} \xi_{-1} + \frac{1}{17} \left(\frac{7}{17} \right)_{1}^{2} \xi_{-2} + \frac{1}{17} \left(\frac{7}{17} \right)_{1}^{2} \xi_{-2} + \cdots }$$

$$-\mathbf{B}^{\frac{3}{2}}\frac{-\mathbf{z}^{2}}{(2-2)(2-0.5)} = \frac{0.52}{(2-0.5)} - \frac{22}{(2-1)}$$

$$|\xi| < 0.5$$
 $\chi(\xi) = -\frac{\xi}{|-2\xi|} + \frac{\xi}{|-0.5\xi|} -(\frac{1}{2})^{\eta+1} + 2^{\eta+1}$ $(\eta < 0)$

$$- \left(\frac{1}{4} \right)_{0} \xi + \left(\frac{1}{4} \right)_{-1} \xi_{2} + \left(\frac{1}{4} \right)_{-2} \xi_{3} + \cdots \right) + \left(\frac{1}{4} \right) \xi_{3} + \left(\frac{1}{4} \right)_{5} \xi_{3} + \cdots \right)$$

$$- \left(\frac{1}{4} \right)_{0} \xi + \left(\frac{1}{4} \right)_{-1} \xi_{3} + \left(\frac{1}{4} \right)_{-2} \xi_{3} + \cdots \right) + \left(\frac{1}{4} \right) \xi_{3} + \cdots \right)$$

$$|\xi| > 2$$
 $|\xi| > 2$ $|-as \epsilon^{-1}| - \frac{2}{|-as \epsilon^{-1}|} + \frac{1}{2} |-as \epsilon^{-1}| + \frac{1}{2} |-as \epsilon^{-1}|$

$$\frac{\left(\frac{1}{2}\right) + \left(\frac{1}{2}\right)^{2} \xi^{-1} + \left(\frac{1}{2}\right)^{3} \xi^{-2} + \cdots + \left(2 + 2^{2} \xi^{-1} + 2^{3} \xi^{-2} + \cdots \right)}{n = 0 \quad n = 1 \quad n = 2}$$

$$\frac{3}{2} \frac{-1}{(2-05)(2-2)} = \begin{pmatrix} \frac{1}{2-0.5} - \frac{1}{2-2} \end{pmatrix}$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| < 0.5 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2 \quad |\xi| > 2$$

$$|\xi| > 2 \quad |\xi| > 2$$

$$\frac{3}{2} \frac{-1}{(2-0.5)(2-2)} = \left(\frac{1}{\xi-0.5} - \frac{1}{\xi-2}\right)$$

$$|\xi| < 0.5 \qquad f(z) = -\frac{2}{1-2\xi} + \frac{0.5}{1-0.5\xi} -2^{n+1} + (\frac{1}{2})^{n+1} \qquad (n > 0)$$

$$-\left(2z^{0} + 2^{1} \xi^{1} + 2^{3} \xi^{2} + \cdots\right) + \left((\frac{1}{2})z^{0} + (\frac{1}{2})^{3} \xi^{1} + (\frac{1}{2})^{3} \xi^{1} + \cdots\right)$$

$$|\xi| < 0.5 \qquad \chi(z) = -\frac{2}{1-2\xi} + \frac{0.5}{1-0.5\xi} -(\frac{1}{2})^{n-1} + 2^{n-1} \qquad (n < 0)$$


$$|\xi| < 0.5 \qquad \chi(\xi) = -\frac{2}{1 - 2\xi} + \frac{0.5}{1 - 0.5\xi} - (\frac{1}{2})^{n-1} + 2^{n-1} \qquad (n < 0)$$


$$- (2'\xi'' + 2^2\xi' + 2^3\xi^2 + \cdots) + ((\frac{1}{2})\xi'' + (\frac{1}{2})^3\xi^1 + (\frac{1}{2})^3\xi^1 + \cdots)$$

$$- ((\frac{1}{2})^1 \underline{z}^0 + (\frac{1}{2})^2 \xi^1 + (\frac{1}{2})^3 \xi^1 + \cdots) + (2^{-1}\underline{z}^0 + 2^{-2}\xi^1 + 2^3\xi^2 + \cdots)$$


$$= 0 \quad n = -1 \quad n = -2$$

$$= 0.5 \quad (n < 0)$$





$$\frac{3}{3} \frac{-1}{(2-0.5)(2-2)} = \left(\frac{\xi-0.5}{1} - \frac{1}{\xi-2}\right)$$




$$\int (\xi) = (-2) \frac{0.5}{0.5 - \xi} + (0.5) \frac{2}{2 - \xi} \qquad \left(|\xi| < 0.5 \right)$$

$$a_n = (-2) \ 2^n + (0.5) \ (\frac{1}{2})^n \ (n \ge 0)$$

$$-2^{n+1} + (\frac{1}{2})^{n+1}$$

$$\frac{3}{2} \frac{-2^{2}}{(2-2)(2-0.5)} = \left(\frac{0.52}{(2-0.5)} - \frac{22}{(2-1)}\right) \qquad |2| > 2$$



$$X(z) = 0.5 \frac{z}{z-0.5} - 2 \frac{z}{z-1} \qquad (|z| > 2)$$




$$Q_n = (0.5) \left(\frac{1}{2}\right)^n - 2 \cdot 2^n \qquad (n \geqslant 0)$$

$$\left(\frac{1}{2}\right)^{n+1} - 2^{n+1}$$

Anti-Causal
$$f(z)$$
 $X(z)$ $|z| > 2$ $|z| < 0.5$

$$\frac{3}{2} \frac{-1}{(2-0.5)(2-2)} = \left(\frac{2-0.5}{2-0.5} - \frac{1}{2-2}\right)$$


|2| >2 |2| >0.5



$$\int (\xi) = (-2) \frac{-0.5}{0.5 - \xi} + (0.5) \frac{-2}{2 - \xi} \qquad (|\xi| > 0.5)$$

$$a_n = (+2) \ 2^n - (0.5) \ (\frac{1}{2})^n \ (n < 0)$$
 $+2^{n+1} - (\frac{1}{2})^{n+1}$

$$\frac{3}{2} \frac{-2^{2}}{(2-2)(2-0.5)} = \left(\frac{0.52}{(2-0.5)} - \frac{22}{(2-1)}\right) \qquad |2| < 2$$



$$X(z) = 0.5 \frac{-\xi}{\xi - 0.5} - 2 \frac{-\xi}{\xi - \lambda} \qquad (|z| < 2)$$



$$\alpha_n = -(0.5)(\frac{1}{2})^n + 2 \cdot 2^n \qquad (n < 0)$$

$$-(\frac{1}{2})^{n+1} + 2^{n+1}$$

$$\bigcirc -\bigcirc = \frac{3}{2} \frac{(3-0.5)(3-2)}{(3-2)} = \boxed{f(3)}$$


$$\frac{3}{2} \frac{-1}{(2-0.5)(2-2)} = \left(\frac{1}{\xi-0.5} - \frac{1}{\xi-2}\right) = \frac{-2}{1-2\xi} + \frac{0.5}{1-0.5\xi}$$

$$f(\bar{z}) = \frac{\frac{(-2)}{1 - (2\bar{z})} + \frac{(\frac{1}{2})}{1 - (\frac{2}{2})}}{= -\sum_{n=0}^{\infty} (2)^{n+i} (\bar{z})^n + \sum_{n=0}^{\infty} (\frac{1}{2})^{n+i} (\bar{z})^n}$$
$$= -\sum_{n=0}^{\infty} (2)^{n+i} \bar{z}^n + \sum_{n=0}^{\infty} (\frac{1}{2})^{n+i} \bar{z}^n$$

$$a_n =$$

$$a_n = -2^{n+i} + \left(\frac{1}{2}\right)^{n+i}$$

$$\frac{3}{2} \frac{-1}{(2-0.5)(2-2)} = \left(\frac{1}{\xi-0.5} - \frac{1}{\xi-2}\right) = \frac{\xi^{-1}}{1-0.5\xi^{-1}} - \frac{\xi^{-1}}{1-2\xi^{-1}}$$



$$\frac{f(\xi)}{f(\xi)} = \frac{\left(\frac{1}{\xi}\right)}{1 - \left(\frac{1}{2\xi}\right)} - \frac{\left(\frac{1}{\xi}\right)}{1 - \left(\frac{3}{\xi}\right)}$$

$$= \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n \left(\frac{1}{\xi}\right)^{n+1} - \sum_{n=0}^{\infty} \left(2\right)^n \left(\frac{1}{\xi}\right)^{n+1}$$

$$= \sum_{n=1}^{\infty} \left(\frac{1}{2}\right)^{n-1} \xi^{-n} - \sum_{n=1}^{\infty} \left(2\right)^{n-1} \xi^{-n}$$

$$= \sum_{n=-1}^{-\infty} (2)^{n+1} \xi^n - \sum_{n=-1}^{-\infty} (\frac{1}{2})^{n+1} \xi^n$$

$$a_n$$

$$\frac{3}{2} \frac{-1}{(2-0.5)(2-2)} = \boxed{\chi(3)} \quad |z| < 0.5 \quad |z| > 2$$
anticausal causal

$$|z| < 0.5$$
 $|z| > 2$

anticausal Causal


$$\frac{3}{2} \frac{-1}{(2-0.5)(2-2)} = \left(\frac{1}{\xi-0.5} - \frac{1}{\xi-2}\right) = \frac{-2}{1-2\xi} + \frac{0.5}{1-0.5\xi}$$

$$\begin{array}{c} \times (\overline{z}) = \frac{\left(-2\right)}{1 - \left(2\overline{z}\right)} + \frac{\left(\frac{1}{a}\right)}{1 - \left(\frac{2}{a}\right)} \\ = -\sum\limits_{n=0}^{\infty} \left(2\right)^{n+i} (\overline{z})^n + \sum\limits_{n=0}^{\infty} \left(\frac{1}{2}\right)^{n+i} (\overline{z})^n \\ = -\sum\limits_{n=0}^{\infty} \left(2\right)^{n+i} \overline{z}^n + \sum\limits_{n=0}^{\infty} \left(\frac{1}{2}\right)^{n+i} \overline{z}^n \end{array}$$

$$= -\sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^{n-1} \xi^{-n} + \sum_{n=0}^{\infty} (2)^{n-1} \xi^{-n}$$

$$(n \le 0)$$
 $a_n = -(\frac{1}{2})^{n-1} + 2^{n-1}$

$$\frac{3}{2} \frac{-1}{(2-05)(2-2)} = \left(\frac{1}{\xi-0.5} - \frac{1}{\xi-2}\right) = \frac{\xi^{-1}}{1-0.5\xi^{-1}} - \frac{\xi^{-1}}{1-2\xi^{-1}}$$



$$(n > 0)$$
 $a_n = (\frac{1}{2})^{n-1} - (2)^{n-1}$

$$(2)-\triangle \frac{3}{2}\frac{-\xi^{2}}{(2-2)(2-0.5)} = \int (3) \frac{|\xi| < 0.5}{\text{causal}} \frac{|\xi| > 2}{\text{anticausal}}$$


$$\frac{3}{2} \frac{-2^{2}}{(2-2)(2-0.5)} = \left(\frac{0.52}{(2-0.5)} - \frac{22}{(2-1)}\right) = -\frac{2}{1-22} + \frac{2}{1-0.52}$$

$$\frac{1}{1 - (2\xi)} = -\frac{(\xi)}{1 - (2\xi)} + \frac{(\xi)}{1 - (\frac{\xi}{2})} \neq \frac{1}{1 - (\frac{\xi}{2})} = -\sum_{n=0}^{\infty} (2)^n (\xi)^{n+1} + \sum_{n=0}^{\infty} (\frac{1}{2})^n (\xi)^{n+1} = -\sum_{n=0}^{\infty} (2)^{n-1} \xi^n + \sum_{n=0}^{\infty} (\frac{1}{2})^{n-1} \xi^n$$

$$= -\sum_{n=0}^{\infty} (2)^{n-1} \xi^n + \sum_{n=0}^{\infty} (\frac{1}{2})^{n-1} \xi^n$$

$$(n > 0)$$
 $a_n = -2^{n-1} + (\frac{1}{2})^{n-1}$

$$\frac{3}{2} \frac{-2}{(2-2)(2-0.5)} = \left(\frac{0.52}{(2-0.5)} - \frac{22}{(2-1)}\right) = \frac{0.5}{1-0.52^{-1}} - \frac{2}{1-22^{-1}}$$



$$(n \leq 0) \qquad a_n = 2^{n-1} - \left(\frac{1}{2}\right)^{n-1}$$

$$-\left(\frac{1}{2}\right)^{n-1}$$

(2) - (B)
$$\frac{3}{2} \frac{-\xi^2}{(2-2)(2-0.5)} = [\chi(\xi)]$$

$$|z| < 0.5$$
 $|z| > 2$

anticausal causal

$$\frac{3}{2} \frac{-2^{2}}{(2-2)(2-0.5)} = \left(\frac{0.52}{(2-0.5)} - \frac{22}{(2-1)}\right) = -\frac{2}{1-22} + \frac{2}{1-0.52}$$

(n < 0) $a_n = -(\frac{1}{2})^{n+1} + 2^{n+1}$

$$\frac{3}{2} \frac{-2^{2}}{(2-2)(2-0.5)} = \left(\frac{0.52}{(2-0.5)} - \frac{22}{(2-1)}\right) = \frac{0.5}{1-0.52^{-1}} - \frac{2}{1-22^{-1}}$$

$$X(\xi) = \frac{\left(\frac{1}{2}\right)}{1 - \left(\frac{1}{2\xi}\right)} - \frac{\left(\frac{2}{2}\right)}{1 - \left(\frac{2}{\xi}\right)}$$

$$= \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^{n+1} \left(\frac{1}{\xi}\right)^n - \sum_{n=0}^{\infty} \left(2\right)^{n+1} \left(\frac{1}{\xi}\right)^n$$

$$= \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^{n+1} \xi^{-n} - \sum_{n=0}^{\infty} \left(2\right)^{n+1} \xi^{-n}$$

$$(n \geqslant 0) \qquad \alpha_n = \left(\frac{1}{2}\right)^{n+1} - 2^{n+1}$$

