

# Statistical Inference Overview

Young W. Lim

2024-06-01 Sat

1 Based on

2 Overview

- Statistical Inference
- Types of Hypothesis Tests

## "Understanding Statistics in the Behavioral Sciences" R. R. Pagano

I, the copyright holder of this work, hereby publish it under the following licenses: GNU head Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.2 or any later version published by the Free Software Foundation; with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts. A copy of the license is included in the section entitled GNU Free Documentation License.

CC BY SA This file is licensed under the Creative Commons Attribution ShareAlike 3.0 Unported License. In short: you are free to share and make derivative works of the file under the conditions that you appropriately attribute it, and that you distribute it only under a license compatible with this one.

# Populations and Samples (1)

- **population**: everything in the group that we want to learn about.
- **sample**: a part of the population.
- Examples of populations and a sample from those populations:

Population	Sample
All of the people in Germany	500 Germans
All of the customers of Netflix	300 Netflix customers
Every car manufacturer	Tesla, Toyota, BMW, Ford

[https://www.w3schools.com/statistics/statistics\\_statistical\\_inference.php](https://www.w3schools.com/statistics/statistics_statistical_inference.php)

## Populations and Samples (2)

- For good statistical analysis, the **sample** needs to be as similar as possible to the population.
- If they are similar enough, we say that the **sample** is representative of the population.
- The **sample** is used to make conclusions about the whole **population**.

[https://www.w3schools.com/statistics/statistics\\_statistical\\_inference.php](https://www.w3schools.com/statistics/statistics_statistical_inference.php)

## Populations and Samples (3)

- If the **sample** is not similar enough to the whole **population**, the **conclusions** could be useless.
- Many words have specific meanings in statistics.
- The word **population** normally refers to a group of people.
- In statistics, it is any specific group that we are interested in learning about.

[https://www.w3schools.com/statistics/statistics\\_statistical\\_inference.php](https://www.w3schools.com/statistics/statistics_statistical_inference.php)

- Using data analysis and statistics to make conclusions about a population is called **statistical inference**.
- The main types of statistical inference are:
  - **Estimation**
  - **Hypothesis testing**

[https://www.w3schools.com/statistics/statistics\\_statistical\\_inference.php](https://www.w3schools.com/statistics/statistics_statistical_inference.php)

# Estimation (1)

- Statistics from a sample are used to **estimate** population parameters.
- *The most likely* value is called a point estimate.
- There is always **uncertainty** when estimating.

[https://www.w3schools.com/statistics/statistics\\_statistical\\_inference.php](https://www.w3schools.com/statistics/statistics_statistical_inference.php)



## Estimation (2)

- The uncertainty is often expressed as **confidence intervals** defined by a *likely* lowest and highest value for the **parameter**.
- An example could be a **confidence interval** for the number of bicycles a Dutch person owns:
  - The **average number** of bikes a Dutch person owns is between 3.5 and 6.

[https://www.w3schools.com/statistics/statistics\\_statistical\\_inference.php](https://www.w3schools.com/statistics/statistics_statistical_inference.php)

# Hypothesis Testing (1)

- a method to check if a claim about a population is true.
- checks how likely it is that a **hypothesis** is true is based on the sample data.
- different types of hypothesis testing.
- the steps of the test depends on:
  - Type of data (categorical or numerical)
  - If you are looking at:
    - A single group
    - Comparing one group to another
    - Comparing the same group before and after a change

[https://www.w3schools.com/statistics/statistics\\_statistical\\_inference.php](https://www.w3schools.com/statistics/statistics_statistical_inference.php)

# Hypothesis Testing (2)

- A hypothesis is a claim about a population parameter.
- A hypothesis test is a formal procedure to check if a hypothesis is true or not.
- Examples of claims that can be checked:
  - The average height of people in Denmark is more than 170 cm.
  - The share of left handed people in Australia is not 10%.
  - The average income of dentists is less the average income of lawyers.

[https://www.w3schools.com/statistics/statistics\\_hypothesis\\_testing.php](https://www.w3schools.com/statistics/statistics_hypothesis_testing.php)

# The Null and Alternative Hypothesis

- Hypothesis testing is based on making two different claims about a population parameter.
- The null hypothesis ( $H_0$ ) and the alternative hypothesis ( $H_1$ ) are the claims.
- The two claims needs to be mutually exclusive, meaning only one of them can be true.
- The alternative hypothesis is typically what we are trying to prove.
- For example, we want to check the following claim:
  - "The average height of people in Denmark is more than 170 cm."

[https://www.w3schools.com/statistics/statistics\\_hypothesis\\_testing.php](https://www.w3schools.com/statistics/statistics_hypothesis_testing.php)

# Summary (1) comparing means

---

tests

---

- **one-sample** test    comparing sample mean, population mean
  - **two-sample** test    comparing two independent sample means
  - **paired** test        comparing two related sample means
- 

---

tests

test conditions

---

- **t-test**            1. when the **population variance** is *known*  
                          2. when the **sample size** is *large*
  - **z-test**            1, when the **population variance** is *unknown*  
                          2. the **sample size** is *small*
- 

<https://www.qualitygurus.com/common-types-of-hypothesis-tests/>

## Summary (2) comparing means

one sample z-test	<u>sample mean</u> , <u>population mean</u> <u>known population var</u> / <u>large sample size</u>
one sample t-test	<u>sample mean</u> , <u>population mean</u> <u>unknown population var</u> / <u>small sample size</u>
two sample z-test	two <i>independent</i> <u>sample means</u> <u>known population var</u> / <u>large sample size</u>
two sample t-test	two <i>independent</i> <u>sample means</u> <u>unknown population var</u> / <u>small sample size</u>
paired t-test	two <i>related</i> <u>sample means</u> <u>unknown population var</u> / <u>small sample size</u>

<https://www.qualitygurus.com/common-types-of-hypothesis-tests/>

## Summary (3) comparing proportions

---

one sample **proportion test**     sample proportion, population proportion  
when  $np \geq 10$  and  $n(1 - p) \geq 10$

---

two sample **proportion test**     two *independent* sample proportions  
when  $np \geq 10$  and  $n(1 - p) \geq 10$

---

---

test conditions

---

the normal approximation is used

when both  $np \geq 10$  and  $n(1 - p) \geq 10$

(data should have at least 10 "successes" and at least 10 "failures" )

---

<https://www.qualitygurus.com/common-types-of-hypothesis-tests/>

## Summary (4)

---

compare variances between

---

sample variance, known population variance      Chi-square test

---

two independent sample variances      F-test

---

observed frequencies, expected frequencies      goodness of fit test

---

observed frequencies, expected frequencies      contingency tables

---

means of three or more independent samples      ANOVA (Analysis of Variance)

---

<https://www.qualitygurus.com/common-types-of-hypothesis-tests/>



# Tests for Comparing Means (1)

- One-sample **z-test**:
  - used to compare the **mean** of a sample to a known population mean
  - used when the population variance is known, or the sample size is *large* ( $n > 30$ ).
- Two-sample **z-test**:
  - used to compare the **means** of two independent samples.
  - used when the population variances are known, or the sample sizes are *large* ( $n > 30$ ).

<https://www.qualitygurus.com/common-types-of-hypothesis-tests/>

# Tests for Comparing Means (2)

- One-sample **t-test**:
  - used to compare the **mean** of a sample to a known population mean.
  - used when the population variance is unknown, and the sample size is *small* ( $n < 30$ ).
- Two-sample **t-test**:
  - used to compare the **means** of two independent **samples**.
  - used when the population variances are unknown, and the sample sizes are *small* ( $n < 30$ ).

<https://www.qualitygurus.com/common-types-of-hypothesis-tests/>

# Tests for Comparing Means (3)

- Paired **t-test**:
  - used to compare the **means** of two *related* samples, such as the before and after measurements of the same group of subjects.
  - used when the population **variances** are unknown, and the sample size is *small* ( $n < 30$ ).

<https://www.qualitygurus.com/common-types-of-hypothesis-tests/>

# Tests for Comparing Proportions (1)

- Let us consider the parameter  $p$  of the **population proportion**
  - eg) we might want to know the **proportion** of males within a total population of adults when we conduct a survey.
- A **test of proportion** will assess whether or not a **sample** from a **population** represents the true **proportion** of the entire **population**

<https://online.stat.psu.edu/statprogram/reviews/statistical-concepts/proportions>

# Examples of proportions

- an example
  - newborn babies are more likely to be boys than girls.
  - a random sample found 13,173 boys were born among 25,468 newborn children
  - the **sample proportion** of boys was 0.5172 ( $= \frac{13173}{25468}$ )
  - is this sample evidence that the birth of boys is more common than the birth of girls in the entire population? ( $0.5172 > 0.4828$ )

<https://online.stat.psu.edu/statprogram/reviews/statistical-concepts/proportions>

# Tests for Comparing Proportions (2-1)

- examples involved testing whether a single population **proportion**  $p$  equals some value .
- different examples of testing whether one population **proportion** equals a second population **proportion**

<https://online.stat.psu.edu/stat415/lesson/9/9.4>

# Tests for Comparing Proportions (2-2)

- Additionally, most of our examples thus far have involved left-tailed tests in which the **alternative hypothesis** involved or right-tailed tests in which the **alternative hypothesis** involved
- Here, let's consider an example that tests the equality of two **proportions** against the **alternative** that they are not equal.

<https://online.stat.psu.edu/stat415/lesson/9/9.4>

## Tests for Comparing Proportions (2-3)

- Time magazine reported the result of a telephone poll of 800 adult Americans.
- The question posed of the Americans who were surveyed was: "Should the federal tax on cigarettes be raised to pay for health care reform?"
- The results of the survey were:

Non-smokers	Smokers
$n_1 = 605$	$n_2 = 195$
$y_1 = 351$ said yes	$y_2 = 41$ said yes
$\hat{p}_1 = \frac{351}{605} = 0.58$	$\hat{p}_2 = \frac{41}{195} = 0.21$

<https://online.stat.psu.edu/stat415/lesson/9/9.4>



# Tests for Comparing Proportions (3)

- One-sample **proportion test** :
  - used to compare the **proportion** of a sample to a known population proportion.
  - the normal approximation is used when both  $np \geq 10$  and  $n(1 - p) \geq 10$  (data should have at least 10 "successes" and at least 10 "failures" ) (in some books, it is 5)

<https://www.qualitygurus.com/common-types-of-hypothesis-tests/>

# Tests for Comparing Proportions (4)

- Two-sample **proportion test** :
  - used to compare the proportions of two independent samples.
  - the normal approximation is used  
when both  $np \geq 10$  and  $n(1 - p) \geq 10$   
(data should have at least 10 "successes" and at least 10 "failures" )  
(in some books, it is 5)

<https://www.qualitygurus.com/common-types-of-hypothesis-tests/>

# Tests for Comparing Variance

- **Chi-square test** for variance :
  - used to compare the **variance** of a sample to a known population variance
- **F-test** for variance :
  - used to compare the **variances** of two *independent samples*

<https://www.qualitygurus.com/common-types-of-hypothesis-tests/>

# Other Common Tests (1)

- **Goodness of fit** test :
- used to determine whether a sample fits a *specific* **distribution**.
- used to compare the observed frequencies of a *categorical variable* to the expected frequencies under a *particular* **distribution**.

<https://www.qualitygurus.com/common-types-of-hypothesis-tests/>

## Other Common Tests (2)

- Testing for **independence of two attributes (Contingency Tables)** :
- used to determine whether there is a relationship between two *categorical variables*.
- often used in the form of a **chi-square** test, which compares the observed frequencies in a **contingency table** to the expected frequencies under the assumption of independence.

<https://www.qualitygurus.com/common-types-of-hypothesis-tests/>

## Other Common Tests (3)

- ANOVA (Analysis of Variance) :
- used to compare the **means** of three or more independent samples.
- used to determine whether there is a significant difference between the **means** of the groups.

<https://www.qualitygurus.com/common-types-of-hypothesis-tests/>

# One-sample z-test

- used to test a hypothesis about the *population mean*
- based on the assumption that the sample is drawn from a **normally distributed** population.
  - the **null hypothesis**  
the *population mean* is equal to a specific value
  - the **alternative hypothesis**  
the *population mean* is not equal to that value

<https://www.qualitygurus.com/common-types-of-hypothesis-tests/>

# Two-sample z-test

- based on the assumption that both samples are drawn from **normally distributed** populations with equal **variances**.
- the two-sample **z-test** requires that the population **standard deviations** be known or that the sample **sizes** be *large* (30 or more),
  - the **null hypothesis**  
the **means** of the two samples are equal
  - the **alternative hypothesis**  
the **means** are not equal

<https://www.qualitygurus.com/common-types-of-hypothesis-tests/>



# One-sample t-test

- used to test a hypothesis about the *population mean*
- based on the assumption that the sample is drawn from a **normally distributed** population
  - the **null hypothesis**  
the *population mean* is equal to a specific value
  - the **alternative hypothesis**  
the *population mean* is not equal to that value

<https://www.qualitygurus.com/common-types-of-hypothesis-tests/>

# Two-sample **t-test**

- based on the assumption that the samples are drawn from populations with **normal distributions**.
- the two-sample **t-test**  
that the population **standard deviations** need not be known or  
that the sample **sizes** need not be large (30 or more),
  - the **null hypothesis**  
the **means** of the two samples are equal
  - the **alternative hypothesis**  
the **means** are not equal

<https://www.qualitygurus.com/common-types-of-hypothesis-tests/>

# Paired t-test

- used to test a hypothesis about the difference between the means of the two samples
- based on the assumption that the differences between the pairs are **normally distributed**
- In a dependent two-sample t-test (a paired t-test), the samples in the two groups being compared are *related* in some way.
  - the **null hypothesis**  
there is no difference between the means of the two samples
  - the **alternative hypothesis**  
there is a difference between the means

<https://www.qualitygurus.com/common-types-of-hypothesis-tests/>

# Two proportions **z-test**

- used to test a hypothesis about the difference between the proportions of the two samples and
- based on the assumption that the samples are drawn from populations with a **normal distribution**
  - the **null hypothesis** :  
there is no difference between the proportions of the two samples
  - the **alternative hypothesis** :  
there is a difference between the proportion

<https://www.qualitygurus.com/common-types-of-hypothesis-tests/>

