

## Multiple Linear Regression II



Image source: <https://commons.wikimedia.org/wiki/File:Autobunnskr1/C3/1A4i2-RO-A201.jpg>

### Lecture 8

Survey Research & Design in Psychology  
James Neill, 2017  
[Creative Commons Attribution 4.0](#)

## Overview



1. Summary of MLR I lecture
2. Semi-partial correlations
3. Residual analysis
4. Interactions
5. Analysis of change
6. Writing up an MLR
7. Summary of MLR II
8. MLR II quiz – Practice questions 2

## Readings

1. As per previous lecture, plus
2. **Howitt & Cramer (2014)**
  - Ch 39: Moderator variables and relationships between two variables

3

## Summary of MLR I lecture

### Purpose of MLR:

To examine linear relations between:

- two or more predictors (IVs; X) and
- a single outcome variable (DV; Y)

### Model:

Develop a theoretical model:

- Path diagram and/or Venn diagram
- Express as one hypothesis per IV

4

## Summary of MLR I

### Check assumptions:

- Level of measurement
- Sample size
- Normality
- Linearity
- Homoscedasticity
- Collinearity
- Multivariate outliers
- Residuals should be normally distributed

5

## Summary of MLR I

### Choose type:

- Standard
- Hierarchical
- Stepwise, Forward, Backward

6

## Summary of MLR I

### Interpret output:


- **Overall:**
  - $R$ ,  $R^2$ , Adjusted  $R^2$
  - Changes in  $R^2$  (if hierarchical)
  - Significance ( $F$ ,  $p$ )
- **For each IV:**
  - Standardised coefficient
    - size, direction and significance
  - Unstandardised coefficient
    - report equation (if useful)
  - Semi-partial correlations ( $sr^2$ )

7

## Semi-partial correlation

Venn Diagrams depict variance and shared variance:

- $a + b + c + d =$  variance in DV
- $a + b + c = R$  (variance in the DV explained by IV1 and IV2)
- $a + c =$  uniquely explained variance
- $b =$  non-uniquely explained variance



$a$  and  $c$  are **semi-partial correlations ( $sr$ )**:

- $a = sr$  between IV1 and DV after controlling for (partialling out) the influence of IV2
- $c = sr$  between IV2 and DV after controlling for (partialling out) the influence of IV1

8

## Semi-partial correlations in MLR

When interpreting MLR coefficients:

- Draw a Path diagram or Venn diagram
- Compare 0-order ( $r$ ) and semi-partial correlations ( $sr$ ) for each IV to help understand relations amongst the IVs and the DV:
  - A semi-partial correlation ( $sr$ ) will be less than or equal to the correlation ( $r$ )
  - If a  $sr$  equals a  $r$ , then the IV independently predicts the DV
  - To the extent that a  $sr$  is less than the  $r$ , the IV's explanation of the DV is shared with other IVs
  - An IV may have a significant  $r$  with the DV, but a non-significant  $sr$ . This indicates that the unique variance explained by the IV may be 0.
- Compare the relative importance of the predictors using betas and/or  $srs$

9


## Semi-partial correlations ( $sr$ ) in MLR

- SPSS provides semi-partial correlations if selected (labelled “part”)
- Square  $sr$  to get  $sr^2$ .
- The  $sr^2$  indicates the % of variance in the DV which is uniquely explained by an IV..
- Compare each  $sr^2$  with the  $r^2$  – do they differ – why?

10

## Semi-partial correlations for MLR in SPSS

Analyze - Linear Regression - Statistics - Part and partial correlations



## Semi-partial correlations for MLR in SPSS - Example

Coefficient <sup>a</sup>				
Correlations				
Model		Zero-order	Partial	Part
1	Zscore: GA: General Ability Total	.285	.107	.090
	Zscore: PSA: Perceptual Speed Ability Total	.419	.319	.280
	Zscore: PMA: Psychomotor Ability Total	.275	.034	.028
	QBK: Total Knowledge of Text Editing Keys (% correct)	.386	.368	.329

a. Dependent Variable: Typing Test: Speed (Words Per Minute)

Semi-partial correlations ( $sr$ ) indicate the relative importance of each of the predictors;  $sr^2$  indicate the % of variance uniquely explained by each predictor.

## Summary: Semi-partial correlation ( $sr$ )

1. In MLR,  $sr$  is labelled "part" in the regression coefficients table SPSS output
2. Square these values to obtain  $sr^2$ , the unique % of DV variance explained by each IV
3. Discuss the extent to which the explained variance in the DV is due to unique or shared contributions of the IVs

13


## Semi-partial correlations ( $sr$ ) in MLR

- $R^2 = .34$
- 16% is of the DV variance is uniquely explained by the IVs (sum of  $sr^2$ s)
- 18% of the DV variance is explained by the combination of the IVs ( $R^2$  - sum of  $sr^2$ s)

Independent variables	$sr^2$
Not Coping (NC)	.07
Somatic Coping (SC)	.06
Wishful Thinking (WT)	.02
Worry (W)	.01

14

## Residual analysis



## Residual analysis

Assumptions about residuals:

- Sometimes positive, sometimes negative but, on average, 0
- Error is random
- Normally distributed about 0

16

## Residual analysis

Analyze - Linear Regression - Plots - Y and X

Linear Regression: Plots

DEPENDENT: \*ZPRED, \*ZRESID, \*DRESID, \*ADJPRED, \*SRESID, \*SDRESID

Scatter 1 of 1

Y: \*ZRESID

X: \*ZPRED

Standardized Residual Plots


Histogram

Normal probability plot

Produce all partial plots

Buttons: Previous, Next, Continue, Cancel, Help

## Residual analysis: Histogram




"The **Normal P-P (Probability) Plot of Regression Standardized Residuals** can be used to assess the assumption of normally distributed residuals. **If the points cluster reasonably tightly along the diagonal line (as they do here), the residuals are normally distributed.** Substantial deviations from the diagonal may be cause for concern."


Allen & Bennett, 2008, p. 183

## Residual analysis: NPP

Normal P-P Plot of Regression Standardized Residual




## Residual analysis: Histogram and NPP



## Residual analysis: Scatterplot of predicted vs. residual values

The absence of any clear patterns in the spread of points indicates that the MLR assumptions of normality, linearity and homoscedasticity are met. (Allen & Bennett, 2008, p. 183)



## Why the big fuss about residuals?

↑ assumption violation

→

↑ Type I error rate  
(i.e., more false positives)

22

## Why the big fuss about residuals?

- Standard error formulae (used for confidence intervals and sig. tests) **WORK** when residuals are well-behaved.
- If the residuals don't meet assumptions these formulae tend to underestimate coefficient standard errors giving lower  $p$ -values and more Type I errors.

23

## Summary: Residual analysis

1. Residuals are the difference between predicted and observed  $Y$  values
2. MLR assumption is that residuals are normally distributed.
3. Examining residuals also helps to assess:
  1. Linearity
  2. Homoscedasticity

24

## Interactions




Image source: [http://commons.wikimedia.org/wiki/File:Color\\_icon\\_orange.png](http://commons.wikimedia.org/wiki/File:Color_icon_orange.png)

## Interactions

- Additivity - when IVs act independently on a DV they do not interact.
- Alternatively, there may be interaction effects - i.e. the magnitude of the effect of one IV on a DV varies as a function of a second IV.
- Also known as a moderation effect.

26

### Interactions: Example 1

Some drugs interact with each other to reduce or enhance other's effects e.g.,

Pseudoephedrine → ↑ Arousal

Caffeine → ↑ Arousal

Pseudoeph. X Caffeine → ↑↑↑ Arousal



X



27

### Interactions: Example 2

Physical exercise (IV1)  
and

Natural environments (IV2)  
may provide

multiplicative benefits  
in reducing stress e.g.,

Natural environment → ↓ Stress

Physical exercise → ↓ Stress

Natural env. x Phys. ex. → ↓↓↓ Stress

28

### Interactions: Example 3

University student satisfaction (IV1)  
and

Level of coping (IV2)

may predict level of stress e.g.,

Satisfaction → ↓ Stress

Coping → ↓ Stress

Satisfaction x Coping → ↓↓↓ Stress

(Dissatisfaction x Not coping → ↑↑↑ Stress)

29

## Interactions

Test interactions in MLR by computing a cross-product term e.g.,:

- Pseudoephedrine (IV1)
- Caffeine (IV2)
- Pseudoephedrine x Caffeine (IV3)

30

## Interactions

cross-product term

$$Y = b_1x_1 + b_2x_2 + b_{12}x_{12} + a + e$$

- $b_{12}$  is the product of the first two slopes ( $b_1 \times b_2$ )
- $b_{12}$  can be interpreted as the amount of change in the slope of the regression of Y on  $b_1$  when  $b_2$  changes by one unit.

31

## Interactions

- Conduct Hierarchical MLR
  - Step 1:
    - Pseudoephedrine
    - Caffeine
  - Step 2:
    - Pseudo x Caffeine (cross-product)
- Examine  $\Delta R^2$ , to see whether the interaction term explains additional variance above and beyond the additive effects of Pseudo and Caffeine.

32

## Interactions


Possible effects of Pseudo and Caffeine on Arousal:

- None
- Pseudo only ( $\uparrow/\downarrow$ )
- Caffeine only ( $\uparrow/\downarrow$ )
- Pseudo + Caffeine ( $\uparrow\uparrow/\downarrow\downarrow$ )
- Pseudo x Caffeine ( $\uparrow\uparrow\uparrow$  (synergistic))
- Pseudo x Caffeine ( $\downarrow\downarrow\downarrow$  (antagonistic))

33


## Interactions - SPSS example

SPSS – Analyze – Transform – Compute



## Interactions - SPSS example

SPSS – Analyze – Regression – Linear



## Interactions - SPSS example

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate	R Square Change
1	.543 <sup>a</sup>	.295	.288	17.34399	.295
2	.547 <sup>b</sup>	.299	.289	17.33241	.004

a. Predictors: (Constant), Caffeine, Pseudoephedrine

b. Predictors: (Constant), Caffeine, Pseudoephedrine, PseudoCaffeine interaction

Model		Unstandardized Coefficients		Standardized Coefficients		t	Sig.
		B	Std. Error	Beta			
1	(Constant)	57.705	5.109			11.286	.000
	Pseudoephedrine	11.511	1.510	.484		7.625	.000
	Caffeine	4.735	1.780	.162		2.660	.008
2	(Constant)	42.730	14.137			3.023	.003
	Pseudoephedrine	18.137	6.025	.732		3.010	.003
	Caffeine	10.126	5.069	.346		1.998	.047
	PseudoCaffeine interaction	-2.313	2.036	-.380		-1.136	.257

Pseudoephedrine and Caffeine are each significantly, positively associated with Arousal, however **there is no interaction**.

Note: This data was fabricated for demonstration purposes.

## Interactions

- Cross-product interaction terms may be highly correlated (collinear) with the corresponding simple IVs, creating problems with assessing the relative importance of main effects and interaction effects.
- An alternative approach is to conduct separate regressions for each level of one of the IVs e.g.,
  - What is the effect of caffeine on Arousal without pseudoephedrine?
  - What is the effect of caffeine on Arousal with pseudoephedrine?

37

## Summary: Interactions

1. In MLR, IVs may interact to:
  1. Increase the IVs' effect on DV
  2. Decrease the IVs' effect on DV
2. Model interactions using hierarchical MLR:
  1. Step 1: Enter IVs
  2. Step 2: Enter cross-product of IVs
  3. Examine change in  $R^2$

38

## Analysis of Change




Image source: <http://commons.wikimedia.org/wiki/File:PMinspirent.jpg>, CC-by-SA 3.0

## Analysis of change

- MLR can be used to analyse the variance of changes in outcome measures over time (e.g., an intervention study with pre-post tests).
- There are two main approaches:
  1. Standard regression:
    1. Compute post-pre difference (or change) scores in the outcome measures and use these change scores as the DV and all IVs in Step 1 of a standard regression.
  2. Hierarchical MLR:
    1. DV is the post measure
    2. Step 1: "Partial out" the baseline score by entering the pre-score.
    3. Step 2: Seek to explain remaining variance in the Post-test DV by entering the IVs of interest and examining the change in  $R^2$ .

40

## Analysis of change


Example research question:

Does the quality of social support (IV2) explain changes in mental health (MH) between the beginning (IV1) and end (DV) of an intervention ?

41

## Analysis of change: Option 1

Calculate the difference scores which indicate the change in mental health.



Use regression to determine the extent to which the IV can explain changes in the DV.

42

## Analysis of change: Option 2

Step 1:  
Baseline  
Mental Health  
(IV1)

Step 1: partial out baseline mental health from post mental health

Post  
Mental  
Health  
(DV)

Step 2:  
Social  
Support  
(IV2)

Step 2: analyse the effect of Social Support on the change (remaining variance) in Mental Health

43

## Analysis of change: Option 2

### Hierarchical MLR

- DV = Mental health after the intervention
- Step 1
  - IV1 = Mental health before the intervention
- Step 2
  - IV2 = Support from group members

44

## Analysis of change: Option 2

### Results of interest

- Change in  $R^2$  – amount of additional variance in the DV (MH2) explained by the predictor (IV2) added in Step 2 (after having baseline IV1 variance has been accounted for in Step 1)
- Regression coefficients - for the predictor (IV2) in Step 2 – indicates variance explained in the DV (MH2) by the IV (SS) after controlling for baseline MH (MH1) in Step 1

45

## Summary: Analysis of change

1. Analysis of changes over time can be assessed by:
  1. Standard regression
    1. Calculate difference scores (Post-score minus Pre-score) and use as a DV
  2. Hierarchical MLR
    1. Step 1: “Partial out” baseline scores
    2. Step 2: Enter other IVs to help predict variance in changes over time.

46

## Writing up an MLR



Image source: <http://commons.wikimedia.org/wiki/File:Writing.svg>, Public domain

## Writing up an MLR: Intro

- Explain the constructs of interest and why they are important
- Explain how / why the IVs may be related to the DV, drawing on theory and research
- State hypotheses (one per IV)
  - Null hypothesis or
  - Directional hypothesis

48



## Writing up an MLR: Results

- Establish the purpose of the analysis
- Provide univariate descriptive statistics ( $M$ ,  $SD$ , Skewness, Kurtosis) for the IVs and DV
- Describe type of MLR
- Test assumptions
- Examine correlations
- Overall model:  $R^2$ , Adjusted  $R^2$ ,  $F$ ,  $p$
- Explain regression coefficients
- Provide table of correlations and regression coefficients (unstandardised and standardised),  $t$ ,  $p$ , and  $sr^2$  values

49

## Writing up an MLR: Discussion

- Summarise the results in relation to the hypotheses and the overall research question, paying attention to:
  - Direction
  - Size
  - Statistical significance
- Interpret the results in the context of relevant psychological theory and research
- Consider alternative explanations and limitations
- Describe implications and/or recommendations

50

## Table of regression coefficients

Table 2

Standard Multiple Linear Regression of Non-productive Coping Strategies on Psychological Distress Reported by Adolescents During Outdoor Education Programs

Variables	PD <sup>a</sup> (DV)	NC	WT	W	SC	B	$\beta$	$sr^{2b}$
						(unique)		
Not Coping (NC)	.43					.41*	.29	.07
Wishful Thinking (WT)	.38	.36				.15*	.15	.02
Worry (W)	.36	.25	.40			.13*	.13	.01
Somatic Coping (SC)	.42	.22	.30	.38		.26*	.26	.06
Intercept						.60*		

Note. <sup>a</sup> $p < .05$ ; <sup>a</sup>PD = Psychological Distress; <sup>b</sup> $sr^2$  = the squared semipartial correlations indicate the unique variance predicted by the independent variable.

## Summary: Writing up an MLR

1. Introduction
  1. Establish purpose
  2. Describe model and hypotheses
2. Results
  1. Univariate descriptive statistics
  2. Correlations
  3. Type of MLR and assumptions
  4. Regression coefficients
3. Discussion
  1. Summarise and interpret, with limitations
  2. Implications and recommendations

52

## Questions



53

## MLR II Quiz – Practice question 1

In an MLR, if the  $r$  between the two IVs is 1,  $R$  will equal the  $r$  between one of the IVs and the DV.

[Hint: Draw a Venn Diagram]

- a) True
- b) False

54

### MLR II Quiz – Practice question 2

In MLR, if two IVs are somewhat correlated with the DV and with one another, the  $srs$  between the IVs and the DV will be \_\_\_\_\_ in magnitude than the  $rs$ :

[Hint: Draw a Venn Diagram]

- a) Equal
- b) Smaller
- c) Larger
- d) Impossible to tell

55

### MLR II Quiz – Practice question 3

In MLR, the unique variance in the DV explained by a particular IV is estimated by its:

- a) Zero-order correlation square ( $r^2$ )
- b) Multiple correlation coefficient squared ( $R^2$ )
- c) Semi-partial correlation squared ( $sr^2$ )

56

### MLR II Quiz – Practice question 4

Interaction effects can be tested in MLR by using IVs that represent:

- a) Cross-products between the IVs and DV in a hierarchical regression
- b) Cross-products of IVs
- c) Semi-partial correlations squared ( $sr^2$ )

57

### MLR II Quiz – Practice question 5

To assess the extent to which social support from group members during an intervention program explain changes in mental health (MH) between the beginning and end of an intervention program, what MLR design could be used?

- a) Hierarchical with pre-MH in Step 1
- b) Hierarchical with cross-products of IVs in Step 2

58

### References

- Allen, P. & Bennett, K. (2008). *SPSS for the health and behavioural sciences*. South Melbourne, Victoria, Australia: Thomson.
- Francis, G. (2007). *Introduction to SPSS for Windows: v. 15.0 and 14.0 with Notes for Studentware* (5th ed.). Sydney: Pearson Education.
- Howell, D. C. (2010). *Statistical methods for psychology* (7th ed.). Belmont, CA: Wadsworth.
- Howitt, D. & Cramer, D. (2011). *Introduction to statistics in psychology* (5th ed.). Harlow, UK: Pearson.

59

### Next lecture

- Statistical power
- Significance testing
- Inferential decision making
- Effect sizes
- Confidence intervals
- Publication bias
- Academic integrity

60

## Open Office Impress

- This presentation was made using Open Office Impress.
- Free and open source software.
- <http://www.openoffice.org/product/impress.html>

