
Summary & Conclusion

Survey Research & Design in Psychology James Neill, 2016 Creative Commons Attribution 4.0

Overview 1. Survey research and design 1. Survey research 2. Survey design 2. Univariate and bivariate 1. Descriptives & graphing 2. Correlation 3. Psychometrics 1. Éxploratory factor analysis 2. Psychometric instrument development 4. Multiple linear regression 1. MLR I 2. MLR II 5. Power & summary 1. Power & effect size 2. Summary and conclusion 2



5

Types of research

- Surveys are used in all types of research:
 - •Experimental
 - Quasi-experimental
 - Non-experimental

What is a survey?

- What is a survey?
 - A standardised stimulus for converting fuzzy psychological phenomenon into hard data.
- History
 - •Survey research has developed into a popular social science research method since the 1920s.

Purposes of research

Purposes of research:

- Information gathering
 - Exploratory
 - Descriptive
- Theory testing & building
 - Explanatory
 - Predictive

Survey research

Pros include:

- · Ecological validity
- Cost efficiency
- · Can obtain lots of data

Cons include:

· Low compliance

Self-administered

anonymity

needs

non-response


- demand characteristics

- access to representative sample

Pros: - cost

Cons:

· Reliance on self-report



Survey questions

- · Objective versus subjective questions:
 - **Objective** there is a verifiably true answer
 - Subjective based on perspective of respondent
- Open versus closed questions:
 - Open empty space for answer
 - Closed pre-set response format options

11

7

- adjustment to cultural differences, special

Survey types

- Level of measurement 1. Categorical/Nominal
- 1. Arbitrary numerical labels
 - 2. Could be in any order

2. Ordinal

- 1. Ordered numerical labels
- 2. Intervals may not be equal
- 3. Interval
 - 1. Ordered numerical labels 2. Equal intervals

4. Ratio

- 1. Data are continuous
- 2. Meaningful 0

12

10

8

Opposite for interviewadministered

surveys

Response formats

- 1. Dichotomous and Multichotomous
- 2. Multiple response
- 3. Verbal frequency scale (Never ... Often)
- 4. Ranking (in order \rightarrow Ordinal)
- 5. Likert scale (equal distances \rightarrow Interval, typically with 3 to 9 options)
- 6. Graphical rating scale (e.g., line)
- 7. Semantic differential (opposing words)
- 8. Non-verbal (idiographic)


Sampling

- 1. Key terms
 - 1. (Target) population
 - 2. Sampling frame
 - 3. Sample
- 2. Sampling
 - 1. Probability 2. Non-probability
 - 1. Simple (random) 1. Convenience
 - 2. Systematic
 - Stratified

14

2. Purposive

3. Snowball





13


Descriptive statistics


- Level of measurement and normality determines how data can be described.
- What is the central tendency?
 Frequencies, Percentages (Non-para)
 Mode, Median, Mean (Para)
- What is the variability?
 - Min, Max, Range, Quartiles (Non-para)
 Standard Deviation, Variance (Para)

-Standard Deviation, variance (Fara)









Covariation

- 1. The world is made of covariations.
- 2. Covariations are the building blocks of more complex relationships which can be analysed through the use of:
 - 1. factor analysis
 - 2. reliability analysis
 - 3. multiple regression

23

Purpose of correlation

The underlying purpose of correlation is to help address the question:

What is the

- relationship or
- association or
- shared variance or

• co-relation

between two variables?

What is correlation?

- 1. Standardised covariance
- 2. Ranges between -1 and +1, with more extreme values indicating stronger relationships
- 3. Correlation does not prove causation – may be opposite causality, bi-directional, or due to other variables.

Types of correlation

- Nominal by nominal: Phi (Φ) / Cramer's V, Chi-squared
- Ordinal by ordinal: Spearman's rank / Kendall's Tau b
- Dichotomous by interval/ratio: Point bi-serial *r*_{pb}
- Interval/ratio by interval/ratio: Product-moment or Pearson's r

26

Correlation steps

- 1. Choose measure of correlation and graphs based on levels of measurement.
- 2. Check graphs (e.g., scatterplot):
 - -Linear or non-linear?
 - -Outliers?
 - -Homoscedasticity?
 - -Range restriction?
 - -Sub-samples to consider?

27

25

Interpreting correlation

- Coefficient of determination
 - Correlation squared
 - -Indicates % of shared variance

<u>Strength</u>	<u>r</u>	<u>r</u> ²
Weak:	.13	1 – 10%
Moderate:	.35	10 - 25%
Strong:	> .5	> 25%

29

Correlation steps

3. Consider

- -Effect size (e.g., Φ , Cramer's V, r, r²)
- -Direction
- -Inferential test (p)
- 4. Interpret/Discuss
 - -Relate back to hypothesis
 - -Size, direction, significance
 - -Limitations e.g.,
 - Heterogeneity (sub-samples)
 - Range restriction
 - Causality?

28



- 1. Levels of measurement
- 2. Normality
- 3. Linearity
 - Effects of outliers
 Non-linearity
- 4. Homoscedasticity
- 5. No range restriction
- 6. Homogenous samples
- 7. Correlation is not causation

Exploratory factor analysis (Lecture 5)

What is factor analysis?

- Factor analysis is a family of multivariate correlational data analysis methods for summarising clusters of covariance.
- FA summarises correlations amongst items into a smaller number of underlying fuzzy constructs (called factors).

32

FAQ

Assumptions

- Sample size
 - 5+ cases per variables
 (ideally 20+ cases per variable)

-N > 200

- Bivariate & multivariate outliers
- Factorability of correlation matrix (Measures of Sampling Adequacy)
- Normality enhances the solution

33

31

Steps / process

- 1. Test assumptions
- 2. Select type of analysis
- 3. Determine no. of factors (Eigen Values, Scree plot, % variance explained)
- 4. Select items (check factor loadings to identify which items belong in which factor; drop items one by one; repeat)
- 5. Name and define factors
- 6. Examine correlations amongst factors
- 7. Analyse internal reliability | Lecture
- 8. Compute composite scores 6 34

Types of FA

- **PAF** (Principal Axis Factoring): Best for theoretical data exploration –uses shared variance
- PC (Principal Components): Best for data reduction

 uses all variance

Rotation

- Orthogonal (Varimax)

 perpendicular (uncorrelated) factors
- Oblique (Oblimin)
 - angled (correlated) factors
- Consider trying both ways
 Are solutions different? Why?

Factor extraction


How many factors to extract?

- Inspect EVs

 look for > 1 or sudden drop (inspect scree plot)
- % of variance explained - aim for 50 to 75%
- Interpretability

 does each factor 'make sense'?
- Theory
 - does the model fit with theory?

Item selection



- a simple factor structure (each variable loads strongly (> +.50) on only one factor)
- each factor has multiple loading variables (more loadings → greater reliability)
- target factor loadings are high (> .6) and cross-loadings are low (< .3), with few intermediate values (.3 to .6).

38

FAQ

Psychometrics instrument development (Lecture 6)

39

FAQ

37

Concepts & their measurement

- 1. Concepts name common elements
- 2. Hypotheses identify relations between concepts
- 3. Brainstorm indicators of a concept
- 4. Define the concept
- 5. Draft measurement items
- 6. Pre-test and pilot test
- 7. Examine psychometric properties
- 8. Redraft/refine and re-test

41

Psychometrics

- 1. Science of psychological measurement
- 2. Goal: Validly measure individual psychosocial differences
- 3. Develop and test psychological measures e.g., using
 - 1. Factor analysis
 - 2. Reliability and validity

40

Measurement error

- 1. Deviation of measure from true score
- 2. Sources:
 - 1. Non-sampling (e.g., paradigm, respondent bias, researcher bias)
 - 2. Sampling (e.g., non-representativeness)
- 3. How to minimise:
 - 1. Well-designed measures
 - 2. Reduce demand effects
 - 3. Representative sampling
 - 4. Maximise response rate
 - 5. Ensure administrative accuracy

Reliability

- 1. Consistency or reproducibility
- 2. Types
 - 1. Internal consistency
 - 2. Test-retest reliability
- 3. Rule of thumb
 - 1. > .6 OK
 - 2. > .8 Very good
- 4. Internal consistency
 - 1. Split-half
 - 2. Odd-even
 - 3. Cronbach's alpha

Validity

- 1. Extent to which a measure measures what it is intended to measure
- 2. Multifaceted
 - 1. Correlations with similar measures
 - 2. Performance in relation to other variables
 - 3. Predicts future

44

46

Composite scores

Ways of creating composite (factor) scores:

1. Unit weighting

1.Total of items or

2. Average of items (recommended for lab report)

2. Regression weighting

1. Each item is weighted by its importance to measuring the underlying factor (based on regression weights) 45

Writing up instrument development

1. Introduction

- Review constructs & previous structures
 Generate research question
- 2. Method
 - 1. Explain measures and their development
- 3. Results
 - 1. Factor analysis
 - 2. Reliability of factors
 - Descriptive statistics for composite scores
 Correlations between factors
- 4. Discussion
 - 1. Theory? / Measure? / Recommendations?

Multiple linear regression (Lectures 7 & 8)

Linear regression

- 1. Best-fitting straight line for a scatterplot of two variables
- 2. Y = bX + a + e 1. Predictor (X; IV) 2. Outcome (Y; DV)
- 3. Least squares criterion
- 4. Residuals are the vertical distance between actual and predicted values

48

47

Level of measurement and dummy coding

- 1. Levels of measurement
 - 1. DV = Continuous (Likert or ratio + normal)
 - 2. IV = Continuous or dichotomous
- 2. Dummy coding
 - 1. Convert complex variable into series of dichotomous IVs

49

Multiple linear regression

1. Multiple IVs to predict a single DV:

 $Y = b_1 x_1 + b_2 x_2 + \dots + b_i x_i + a + e$

- 2. Overall fit: R, R^2 , and Adjusted R^2
- 3. Coefficients
 - Relation between each IV and the DV, adjusted for the other IVs
 B, β, t, p, and sr2
- 4. Types
 - 1. Standard
 - 2. Hierarchical
 - 3. Stepwise / Forward / Backward

50

General steps

- 1. Develop model and hypotheses
- 2. Check assumptions
- 3. Choose type
- 4. Interpret output
- 5. Develop a regression equation (if needed)

51

Summary: Semi-partial correlation (*sr*)

- 1. In MLR, *sr* is labelled "part" in the regression coefficients table SPSS output
- 2. *sr*² is the unique % of the DV variance explained by each IV

52

Residual analysis

- 1. Residuals are the difference between predicted and observed Y values
- 2. MLR assumption is that residuals are normally distributed.
- 3. Examining residuals also helps assess:
 - 1. Normality
 - 2. Linearity
 - 3. Homoscedasticity

53

Interactions

- 1. In MLR, IVs may interact to:
 - 1. Increase effect on DV
 - 2. Decrease effect on DV
- 2. Model interactions with hierarchical MLR:
 - 1. Step 1: Enter IVs
 - 2. Step 2: Enter cross-product of IVs
 - 3. Examine change in R^2

Analysis of change

Variance in changes over time can be assessed by:

1. Standard regression 1. Calculate difference scores (Time 2 minus Time 1) and use as DV

2. Hierarchical MLR

1. Step 1: "Partial out" baseline scores

(Lecture 9)

2. Step 2: Enter other IVs to help predict variance in changes over time.


55


Writing up an MLR

1. Introduction:

- 1. Consider theoretical relationship between possible predictors and the outcome variable of interest
- 2. One hypothesis per predictor
- 2. Results:
 - 1. Univariate descriptive statistics
 - 2. Correlations
 - 3. Type of MLR and assumptions
 - 4. Overall model 5. Regression coefficients

56







- 1. Power = probability of detecting a real effect as statistically significant
- 2. Increase by:
 - -↑ N
 - $-\uparrow$ critical α
- -↑ES Power
- >.8 "desirable"
- ~.6 is more typical
- · Can be calculated prospectively and

retrospectively

60

Effect size

- 1. Standardised size of difference or strength of relationship
- 2. Inferential tests should be accompanied by ESs and CIs
- Common bivariate ESs include:
 Cohen's d
 Correlation r
- Cohen's *d* not in SPSS use an online effect size calculator

Confidence interval

- 1. Gives 'range of certainty'
- 2. Can be used for *B*, *M*, ES etc.
- 3. Can be examined1. Statistically (upper and lower limits)2. Graphically (e.g., error-bar graphs)

62


Publication bias

- 1. Tendency for statistically significant studies to be published over non-significant studies
- 2. Indicated by gap in funnel plot \rightarrow file-drawer effect
- 3. Counteracting biases in scientific publishing:
 - low-power studies tend to underestimate real effects
 - bias towards publish sig. effects over nonsig. effects 63

Academic integrity

- 1. Violations of academic integrity are evident and prevalent amongst those with incentives to do so:
 - 1. Students
 - 2. Researchers
 - 3. Commercial sponsors
- 2. Adopt a balanced, critical approach, striving for objectivity and academic integrity

64



Learning outcomes

- 1. Design and conduct survey-based research in psychology;
- 2. Use SPSS to conduct and interpret data analysis using correlation-based statistics, including reliability, factor analysis and multiple regression analysis;
- 3. Communicate in writing the results of survey-based psychological research

65

Graduate attributes

- 1. Display initiative and drive, and use organisation skills to plan and manage workload
- 2. Employ up-to-date and relevant knowledge and skills
- 3. Take pride in professional and personal integrity
- 4. Use creativity, critical thinking, analysis and research skills to solve theoretical and real-world problems 67

Feedback

• Please provide feedback about what worked well and what could be improved via the Unit Satisfaction Survey

Open Office Impress

- This presentation was made using Open Office Impress.
- Free and open source software.
- <u>http://www.openoffice.org/product/impress.html</u>

