

# Z Transform (H.1)

## Definition

20170208

Copyright (c) 2016 - 2017 Young W. Lim.

Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.2 or any later version published by the Free Software Foundation; with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts. A copy of the license is included in the section entitled "GNU Free Documentation License".

Based on  
Complex Analysis for Mathematics and Engineering  
J. Mathews

# z - Transform

$$X(z) = \sum_{k=-\infty}^{+\infty} x[k] z^{-k}$$

$$z = |r| e^{j2\pi F} \\ = |r| e^{j\Omega}$$

$$x[n] \longleftrightarrow X(z)$$

One Sided z-transform

$$X(z) = \sum_{k=0}^{+\infty} x[k] z^{-k}$$

# Inverse z-Transform

$$X(z) = \mathcal{Z}[\{x_n\}_{n=0}^{\infty}]$$

$$= \sum_{n=0}^{\infty} x_n z^{-n}$$

$$= \sum_{n=0}^{\infty} x[n] z^{-n}$$

$$x[n] \longrightarrow X(z)$$

$$x_n = x[n]$$

$$= \mathcal{Z}^{-1}[X(z)]$$

$$= \frac{1}{2\pi i} \int_{\mathcal{C}} X(z) z^{n+1} dz$$

$$x[n] \longleftarrow X(z)$$

# Admissible Form of $z$ -transform

$$X(z) = \sum_{n=0}^{\infty} x[n] z^{-n}$$

$X(z)$ : admissible  $z$ -transform

if  $X(z)$  is a rational function

$$X(z) = \frac{P(z)}{Q(z)} = \frac{b_0 + b_1 z^1 + b_2 z^2 + \dots + b_{p-1} z^{p-1} + b_p z^p}{a_0 + a_1 z^1 + a_2 z^2 + \dots + a_{q-1} z^{q-1} + a_q z^q}$$

$P(z)$ : a polynomial of degree  $p$

$Q(z)$ : a polynomial of degree  $q$

# Residue Theorem

$D$ : Simply connected domain


$C$ : Simple closed contour (CCW) in  $D$

if  $f(z)$  is **analytic** inside  $C$  and on  $C$   
except at the points  $z_1, z_2, \dots, z_k$  in  $C$

then

$$\frac{1}{2\pi i} \int_C f(z) dz = \sum_{j=1}^k \text{Res}(f(z), z_j)$$


Singular points of  $f(z)$ :  $z_1, z_2, \dots, z_k$



# Integration of a function of a complex var.


$$\oint_C f(z) dz = 2\pi i \sum_{k=1}^n \text{Res}(f(z), z_k)$$

finite number  $k$  of  
singular points  $z_k$   
residue theorem



$$\oint_C f(z) dz = 0 \quad \text{if } f(z) \text{ is analytic within and on } C$$

no singularity



$$\oint_C f(z) dz = 0 \quad \text{if } f(z) = F'(z) \text{ on } C$$

:  $F(z)$  is an antiderivative of  $f(z)$   
fundamental theorem of calculus

$\oint_C f(z) dz = 0$  if  $f(z)$  is continuous in  $D$  and  
 $f(z) = F'(z)$  :  $F(z)$  is an antiderivative of  $f(z)$   
fundamental theorem of calculus



# Series Expansion

can expand  $f(z)$  about any point  $z_m$   
over powers of  $(z - z_m)$

whether or not  $f(z)$  is singular at  $z_m$   
or at other points between  $z$  and  $z_m$

$$f(z) = \sum_{n=\nu_1}^{\infty} a_n^{(m)} (z - z_m)^n$$

① Laurent Series Expansion of  $f(z)$  at  $z_m$   
general  $\nu_1$  - depend on  $f(z)$  and  $z_m$

②  $z$ -transform of  $a_n^{(m)}$   
general  $\nu_1$  - depend on  $f(z)$

$$z_m = 0$$

③ Taylor Series Expansion of  $f(z)$  at  $z_m$   
positive  $\nu_1$  - depend on  $f(z)$  and  $z_m$  ( $\nu_1 > 0$ )

④ MacLaurin Series Expansion of  $f(z)$  at  $z_m$   
positive  $\nu_1$  - depend on  $f(z)$  ( $\nu_1 > 0$ )

$$z_m = 0$$

$$f(z) = \sum_{n=n_1}^{\infty} a_n^{(m)} (z - z_m)^n$$

$n_1 > 0$  pos powers

$$z_m = 0$$

① Laurent Series	③ Taylor Series
② z-transform	④ MacLaurin Series

\* Expansion of  $f(z)$  about any point  $z_m$   
over powers of  $(z - z_m)$

$$f(z) = \sum_{n=n_1}^{\infty} a_n^{(m)} (z - z_m)^n$$

$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz$$

for general  $f(z)$

$$a_n^{(m)} = \sum_k \operatorname{Res} \left( \frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

for general  $f(z)$

$$a_n^{(m)} = \frac{1}{n!} f^{(n)}(z_m) \quad n_1 \geq 0$$

for analytic  $f(z)$  within  $C$

analytic  $f(z) \longrightarrow \frac{f(z)}{(z - z_m)^{n+1}}$  has a pole at  $z_m$   
order of  $n+1$

$$f(z) = \sum_{n=n_1}^{\infty} a_n^{(m)} (z - z_m)^n$$

$z_m$ : possible poles of  $f(z)$   
not necessarily poles

$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z')}{(z' - z_m)^{n+1}} dz'$$
$$= \sum_k \operatorname{Res} \left( \frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$




$z_k$ : poles of  $\frac{f(z)}{(z - z_m)^{n+1}}$   
within  $C$

$$= \frac{1}{n!} f^{(n)}(z_m) \quad n_1 \geq 0$$


# Residue Theorem and Laurent Series

assumed there are  $(m)$  singularities (poles) of  $f(z)$  in a region


let  $C_m$  is taken to enclose only one pole  $z_m$



$a_n^{(1)}$  expanded at  $z_1$   
 $C_1$  encloses  $z_1$  only  
 $\tilde{a}_{-1}^{(1)} = \text{Res}(f(z), z_1)$




$a_n^{(2)}$  expanded at  $z_2$   
 $C_2$  encloses  $z_2$  only  
 $\tilde{a}_{-1}^{(2)} = \text{Res}(f(z), z_2)$




$a_n^{(3)}$  expanded at  $z_3$   
 $C_3$  encloses  $z_3$  only  
 $\tilde{a}_{-1}^{(3)} = \text{Res}(f(z), z_3)$

# Series Expansion at $z_m$



$$f(z) = \sum_{n=\eta_1}^{\infty} a_n^{(m)} (z - z_m)^n$$



$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz$$

$$= \sum_k \text{Res} \left( \frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

Let  $z_1, z_2, z_3$  poles of  $f(z)$

Then the poles of  $\frac{f(z)}{(z - z_m)^{n+1}}$

$n \geq 0$	$z_1, z_2, z_3, z_m$
$n < 0$	$z_1, z_2, z_3$

$$f(z) = \sum_{n=\eta_1}^{\infty} a_n^{[m]} (z - z_m)^n \quad \boxed{n \leftarrow k}$$

$$f(z) = \sum_{k=\eta_1}^{\infty} a_k^{[m]} (z - z_m)^k$$

for a given  $n$

$$\frac{f(z)}{(z - z_m)^{n_H}} = \sum_{k=\eta_1}^{\infty} a_k^{[m]} (z - z_m)^{k-n-1} \quad \begin{array}{l} k: \text{index variable} \\ n: \text{fixed value} \end{array}$$

$$\oint_C \frac{f(z)}{(z - z_m)^{n_H}} dz = \oint_C \sum_{k=\eta_1}^{\infty} a_k^{[m]} (z - z_m)^{k-n-1} dz$$

$$= \sum_{k=\eta_1}^{\infty} \oint_C a_k^{[m]} (z - z_m)^{k-n-1} dz$$

$$\boxed{k=n}$$


$$\oint_C \frac{f(z)}{(z - z_m)^{n_H}} dz = \oint_C a_n^{[m]} \frac{1}{(z - z_m)} dz = 2\pi i \cdot a_n^{[m]}$$

$$\boxed{a_n^{[m]} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{n_H}} dz}$$

$$\oint_C \left[ \dots (z - z_m)^{-3} + (z - z_m)^{-2} + \frac{1}{(z - z_m)} + 1 + (z - z_m) + (z - z_m)^2 + \dots \right] dz$$

$$= \oint_C \frac{1}{(z - z_m)} dz = 2\pi i$$

# Series Expansion at $z=0$



$$f(z) = \sum_{n=n_1}^{\infty} a_n^{(m)} z^n$$

$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{z^{n+1}} dz$$
$$= \sum_k \operatorname{Res} \left( \frac{f(z)}{z^{n+1}}, z_k \right)$$

Poles  $z_k$

$$n \geq 0 \quad z_1, z_2, z_3, \circ$$

$$n < 0 \quad z_1, z_2, z_3$$



# Residue & Series Expansion at $z_m$

expansion at  $z_m$

$$\eta = -1$$

$$\eta + 1 = 0 \quad (z - z_m)^{\eta+1} = 1$$

$$a_n^{\{m\}} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{\eta+1}} dz$$


$$= \sum_k \text{Res} \left( \frac{f(z)}{(z - z_m)^{\eta+1}}, z_k \right)$$

$$a_{-1}^{\{m\}} = \frac{1}{2\pi i} \oint_C f(z) dz$$

$$= \sum_k \text{Res} (f(z), z_k)$$

$$a_{-1}^{\{m\}} = \frac{1}{2\pi i} \oint_C f(z) dz = \sum_k \text{Res} (f(z), z_k)$$

$$a_{-1}^{\{m\}} = \text{Res} (f(z), z_m)^\vee = \sum_k \text{Res} (f(z), z_k)$$



$$f(z) = \sum_{n=\eta_1}^{\infty} a_n^{\{m\}} (z - z_m)^n$$

$$a_n^{\{m\}} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{\eta+1}} dz$$

$$= \sum_k \text{Res} \left( \frac{f(z)}{(z - z_m)^{\eta+1}}, z_k \right)$$

Residue  $\rightarrow$  Laurent Series  $\rightarrow$  annulus region

if  $C$  encloses only one pole  $z_0$ ,  
and the expansion at that pole  $z_0$  is assumed,  
then

$$a_{-1}^{\{0\}} = \frac{1}{2\pi i} \oint_{C_0} f(z) dz = \text{Res}(f(z), z_0)$$

Let  $\tilde{a}_{-1}^{\{m\}} = \text{Res}(f(z), z_m)$  notation  $\sim$

the residue of  $f(z)$  at  $z_m$


Using  $C_m$  which is in the annulus ROC

$$f(z) = \sum_{n=-\infty}^{+\infty} a_n^{\{m\}} (z - z_m)^n$$



$$\begin{aligned}\tilde{a}_{-1}^{\{1\}} &= \text{Res}(f(z), z_1) \\ &= \frac{1}{2\pi i} \oint_{C_1} f(z) dz\end{aligned}$$

$$f(z) = \sum_{n=-\infty}^{+\infty} a_n^{\{1\}} (z-z_1)^n$$




$$\begin{aligned}\tilde{a}_{-1}^{\{2\}} &= \text{Res}(f(z), z_2) \\ &= \frac{1}{2\pi i} \oint_{C_2} f(z) dz\end{aligned}$$

$$f(z) = \sum_{n=-\infty}^{+\infty} a_n^{\{2\}} (z-z_2)^n$$



$$\begin{aligned}\tilde{a}_{-1}^{\{3\}} &= \text{Res}(f(z), z_3) \\ &= \frac{1}{2\pi i} \oint_{C_3} f(z) dz\end{aligned}$$


$$f(z) = \sum_{n=-\infty}^{+\infty} a_n^{\{3\}} (z-z_3)^n$$



$$f(z) = \sum_{n=n_1}^{\infty} a_n^{(m)} (z - z_m)^n$$

$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_m)^{n+1}} dz$$

$$= \sum_k \operatorname{Res} \left( \frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$




$$a_{-1}^{(m)} = \frac{1}{2\pi i} \oint_C f(z) dz$$


$$= \sum_k \operatorname{Res} (f(z), z_k)$$



$$\tilde{a}_{-1}^{(1)} = \operatorname{Res}(f(z), z_1)$$



$$\tilde{a}_{-1}^{(2)} = \operatorname{Res}(f(z), z_2)$$





$$\tilde{a}_{-1}^{(3)} = \operatorname{Res}(f(z), z_3)$$

$$a_{-1}^{(m)} = \tilde{a}_{-1}^{(1)} + \tilde{a}_{-1}^{(2)} + \tilde{a}_{-1}^{(3)}$$

$$a_{-1}^{(m)} = \operatorname{Res}(f(z), z_1)$$

$$+ \operatorname{Res}(f(z), z_2)$$

$$+ \operatorname{Res}(f(z), z_3)$$



$$\oint_C f(z) dz = 2\pi j \sum_{k=1}^M \tilde{a}_{-1}^{(k)} = 2\pi j \sum_{k=1}^M \text{Res}(f(z), z_k)$$

residue theorem

$$a_n = \sum_{k=1}^M \text{Res} \left( \frac{f(z)}{(z-z_k)^{n+1}}, z_k \right)$$

Laurent coefficient

$C$  encloses  $k$  poles


$C_k$  encloses only the  $k$ -th pole

$\tilde{a}_{-1}^{(k)}$  the residue of the  $k$ -th pole enclosed by  $C$ ,  $z_k$

$$f(z) = \sum_{n=n_1}^{\infty} a_n (z - z_m)^n$$

$$a_n = \frac{1}{2\pi i} \oint_C \frac{f(z')}{(z' - z_m)^{n+1}} dz'$$

$$= \sum_k \text{Res} \left( \frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$



$C$  is in the same region of analyticity of  $f(z)$   
typically a circle centered on  $z_m$

$z_k$  within  $C$  : singularities of  $\frac{f(z)}{(z - z_m)^{n+1}}$

$n_1 = n_{f,m}$  depends on  $f(z)$ ,  $z_m$

$a_n$  depends on  $f(z)$ ,  $z_m$ , region of analyticity

Whether  $f(z)$  is singular at  $z = z_m$  or not  
or at other points between  $z$  and  $z_m$

We can expand  $f(z)$  about any point  $z_m$   
over powers of  $(z - z_m)$ .




$$f(z) = \sum_{n=n_1}^{\infty} a_n (z - z_m)^n$$

$$a_n = \frac{1}{2\pi i} \oint_C \frac{f(z')}{(z' - z_m)^{n+1}} dz'$$


$$= \sum_k \operatorname{Res} \left( \frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

$z_k$  within  $C$  : singularities of  $\frac{f(z)}{(z - z_m)^{n+1}}$

$$\begin{cases} \text{poles of } f(z) \cup z = z_m & n \geq 0 \\ \text{poles of } f(z) & n < 0 \end{cases}$$







$\text{Res}(f, z_1) + \text{Res}(f, z_2) + \text{Res}(f, z_m)$



$\text{Res}(f, z_1) + \text{Res}(f, z_2) + \text{Res}(f, z_3) + \text{Res}(f, z_m)$








$$f(z) = \sum_{n=-\infty}^{\infty} a_n^{(m)} (z - z_m)^n$$

$$a_n^{(m)} = \frac{1}{2\pi i} \oint_C \frac{f(z')}{(z' - z_m)^{n+1}} dz'$$

$$= \sum_k \operatorname{Res} \left( \frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$



## Laurent's Theorem

$f$ : analytic within the annular domain  $D$

$$r < |z - z_0| < R$$

then

$$f(z) = \sum_{k=-\infty}^{+\infty} a_k (z - z_0)^k, \text{ valid for } r < |z - z_0| < R$$


$$a_k = \frac{1}{2\pi i} \oint_C \frac{f(s)}{(s - z_0)^{k+1}} ds, \quad k = 0, \pm 1, \pm 2, \dots$$

$C$ : a simple closed curve  
that lies entirely within  $D$   
that encloses  $z_0$

$$a_{-1} = \frac{1}{2\pi i} \oint_C f(s) ds \quad \rightarrow \quad \oint_C f(s) ds = 2\pi i \cdot a_{-1}$$

$$a_{-1} = \frac{1}{2\pi i} \oint_C f(s) ds = \text{Res}(f(z), z_0)$$

$$= \begin{cases} \lim_{z \rightarrow z_0} (z - z_0) f(z) & \text{(simple)} \\ \frac{1}{(n-1)!} \lim_{z \rightarrow z_0} \frac{d^{n-1}}{dz^{n-1}} (z - z_0)^n f(z) & \text{(order } n) \end{cases}$$



$z_0$ : expansion point

$z$ : evaluation point

which poles of  $f(z)$  lie between the point of evaluation  $z$  and the point  $z_0$  about which the expansion is formed

$\frac{f(z')}{(z' - z_0)}$  is analytic between  $C_1$  &  $C_2$

deformation theorem

$C_1 - C_2$  coincide

common contour  $\curvearrowright$

## Cauchy's Residue Theorem



$f(z)$ : **analytic** on and within  $C$   
except a finite number of **singular points**  
 $z_1, z_2, \dots, z_n$  within  $C$

then

$$\int_C f(z) dz = 2\pi i \sum_{k=1}^n \text{Res}(f(z), z_k)$$

$D$ : a simply connected domain

$C$ : a simple closed contour in  $D$



$z_1$

$$f(z) = \sum_{k=-\infty}^{+\infty} a_k (z - z_1)^k$$

$$a_{-1}^{f(z)} = \frac{1}{2\pi i} \oint_{C_1} f(s) ds = \text{Res}(f(z), z_1)$$

$z_2$

$$f(z) = \sum_{k=-\infty}^{+\infty} a_k (z - z_2)^k$$


$$a_{-1}^{f(z)} = \frac{1}{2\pi i} \oint_{C_2} f(s) ds = \text{Res}(f(z), z_2)$$

$z_3$

$$f(z) = \sum_{k=-\infty}^{+\infty} a_k (z - z_3)^k$$

$$a_{-1}^{f(z)} = \frac{1}{2\pi i} \oint_{C_3} f(s) ds = \text{Res}(f(z), z_3)$$

$z_1$




Laurent series expansion at  $z_1$

$$f(z) = \sum_{k=-\infty}^{+\infty} a_k (z - z_1)^k$$

$$a_{-1} = \frac{1}{2\pi i} \oint_{C_1} f(s) ds = \text{Res}(f(z), z_1)$$

$z_2$




Laurent series expansion at  $z_2$

$$f(z) = \sum_{k=-\infty}^{+\infty} a_k (z - z_2)^k$$

$$a_{-1} = \frac{1}{2\pi i} \oint_{C_2} f(s) ds = \text{Res}(f(z), z_2)$$


$z_3$




Laurent series expansion at  $z_3$

$$f(z) = \sum_{k=-\infty}^{+\infty} a_k (z - z_3)^k$$


$$a_{-1} = \frac{1}{2\pi i} \oint_{C_3} f(s) ds = \text{Res}(f(z), z_3)$$




$$\int_C f(z) dz = 2\pi i \sum_{k=1}^n \text{Res}(f(z), z_k)$$




$$\int_C f(z) dz = 2\pi i \operatorname{Res}(f(z), z_1) + 2\pi i \operatorname{Res}(f(z), z_2) + 2\pi i \operatorname{Res}(f(z), z_3)$$



$$\int_{C'} f(z) dz = 2\pi i \operatorname{Res}(f(z), z_1) + 2\pi i \operatorname{Res}(f(z), z_2)$$



$$\int_{C''} f(z) dz = 2\pi i \operatorname{Res}(f(z), z_1)$$



$$\int_{C'''} f(z) dz = 0$$

# Inverse z-Transform $x[n] = \frac{1}{2\pi i} \int_C X(z) z^{n-1} dz$

$$X(z) = \sum_{k=0}^{\infty} x_k z^{-k}$$

$$z^{n-1} X(z) = \left( \sum_{k=0}^{\infty} x_k z^{-k} \right) z^{n-1}$$

$$\int z^{n-1} \text{LHS} dz = \int \text{RHS} z^{n-1} dz$$

$$= \sum_{k=0}^{\infty} x_k z^{-k+n-1}$$

$$[0, \infty) = [0, n-1] \cup [n] \cup [n+1, \infty)$$

$$= \sum_{k=0}^{n-1} x_k z^{-k+n-1} + \sum_{k=n}^n x_k z^{-k+n-1} + \sum_{k=n+1}^{\infty} x_k z^{-k+n-1}$$

$$= \sum_{k=0}^{n-1} x_k z^{-k+n-1} + \frac{x_n}{z^1} + \sum_{k=n+1}^{\infty} \frac{x_k}{z^{k-n+1}}$$

$$\int_C X(z) z^{n-1} dz = \int_C \sum_{k=0}^{n-1} x_k z^{-k+n-1} dz + \int_C \frac{x_n}{z^1} dz + \int_C \sum_{k=n+1}^{\infty} \frac{x_k}{z^{k-n+1}} dz$$

$$= \sum_{k=0}^{n-1} x_k \int_C z^{-k+n-1} dz + x_n \int_C \frac{1}{z^1} dz + \sum_{k=n+1}^{\infty} x_k \int_C \frac{1}{z^{k-n+1}} dz$$

$$= \sum_{k=0}^{n-1} x_k \cdot 0 + x_n \cdot 2\pi i + \sum_{k=n+1}^{\infty} x_k \cdot 0$$

$$x[n] = \frac{1}{2\pi i} \int_C X(z) z^{n-1} dz$$



## Z-transform

$$z_m = 0$$

$$\begin{aligned} x[n] &= \frac{1}{2\pi i} \oint_C f(z) z^{n-1} dz \\ &= \sum_k \operatorname{Res}(f(z) z^{n-1}, z_k) \end{aligned}$$

$n > 0$   $z_k$ : poles of  $f(z)$

$n = 0$   $z_k$ : poles of  $f(z)$  +  $z = 0$   
 $z^{n-1} = z^{-1} = \frac{1}{z}$

$x[n]$  includes  $u[n] \rightarrow X[z]$  contains  $z$  on its numerator

Also, think about modified partial fraction  $\frac{X[z]}{z}$

## Laurent Expansion

expansion at  $z_m$

$$\begin{aligned} a_n^{\{m\}} &= \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z-z_m)^{n+1}} dz \\ &= \sum_k \operatorname{Res}\left(\frac{f(z)}{(z-z_m)^{n+1}}, z_k\right) \end{aligned}$$





$$z_m = 0$$

$$\begin{aligned} a_n^{\{0\}} &= \frac{1}{2\pi i} \oint_C \frac{f(z)}{z^{n+1}} dz \\ &= \sum_k \operatorname{Res}\left(\frac{f(z)}{z^{n+1}}, z_k\right) \end{aligned}$$


$$\begin{aligned} a_{-n}^{\{0\}} &= \frac{1}{2\pi i} \oint_C f(z) z^{n-1} dz \\ &= \sum_k \operatorname{Res}(f(z) z^{n-1}, z_k) \end{aligned}$$

$$\begin{aligned} a_{-n}^{\{0\}} &= \frac{1}{2\pi i} \oint_C \frac{f(z)}{z^{-n+1}} dz \\ &= \sum_k \operatorname{Res}\left(\frac{f(z)}{z^{-n+1}}, z_k\right) \end{aligned}$$

# Different D, Different Laurent Series



$$\begin{aligned} x[n] &= \frac{1}{2\pi i} \int_C X(z) z^{n-1} dz \\ &= \sum_{z_k} \text{Res}(X(z) z^{n-1}, z_k) \end{aligned}$$



z-transform

$$f(z) = \frac{-1}{(z-1)(z-2)}$$

Complex Variables and Ap  
Brown & Churchill

$$f(z) = \frac{-1}{(z-1)(z-2)} = \frac{1}{z-1} - \frac{1}{z-2}$$

$$D_1: |z| < 1$$

$$D_2: 1 < |z| < 2$$

$$D_3: 2 < |z|$$

$$\textcircled{1} D_1 \quad |z| < 1, \quad \left|\frac{z}{2}\right| < 1$$

$$\begin{aligned} f(z) &= \frac{1}{z-1} - \frac{1}{z-2} = \frac{-1}{1-z} + \frac{1}{1-\left(\frac{z}{2}\right)} \\ &= -\sum_{n=0}^{\infty} z^n + \sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} = \sum_{n=0}^{\infty} (2^{-n-1} - 1)z^n \quad |z| < 1 \end{aligned}$$

$$\textcircled{2} D_2 \quad 1 < |z| < 2 \Rightarrow \left|\frac{1}{z}\right| < 1, \quad \left|\frac{z}{2}\right| < 1$$


$$\begin{aligned} f(z) &= \frac{1}{z-1} - \frac{1}{z-2} = \frac{1}{z} \cdot \frac{1}{1-\left(\frac{1}{z}\right)} + \frac{1}{2} \frac{1}{1-\left(\frac{z}{2}\right)} \\ &= \sum_{n=0}^{\infty} \frac{1}{z^{n+1}} + \sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} \\ &= \sum_{n=1}^{\infty} \frac{1}{z^n} + \sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} \end{aligned}$$

$$\textcircled{3} D_3 \quad 2 < |z| \quad \left|\frac{2}{z}\right| < 1 \quad \left|\frac{1}{z}\right| < 1$$

$$\begin{aligned} f(z) &= \frac{1}{z-1} - \frac{1}{z-2} = \frac{1}{z} \frac{1}{1-\left(\frac{1}{z}\right)} - \frac{1}{z} \frac{1}{1-\left(\frac{2}{z}\right)} \\ &= \sum_{n=0}^{\infty} \frac{1}{z^{n+1}} - \sum_{n=0}^{\infty} \frac{2^n}{z^{n+1}} = \sum_{n=0}^{\infty} \frac{1-2^n}{z^{n+1}} \\ &= \sum_{n=1}^{\infty} \frac{1-2^{n-1}}{z^n} \end{aligned}$$

$$f(z) = \frac{-1}{(z-1)(z-2)}$$

①  $D_1 \quad |z| < 1, \quad \left|\frac{z}{2}\right| < 1$




$$\frac{f(z)}{z^{n+1}} = \frac{-1}{(z-1)(z-2)z^{n+1}}$$

$$f(z) = \frac{1}{z-1} - \frac{1}{z-2} = \frac{-1}{1-z} + \frac{1}{1-\frac{z}{2}}$$

$$= -\sum_{n=0}^{\infty} z^n + \sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} = \sum_{n=0}^{\infty} (2^{-n-1} - 1)z^n \quad |z| < 1$$

$$a_n = \frac{f(z)}{z^{n+1}} = \frac{1}{(z-1)(z-2)z^{n+1}} \quad \frac{1}{z-1} - \frac{1}{z-2}$$

$$a_n = \sum_{k=1}^M \operatorname{Res} \left( \frac{f(z)}{(z-z_k)^{n+1}}, z_k \right) = \operatorname{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 0 \right)$$



$$a_n = \sum_{k=1}^M \operatorname{Res} \left( \frac{f(z)}{(z-z_k)^{n+1}}, z_k \right) = \operatorname{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 0 \right)$$

$n \geq 0$  then the pole  $z=0$

$$\frac{1}{(n-1)!} \lim_{z \rightarrow z_0} \frac{d^{n-1}}{dz^{n-1}} (z-z_0)^n f(z) \quad (\text{order } n)$$

$$\frac{d}{dz} ((z-1)^{-1} - (z-2)^{-1}) = (-1) ((z-1)^{-2} - (z-2)^{-2})$$

$$\frac{d^2}{dz^2} ((z-1)^{-1} - (z-2)^{-1}) = (-1)(-2) ((z-1)^{-3} - (z-2)^{-3})$$

$$\frac{d^3}{dz^3} ((z-1)^{-1} - (z-2)^{-1}) = (-1)(-2)(-3) ((z-1)^{-4} - (z-2)^{-4})$$

$$\frac{d^n}{dz^n} ((z-1)^{-1} - (z-2)^{-1}) = (-1)^n n! ((z-1)^{-n-1} - (z-2)^{-n-1})$$


$$\begin{aligned} \frac{1}{n!} \lim_{z \rightarrow 0} \frac{d^n}{dz^n} ((z-1)^{-1} - (z-2)^{-1}) &= (-1)^n \lim_{z \rightarrow 0} ((z-1)^{-n-1} - (z-2)^{-n-1}) \\ &= (-1)^n ((-1)^{-n-1} - (-2)^{-n-1}) \\ &= -1 + 2^{-n-1} \end{aligned}$$

$$a_n = -1 + 2^{-n-1} \quad (n \geq 0)$$

$$f(z) = \sum_{n=-n_1}^{\infty} a_n z^n = \sum_{n=0}^{\infty} (2^{-n-1} - 1) z^n$$


$$f(z) = \frac{-1}{(z-1)(z-2)}$$

$$\textcircled{2} D_2 \quad 1 < |z| < 2 \Rightarrow \left| \frac{1}{z} \right| < 1, \quad \left| \frac{z}{2} \right| < 1$$



$$\begin{aligned} f(z) &= \frac{1}{z-1} - \frac{1}{z-2} = \frac{1}{z} \cdot \frac{1}{1-\left(\frac{1}{z}\right)} + \frac{1}{2} \frac{1}{1-\left(\frac{z}{2}\right)} \\ &= \sum_{n=0}^{\infty} \frac{1}{z^{n+1}} + \sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} \\ &= \sum_{n=1}^{\infty} \frac{1}{z^n} + \sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}} \end{aligned}$$

$$\begin{aligned} a_n &= \sum_{k=1}^M \text{Res} \left( \frac{f(z)}{(z-z_k)^{n+1}}, z_k \right) = \text{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 0 \right) \\ &\quad + \text{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 1 \right) \end{aligned}$$



$$a_n = \sum_{k=1}^M \operatorname{Res} \left( \frac{f(z)}{(z-z_k)^{n+1}}, z_k \right) = \operatorname{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 0 \right) + \operatorname{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 1 \right)$$

$$\frac{1}{(n-1)!} \lim_{z \rightarrow z_0} \frac{d^{n-1}}{dz^{n-1}} (z-z_0)^n f(z) \quad (\text{order } n)$$

$$\begin{aligned} \frac{1}{n!} \lim_{z \rightarrow 0} \frac{d^n}{dz^n} ((z-1)^{-1} - (z-2)^{-1}) &= (-1)^n \lim_{z \rightarrow 0} ((z-1)^{-n-1} - (z-2)^{-n-1}) \\ &= (-1)^n ((-1)^{-n-1} - (-2)^{-n-1}) \\ &= -1 + 2^{-n-1} \end{aligned}$$

$$\operatorname{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 0 \right) = -1 + 2^{-n-1} \quad (n \geq 0)$$

$$\operatorname{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 1 \right) = \lim_{z \rightarrow 1} (z-1) \frac{-1}{(z-1)(z-2)z^{n+1}} = 1$$


$n=-3$	$n=-2$	$n=-1$	$n=0$	$n=1$	$n=2$	
0	0	0	$-1+2^{-1}$	$-1+2^{-2}$	$-1+2^{-3}$	$\operatorname{Res} \left( \frac{f(z)}{z^{n+1}}, 0 \right)$
1	1	1	1	1	1	$\operatorname{Res} \left( \frac{f(z)}{z^{n+1}}, 1 \right)$
1	1	1	$2^{-1}$	$2^{-2}$	$2^{-3}$	

$$\begin{cases} a_n = 2^{-n-1} & n \geq 0 \\ a_n = 1 & n < 0 \end{cases} \quad \begin{cases} 2^{-n-1} z^n \\ z^{-n} \end{cases}$$

$$f(z) = \sum_{n=1}^{\infty} \frac{1}{z^n} + \sum_{n=0}^{\infty} \frac{z^n}{2^{n+1}}$$


$$f(z) = \frac{-1}{(z-1)(z-2)}$$

$$\textcircled{3} \quad D_3 \quad 2 < |z| \quad \left| \frac{2}{z} \right| < 1 \quad \left| \frac{1}{z} \right| < 1$$



$$\begin{aligned} f(z) &= \frac{1}{z-1} - \frac{1}{z-2} = \frac{1}{z} \frac{1}{1-(\frac{1}{z})} - \frac{1}{z} \frac{1}{1-(\frac{2}{z})} \\ &= \sum_{n=0}^{\infty} \frac{1}{z^{n+1}} - \sum_{n=0}^{\infty} \frac{2^n}{z^{n+1}} = \sum_{n=0}^{\infty} \frac{1-2^n}{z^{n+1}} \\ &= \sum_{n=1}^{\infty} \frac{1-2^{n-1}}{z^n} \end{aligned}$$

$$\begin{aligned} a_n &= \sum_{k=1}^M \text{Res} \left( \frac{f(z)}{(z-z_k)^{n+1}}, z_k \right) = \text{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 0 \right) \\ &\quad + \text{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 1 \right) \\ &\quad + \text{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 2 \right) \end{aligned}$$





$$\text{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 0 \right) = -1 + 2^{-n+1} \quad (n \geq 0)$$

$$\text{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 1 \right) = \lim_{z \rightarrow 1} (z-1) \frac{-1}{(z-1)(z-2)z^{n+1}} = 1$$


$$\text{Res} \left( \frac{-1}{(z-1)(z-2)z^{n+1}}, 2 \right) = \lim_{z \rightarrow 2} (z-2) \frac{-1}{(z-1)(z-2)z^{n+1}} = -\frac{1}{2^{n+1}}$$

$n=-3$	$n=-2$	$n=-1$	$n=0$	$n=1$	$n=2$	
0	0	0	$-1+2^1$	$-1+2^2$	$-1+2^3$	$\text{Res} \left( \frac{f(z)}{z^{n+1}}, 0 \right)$
1	1	1	1	1	1	$\text{Res} \left( \frac{f(z)}{z^{n+1}}, 1 \right)$
$-2^2$	$-2$	$-1$	$-2^1$	$-2^2$	$-2^3$	$\text{Res} \left( \frac{f(z)}{z^{n+1}}, 2 \right)$
$1-2^2$	$1-2$	0	0	0	0	

$$a_n = 1 - 2^{-n+1} \quad n < 0 = \sum_{n=1}^{\infty} \frac{1-2^{n+1}}{z^n}$$

$$f(z) = \sum_{n=-1}^{-\infty} (1-2^{-n+1}) z^n = \sum_{n=1}^{\infty} \frac{1-2^{n+1}}{z^n}$$

$$f(z) = \frac{-1}{(z-1)(z-2)}$$



$$x[n]$$

$$= \frac{1}{2\pi i} \int_C \boxed{X(z) z^{n-1}} dz$$

$$= \sum_{j=1}^k \text{Res}(\boxed{X(z) z^{n-1}}, z_j)$$


$$X(z) = \frac{-1}{(z-1)(z-2)}$$

$$X(z) z^{n-1} = \frac{-1}{(z-1)(z-2)} z^{n-1}$$

$$\text{Res}(\boxed{X(z) z^{n-1}}, 1) = (z-2) \frac{-1}{(z-1)(z-2)} z^{n-1} \Big|_{z=1} = 1$$

$$\text{Res}(\boxed{X(z) z^{n-1}}, 2) = (z-1) \frac{-1}{(z-1)(z-2)} z^{n-1} \Big|_{z=2} = -2^{n-1}$$

$$x[n] = 1 - 2^{n-1}$$



ROC (Region of Convergence)

$$|z| > 2 \Rightarrow \frac{2}{|z|} < 1$$

$$\left(\frac{2}{z}\right)^0 + \left(\frac{2}{z}\right)^1 + \left(\frac{2}{z}\right)^2 + \dots \longrightarrow \frac{1}{1 - \frac{2}{z}}$$

Converge

$$|z| > 2 \Rightarrow \frac{1}{|z|} < 1$$

$$\left(\frac{1}{z}\right)^0 + \left(\frac{1}{z}\right)^1 + \left(\frac{1}{z}\right)^2 + \dots \longrightarrow \frac{1}{1 - \frac{1}{z}}$$

Converge

$$\begin{aligned} f(z) &= \frac{1}{z-1} - \frac{1}{z-2} = \frac{1}{z} \frac{1}{1 - (\frac{1}{z})} - \frac{1}{z} \frac{1}{1 - (\frac{2}{z})} \\ &= \sum_{n=0}^{\infty} \frac{1}{z^{n+1}} - \sum_{n=0}^{\infty} \frac{2^n}{z^{n+1}} = \sum_{n=0}^{\infty} \frac{1 - 2^n}{z^{n+1}} \\ &= \sum_{n=1}^{\infty} \frac{1 - 2^{n-1}}{z^n} \end{aligned}$$

$$\left(\frac{1}{z}\right) + \left(\frac{1}{z}\right)^2 + \left(\frac{1}{z}\right)^3 + \dots + \frac{1}{2} \left\{ \left(\frac{2}{z}\right) + \left(\frac{2}{z}\right)^2 + \left(\frac{2}{z}\right)^3 + \dots \right\} \longrightarrow \frac{1}{z-1} - \frac{1}{z-2} = \frac{-1}{(z-1)(z-2)}$$

Converge

$$(1-2^0)z^{-1} + (1-2^1)z^{-2} + (1-2^2)z^{-3} + \dots \longrightarrow \frac{-1}{(z-1)(z-2)} \quad (|z| > 2)$$


Converge

$$x[n] = 1 - 2^n \quad \longleftrightarrow \quad X(z) = \frac{-1}{(z-1)(z-2)} \quad (|z| > 2)$$

①

$$f(z) = \frac{12}{z(2-z)(1+z)} = \frac{4}{z} \left( \frac{1}{1+z} + \frac{1}{2-z} \right)$$

pole:  $z=0$ ,  $z=2$ ,  $z=-1$



2

$$0 < |z| < 1$$


$$f(z) = -3 + 9z/2 - 15z^2/4 + 33z^3/8 + \dots + 6/z$$

$$|z| > 2$$

$$\frac{1}{1+z} = \frac{1}{z} \frac{1}{(1+z^{-1})} \quad \frac{1}{2+z} = -\frac{1}{z} \frac{1}{1-2z^{-1}}$$


$$f(z) = -(12/z^3) (1 + 1/z + 3/z^2 + 5/z^3 + 11/z^4 + \dots)$$

3




$$0 < |z| < 1$$

$$f(z) = -3 + 9z/2 - 15z^2/4 + 33z^3/8 + \dots + 6/z$$



5



$$|z| > 2$$



$$\frac{1}{1+z} = \frac{1}{z} \frac{1}{(1+z^{-1})}$$

$$\frac{1}{2+z} = -\frac{1}{z} \frac{1}{1-2z^{-1}}$$



causal  $x[n]=0$  ( $n < 0$ )

anti-causal  $x[n]=0$  ( $n > 0$ )




ROC: outside a circle

ROC: inside a circle

bi-causal  $x[n]$

Overlapped ROC



$$f(z) = \sum_{n=-\infty}^{\infty} a_n^{(m)} (z - z_m)^n$$

$$f(z) = \sum_{n=-\infty}^{\infty} a_n z^n \quad z_m = 0 \quad a_n^{(0)} \rightarrow a_n$$

## Laurent Series at $z=0$

$$f(z) = \dots + a_{-2} z^{-2} + a_{-1} z^{-1} + a_0 z^0 + a_1 z^1 + a_2 z^2 + a_3 z^3 + \dots$$

## z-transform

Bi-causal

$$X(z) = \dots + x[-2] z^2 + x[-1] z^1 + x[0] z^0 + x[1] z^{-1} + x[2] z^{-2} + x[3] z^{-3} + \dots$$

Causal

$$X(z) = x[0] z^0 + x[1] z^{-1} + x[2] z^{-2} + x[3] z^{-3} + \dots$$

Anti-causal

$$X(z) = \dots + x[-1] z^2 + x[-1] z^1 + x[0] z^0$$

$$a_n \leftrightarrow x[-n]$$

$$a_{-n} \leftrightarrow x[n]$$

$$f(z) = \sum_{n=n_1}^{\infty} a_n^{\{m\}} (z - z_m)^n$$

$$a_n^{\{m\}} = \frac{1}{2\pi i} \oint_C \frac{f(z')}{(z' - z_m)^{n+1}} dz'$$

$$= \sum_k \operatorname{Res} \left( \frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

Analytic at  $z_m$

$$n_1 \geq 0$$

Taylor Series

$$\text{general } n_1, \quad z_m = 0$$

MacLaurin Series

Singular at  $z_m$

$$\text{general } n_1$$

Laurent Series

$$\text{general } n_1, \quad z_m = 0$$

z-Transform

$$f(z) = \sum_{n=-\infty}^{\infty} a_n^{(m)} (z - z_m)^n$$

$$\begin{aligned} a_n^{(m)} &= \frac{1}{2\pi i} \oint_c \frac{f(z')}{(z' - z_m)^{n+1}} dz' \\ &= \sum_k \operatorname{Res} \left( \frac{f(z)}{(z - z_m)^{n+1}}, z_k \right) \end{aligned}$$

$$z_m = 0$$

$$a_{-n}^{(0)} = h(n)$$

$$n \rightarrow -n$$

$$H(z) = \sum_{n=-\infty}^{\infty} h(-n) z^n$$

$$\begin{aligned} h(n) &= \frac{1}{2\pi i} \oint_c \frac{H(z')}{z'^{n+1}} dz' \\ &= \sum_k \operatorname{Res} \left( \frac{H(z)}{z^{n+1}}, z_k \right) \end{aligned}$$

$$H(z) = \sum_{n=-\infty}^{\infty} h(n) z^{-n}$$

$$\begin{aligned} h(n) &= \frac{1}{2\pi i} \oint_c H(z') z'^{n-1} dz' \\ &= \sum_k \operatorname{Res} (H(z) z^{n-1}, z_k) \end{aligned}$$

$C$  is in the same region of analyticity of  $f(z)$   
typically a circle centered on  $z_m$

$z_k$  within  $C$  : singularities of  $\frac{f(z)}{(z-z_k)^{n+1}}$

$C$  is in the same region of analyticity of  $H(z)$   
typically a circle centered on  $z_m$

generally a circle centered on the origin  
may enclose any or all singularities of  $H(z)$   
often the unit circle

$z_k$  within  $C$  : singularities of  $H(z)z^{n-1}$

$$H(z) = \sum_{n=-\infty}^{\infty} h(n) z^{-n} \quad z \in \text{R.O.C.}$$

$$h(n) = \frac{1}{2\pi i} \oint_C H(z') z'^{n-1} dz' \quad C \text{ in R.O.C.}$$
$$= \sum_k \text{Res}(H(z) z^{n-1}, z_k)$$


- ① a power series representation of a function  $f(z)$  of a complex variable  $z$
- ② a transform  $H(z)$  of a sequence of 1


$$X(z) = \frac{z}{z - \frac{1}{2}} \quad \text{pole } z_0 = \frac{1}{2}$$

$$\begin{aligned} x[n] &= \text{Res} \left( X(z) z^{n-1}, z_0 \right) = \text{Res} \left( \frac{z}{z - \frac{1}{2}} z^{n-1}, \frac{1}{2} \right) \\ &= \text{Res} \left( \frac{z^n}{z - \frac{1}{2}}, \frac{1}{2} \right) = \lim_{z \rightarrow \frac{1}{2}} (z - \frac{1}{2}) \frac{z^n}{z - \frac{1}{2}} = \left( \frac{1}{2} \right)^n \end{aligned}$$

$$x[n] = \frac{1}{2^n} \quad n \geq 0$$

$$\begin{aligned} \left( \frac{1}{2} \right)^0 z^0 + \left( \frac{1}{2} \right)^1 z^{-1} + \left( \frac{1}{2} \right)^2 z^{-2} + \left( \frac{1}{2} \right)^3 z^{-3} + \dots &= \frac{1}{1 - \left( \frac{1}{2} z^{-1} \right)} \\ &= \frac{z}{z - \frac{1}{2}} \end{aligned}$$





$$x_{s,c}(t) = \sum_{n=-\infty}^{+\infty} x(n) \delta_c(t - n\Delta t)$$

$$\begin{aligned} X_{s,c}(s) &= \mathcal{L}\{x_{s,c}(t)\} = \int_{-\infty}^{\infty} \left[ \sum_{n=-\infty}^{+\infty} x(n) \delta_c(t - n\Delta t) \right] e^{-st} dt \\ &= \sum_{n=-\infty}^{+\infty} x(n) \int_{-\infty}^{\infty} \delta_c(t - n\Delta t) e^{-st} dt \\ &= \sum_{n=-\infty}^{+\infty} x(n) e^{-sn\Delta t} \quad e^{s\Delta t} \triangleq z \end{aligned}$$

$$X_{s,c}(s) = \sum_{n=-\infty}^{+\infty} x(n) z^{-n} \Big|_{z=e^{s\Delta t}}$$

$$X_{s,c}(s) = X(z) \Big|_{z=e^{s\Delta t}}$$



$$X_{s,c}(s) = \mathcal{L}\{x_{s,c}(t)\} = X(z) \Big|_{z=e^{s\Delta t}}$$

$x_{s,c}(t)$  an impulse train

whose coefficients are given by  $x[n] = x_c(n\Delta t)$

z-transform : a special Laurent series

$$z_m = 0$$

$$a_{-n}^{\{0\}} = h(n)$$

$$n \rightarrow -n$$

$$f(z) = \sum_{n=n_1}^{\infty} a_n^{\{m\}} (z - z_m)^n$$

$$a_n^{\{m\}} = \frac{1}{2\pi i} \oint_c \frac{f(z')}{(z' - z_m)^{n+1}} dz'$$

$$= \sum_k \operatorname{Res} \left( \frac{f(z)}{(z - z_m)^{n+1}}, z_k \right)$$

Time Reversal  $\leftarrow$  Laplace Transform

the transform functions

$$X(s) = \int \text{over negative powers } e^{-st} \quad \text{for } t > 0$$


$$X(z) = \int \text{over negative powers } z^{-n} \quad \text{for } n > 0$$

the time expansion functions

$$x(t) = \int \text{over negative powers } e^{-st} \quad \text{for } t > 0$$

$$x[n] = \int \text{over negative powers } z^{-n} \quad \text{for } n > 0$$

Time Reversal  $\leftarrow$   $z^{-1}$ : unit delay, char eq (modes in  $z^k$ )



Stable system :  $h[n]$  must be absolutely summable

$$|e^{j\omega n}| = 1$$

$$|z^n| \quad z = 1$$

$$\infty > M_h > \sum_{n=-\infty}^{\infty} |h[n]| \quad \text{absolutely summable}$$

$$= \sum_{n=-\infty}^{+\infty} |h[n] e^{-j\omega n}|$$

$$\geq \left| \sum_{n=-\infty}^{+\infty} h[n] e^{-j\omega n} \right|$$

$$= \left| H(z) \Big|_{z=e^{j\omega}} \right|$$

$$\infty > \left| H(z) \Big|_{z=e^{j\omega}} \right|$$

a stable system,

$H(z)$  must converge on the unit circle  $|z|=1$

ROC (Region of Convergence) must include the unit circle

regardless of causality of  $h[n]$

$$H(z) \Big|_{|z|=1} = H(e^{j\hat{\omega}}) \quad \text{DTFT of } h[n]$$

discrete    all stable sequence must have convergent DTFTs

continuous    all stable signal must have convergent CTFTs

$$C \leftarrow \text{unit circle} \quad z = e^{j\hat{\omega}}$$

$ZT^{-1}$        $DTFT^{-1}$       identical formulas

$h[n]$  causal

$$H(z) = \sum_{n=-\infty}^{+\infty} h[n] z^{-n} = \sum_{n=0}^{+\infty} h[n] z^{-n} \quad n \in [0, \infty)$$

for finite values of  $n$ ,

each term must be finite as long as  $z \neq 0$

For the sum to converge,

$h[n] z^{-n}$  must vanish as  $n \rightarrow \infty$

$$|z| > r_h \quad z_h = r_h e^{j\theta}$$

$z_h^n$  is the largest magnitude

geometrically increasing component

$n^m z^n$  : the most general term

for impulse responses

$n \rightarrow \infty$   $z^n$  dominant over  $n^m$  for finite  $m$

Geometric components — as poles

$$\sum \{ z_0^n u[n] \} = \frac{1}{1 - (\frac{z_0}{z})} = \frac{z}{z - z_0}$$

ROC of a causal sequence  $h[n]$

outside the radius of the largest magnitude pole of  $H(z)$

ROC of a causal signal  $h(t)$

to the right of the rightmost pole of  $H_c(s)$

if  $h[n]$  is a stable, causal sequence,

the unit circle must be included in the ROC

• Causal  $h[n]$


ROC: outside of  
a circle

• Stable  $h[n]$

all poles inside  
the unit circle

ROC circle must be

smaller than the unit circle



⇒ all the geometric components of  $h[n]$  : modes  
must decay with increasing  $n$

all the poles of  $H(z)$  must be within the unit circle

all the poles of  $H_c(s)$  must be in the left half plane

• anti-causal  $h[n]$


ROC: inside of  
a circle

• Stable  $h[n]$

all poles outside  
the unit circle

ROC circle must be

larger than the unit circle



⇒ all the geometric components of  $h[n]$  : modes  
must decay with decreasing  $n$




• bi-causal  $h[n]$

$$h_c[n] + h_{ac}[n]$$

outside      inside

max mag < min mag

Overlapped ROC



• Stable  $h[n]$

all poles outside


the unit circle

ROC circle must include the unit circle


## • bi-causal $h[n]$

$$h[n] = h_c[n] + h_{ac}[n]$$

causal comp.                      anti-causal comp




outside a circle



inside a circle


$$\text{max mag} < \text{min mag}$$

Overlapped ROC



$$\text{max mag} > \text{min mag}$$

non-overlapping ROC



## • Stable $h[n]$

all poles outside the large circle  
inside the small circle

ROC circle must include the unit circle

only one annulus include the unit circle

only one stable sequence

# Existence of the z-Transform

$$X(z) = \sum_{n=0}^{\infty} x[n]z^{-n} = \sum_{n=0}^{\infty} \frac{x[n]}{z^n}$$

the existence of the z-transform is guaranteed if

$$|X(z)| \leq \sum_{n=0}^{\infty} \frac{|x[n]|}{|z|^n} < \infty \quad \text{for some } |z|$$

any signal  $x[n]$  that grows no faster than an exponential signal  $r_0^n$ , for some  $r_0$  satisfies the above condition

if  $|x[n]| \leq r_0^n$  for some  $r_0$

$$\text{then } |X(z)| \leq \sum_{n=0}^{\infty} \left(\frac{r_0}{|z|}\right)^n = \frac{1}{1 - \frac{r_0}{|z|}} \quad |z| > r_0$$

therefore  $X(z)$  exists for  $|z| > r_0$

Almost all practical signals satisfy this condition

$$|x[n]| \leq r_0^n \quad \text{for some } r_0$$

and z-transformable

Some signal models (e.g.  $r^{n^2}$ ) grows faster than

the exponential signal  $r_0^n$  (for any  $r_0$ )

and do not satisfy this condition

and are not z-transformable

Such signals are of little practical or theoretical interest

Even such signals over a finite interval are z-transformable

# Region of Convergence

Laplace Transform	$Ae^{\alpha t}u(t)$	$\alpha > 0$
z - Transform	$A\alpha^n u[n]$	$ \alpha  > 0$
DTFT (X)		

$$X(z) = A \sum_{n=-\infty}^{\infty} \alpha^n u[n] z^{-n} = A \sum_{n=0}^{\infty} \alpha^n z^{-n} = A \sum_{n=0}^{\infty} \left(\frac{\alpha}{z}\right)^n$$

converge  $\left|\frac{\alpha}{z}\right| < 1$   $|z| > |\alpha|$   
open exterior of  
a circle of radius  $|\alpha|$

the sum of a geometric series

$$X(z) = A \frac{1}{1 - \frac{\alpha}{z}} = \frac{A}{1 - \alpha z^{-1}} = A \frac{z}{z - \alpha} \quad |z| > |\alpha|$$

DTFT

$$X(j\hat{\omega}) = \sum_{n=-\infty}^{+\infty} x[n] e^{-j\hat{\omega}n}$$

# DTFT

DTFT of the unit sequence  $u[n]$

$$X(e^{-j\hat{\omega}n}) = \sum_{n=-\infty}^{+\infty} u[n] e^{-j\hat{\omega}n} = \sum_{n=0}^{\infty} e^{-j\hat{\omega}n}$$

not converge

$$\hat{\omega} = 0 \quad \sum_{n=0}^{\infty} 1^n \quad \text{diverge}$$

$$\hat{\omega} = \pi \quad \sum_{n=0}^{\infty} (-1)^n \quad \text{oscillates}$$

$$\hat{\omega} = \frac{\pi}{2} \quad \sum_{n=0}^{\infty} (j)^n$$

The DTFTs of some commonly used functions do not exist in the strict sense.


But even though the DTFT does not exist, the  $z$ -transform does exist.

$$X(z) = \sum_{n=-\infty}^{+\infty} u[n] z^{-n} = \sum_{n=0}^{\infty} z^{-n}$$

$$|z| > 1 \quad X(z) = \frac{z}{z-1} = \frac{1}{1-z^{-1}}$$

$$X(z) = \frac{z}{z-1} \quad \text{pole } z=1, \quad \text{zero } z=0$$

$$X(z) = \frac{1}{1-z^{-1}} \quad \text{useful when a system is synthesized from a } z\text{-domain transfer function}$$



$$f[n] = \frac{1}{2} \quad -\infty < n < \infty$$

$$g[n] = \begin{cases} \frac{1}{2} & n \geq 0 \\ -\frac{1}{2} & n < 0 \end{cases}$$

$$u[n] = f[n] + g[n]$$

$$\delta[n] = g[n] - g[n-1]$$

$$1 = G(e^{j\omega}) - e^{-j\omega} G(e^{j\omega})$$

$$G(e^{j\omega}) = \frac{1}{1 - e^{-j\omega}}$$

$$F(e^{j\omega}) = \pi \sum_{k=-\infty}^{+\infty} \delta(\omega - 2\pi k) \quad (\text{impulse train})$$

$$U(e^{j\omega}) = \frac{1}{1 - e^{-j\omega}} + \pi \sum_{k=-\infty}^{+\infty} \delta(\omega - 2\pi k)$$

# Discrete Time Exponential $r^n$

continuous time exponential  $e^{\lambda t}$

$$\begin{aligned}e^{\lambda t} &= r^t & (e^{\lambda})^t &= r^t \\e^{\lambda} &= r \\ \lambda &= \ln r\end{aligned}$$

$$\begin{aligned}e^{-0.03t} &= (0.9408)^t \\ 4^t &= e^{1.386t}\end{aligned}$$

continuous time analysis  $e^{\lambda t}$

discrete time analysis  $r^n$

$$\begin{aligned}e^{\lambda n} &= r^n & (e^{\lambda})^n &= r^n \\e^{\lambda} &= r \\ \lambda &= \ln r\end{aligned}$$

$e^{\lambda n}$

Exponentially grows if  $\text{Re } \lambda > 0$  ( $\lambda$  in RHP)

exponentially decays if  $\text{Re } \lambda < 0$  ( $\lambda$  in LHP)

Oscillates or constant if  $\text{Re } \lambda = 0$  ( $\lambda$  in imag axis)

the location of  $\lambda$  in the complex plain indicates whether

①  $e^{\lambda t}$  will grow exponentially

②  $e^{\lambda t}$  will decay exponentially

③  $e^{\lambda t}$  will oscillates with constant amplitude

constant signal : oscillation with zero frequency

$e^{j\Omega n}$   $\lambda = j\Omega$  imaginary axis

(constant amplitude oscillating signal)

$$e^{j\Omega n} = (e^{j\Omega})^n = \gamma^n \quad \gamma = e^{j\Omega} \quad |\gamma| = 1$$

$\lambda = j\Omega$  imaginary axis  $\rightarrow |\gamma| = 1$  unit circle

if  $\gamma$  lies on the unit circle,

$\gamma^n$  oscillates with constant amplitude

the imaginary axis in the  $\lambda$  plane

the unit circle in the  $\gamma$  plane



$e^{\lambda n}$      $\lambda = a + jb$  in the LHP ( $a < 0$ )  
exponentially decaying

$$r = e^{\lambda} = e^{a+jb} = e^a e^{jb}$$

$$|r| = |e^{\lambda}| = |e^a \cdot e^{jb}| = |e^a| = e^a$$

$|r| = e^a < 1$     inside the Unit circle

$r^n$  : exponentially decaying

$|r| = e^a > 1$     outside the Unit circle

$r^n$  : exponentially growing

$\lambda$ -plane

the imaginary axis

the LHP

the RHP



$z$ -plane

the unit circle

inside of the unit circle

outside of the unit circle



